

Massive 3D Gravity Big-Bounce

H.L.C.Louzada*, U.Camara dS^{*1} and G.M.Sotkov^{*2}

* Departamento de Física - CCE
Universidade Federal do Espírito Santo
29075-900, Vitoria - ES, Brazil

ABSTRACT

The properties of an extension of the New Massive 3D Gravity by scalar matter with Higgs-like self-interaction are investigated. Its perturbative unitarity consistency is verified for a family of cosmological Bounce solutions found by the superpotential method. They correspond to the lower bound $\lambda = -1$ of the BHT unitarity window and describe eternally accelerated 3D Universe between two initial/final stable dS_3 vacua states.

¹e-mail: ulyssescamara@gmail.com

²e-mail: sotkov@cce.ufes.br, gsotkov@yahoo.com.br

The perturbative quantization of the Einstein gravity is known to introduce at one-loop order in the gravitational coupling $\kappa^2 = 16\pi G$ “quadratic” counterterms $\mathcal{L}_{ct} = \sqrt{-g}(\alpha R^2 + \beta R^{\mu\nu} R_{\mu\nu} + \dots)$ which cure its UV divergencies but violate unitarity (for $d \geq 4$) [1], [2]. In three dimensions the problem of perturbative consistency of quantum gravity model including specific “high derivatives” terms:

$$S_{NMG}^{eff}(g_{\mu\nu}, \lambda_0, \Lambda) = \frac{1}{\kappa^2} \int dx^3 \sqrt{-g} [\epsilon R + 64\lambda_0 K - 2\Lambda] \quad , \quad K = R_{\mu\nu} R^{\mu\nu} - \frac{3}{8} R^2 \quad (1)$$

called New Massive Gravity (NMG) was recently reexamined by Bergshoeff, Hohm and Townsend (BHT) [3], [4]. It turns out that it is super-renormalizable and unitary (ghost free) under certain restrictions on the values of the cosmological constant Λ and of the new scale (mass) parameter λ_0 for the both choices $\epsilon = \pm 1$ of the “right” or “wrong” sign of the R -term [4], [5], [6]. The BHT - model (1) has two propagating degrees of freedom and unlike 3D Einstein gravity (*i.e.* $\lambda_0 = 0$ case) it admits interesting (vacuum) solutions - black holes, gravitational waves, etc. [7], [8]. Its Newtonian limit for $\epsilon = -1$ reproduces an attractive gravitational force [9].

The New Massive Gravity (1) can be realized perturbatively as a massive spin two quantum field $\hat{h}_{\mu\nu}$ ($\hat{g}_{\mu\nu} = \bar{g}_{\mu\nu} + \kappa \hat{h}_{\mu\nu}$) - theory with unusual kinetic term and specific self-interactions in external constant curvature background $\bar{g}_{\mu\nu}$. Such a procedure provides well defined rules for studying the quantum graviton ($\hat{h}_{\mu\nu}$) properties and eventually it might shed some light on the nature of the “quantum 3D geometry” behind it. Thus we have an example of consistent (3D) quantum gravity model that shares many of the (desired) properties of 4D gravity. It is therefore natural to address the question about the perturbative quantum consistency of the New Massive Gravity coupled to matter fields. Since we are interested to study the Cosmological consequences of such BHT-matter model, the simplest and most representative candidate of matter QFT is the one of scalar field $\sigma(x^\mu)$ with Higgs - like self-interaction:

$$\mathcal{L}_{matt}(\sigma, g_{\mu\nu} | \gamma, m_\sigma) = \sqrt{-g} \left(-\frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma - V(\sigma) \right) \quad , \quad V(\sigma) = \frac{\gamma}{4} \left(\sigma^2 - \frac{m_\sigma^2}{2\gamma} \right)^2 - 2\Lambda \quad (2)$$

The parameters γ , m_σ and Λ above denote the renormalized coupling constant γ , “Higgs mass” m_σ and the vacuum energy Λ and they are assumed to depend on the “graviton mass” scale $\lambda_0 = \frac{1}{64m^2}$. Since in four dimensions (in curved space) this model is known to be renormalizable [10], [11] one expects that it is power - counting super-renormalizable in three dimensions. Note that the relevant (lowest order in γ) counter-terms to \mathcal{L}_{matt} are again in the form $\tilde{\alpha} R^2 + \tilde{\beta} R_{\mu\nu} R^{\mu\nu}$ [10], *i.e.* they do coincide with the gravitational ones (in the case of minimal coupling $\xi = 0$). We further assume that the corresponding (vacuum) “semiclassical” equations [10], [12]:

$$\bar{R}_{\mu\nu} - \frac{1}{2} \bar{g}_{\mu\nu} \bar{R} + \Lambda \bar{g}_{\mu\nu} = \frac{1}{2} \kappa^2 \langle vac | \hat{T}_{\mu\nu}(h_{\mu\nu}, \sigma) | vac \rangle \quad (3)$$

for this BHT - matter model (to lowest order in the both couplings κ^2 and γ) with the UV counter-terms added and when adiabatic vacuum states $|vac \rangle = |g_{\mu\nu}^{vac}, \sigma_\pm \rangle$, ($V'(\sigma_\pm) = 0$) exist, are represented by the *classical* equations obtained from the following “effective” Matter-Gravity action:

$$S_{NMG M}^{eff} = \frac{1}{\kappa^2} \int dx^3 \sqrt{-g} \left[\epsilon R + 64\lambda_0 K - \frac{1}{2} |\nabla \sigma|^2 - V(\sigma) \right] \quad (4)$$

The BHT - cosmological constant Λ is now included in the potential (2), *i.e.* $V(\sigma_\pm^2 = \frac{m_\sigma^2}{2\gamma}) = -2\Lambda$ for the case $\epsilon \lambda_0 < 0$. In order to describe in a compact form the corrections (*i.e.* backreaction)

to the classical background $\bar{g}_{\mu\nu}$ ($\bar{R}_{\mu\nu} = R_{\mu\nu}(\bar{g}_{\mu\nu})$ etc.) due to the quantum fluctuations of both $\hat{h}_{\mu\nu}$ and $\hat{\sigma}$, as usually (see ref. [12]) all the higher derivatives contributions are included in the semiclassical “source” in the r.h.s. of eqs.(3).

The present paper is devoted to the investigation of the properties of homogeneous and isotropic spatially flat $k = 0$ $(-, +, +)$ 3D Universe :

$$ds_3^2 = -dt^2 + a_{eff}^2(t)(dx^2 + dy^2) \quad (5)$$

described by the matter extension (4) of BHT-model (1). Our *problem* is therefore to find analytic solutions of the corresponding 3D “Massive Cosmological” equations³ :

$$\begin{aligned} \ddot{\sigma} + \dot{\sigma}\dot{\varphi} + V'(\sigma) &= 0 \\ \ddot{\varphi}(1 + 8\epsilon\lambda_0\dot{\varphi}^2) + \frac{1}{2}\dot{\varphi}^2(1 + 4\epsilon\lambda_0\dot{\varphi}^2) + \epsilon\left(\frac{1}{2}\dot{\sigma}^2 - V(\sigma)\right) &= 0 \\ \epsilon(\dot{\sigma}^2 + 2V(\sigma)) - \dot{\varphi}^2(1 + 4\epsilon\lambda_0\dot{\varphi}^2) &= 0 \end{aligned} \quad (6)$$

derived from the action (4) above. We are further interested in studying specific features of these solutions as for example: the effects of the matter on the BHT - unitarity conditions (u.c.’s), the conditions of absence of initial/final singularities, how many different periods of acceleration and/or deceleration they describe etc.

Observe that eqs. (6) for the scale factor $a(t) = e^{\frac{\varphi(t)}{2}}$ and for the scalar field $\sigma(t)$ are of second order although the action (4) and the eqs. (3) include (up to) fourth order time derivatives of $g_{\mu\nu}$. The special BHT unitarity motivated choice of the relative coefficients $\beta = -\frac{3}{8}\alpha = 64\lambda_0$ (that fix the form of the K-term for $D = 3$) together with the FRW ansatz (5) are responsible for the cancellation of all the terms involving $\varphi^{(III)}$ and $\varphi^{(IV)}$ derivatives and they cure another disease of all “higher derivatives” gravities. The family of solutions we are considering does have well defined initial values Cauchy problem (*i.e.* “causality”) for *all the values* of the parameters $\epsilon, \lambda_0, \gamma, m_\sigma^2$, *i.e.* independently of whether or not the BHT u.c.’s [4] are satisfied. The construction of all the solutions of this second order system of nonlinear eqs. (6) is however rather difficult problem, although numerical methods are indeed available and largely used for this type of dynamical systems. One has to be able to further analyze how their properties depend on the initial conditions $(\sigma_0, \dot{\sigma}_0, \varphi_0, \dot{\varphi}_0)$ and on the values of the parameters $\epsilon, \lambda_0, \gamma, m_\sigma$ as well.

At this place we take a different route known as “superpotential” method, which provides very special analytic (non-perturbative in γ and λ_0) solutions of eqs. (6). It is widely used in the construction of domain walls (DW) solutions of different (super) gravity models in arbitrary dimensions (see for example refs. [13], [14] for $\lambda_0 = 0$ case). It consists in the introduction of an auxiliary function $W(\sigma)$ (called superpotential) such that:

$$\begin{aligned} V(\sigma) &= 2\epsilon W^2(1 + 16\epsilon\lambda_0 W^2) - 2(W')^2(1 + 32\epsilon\lambda_0 W^2)^2 \\ \dot{\varphi} &= -2\epsilon W(\sigma) \\ \dot{\sigma} &= 2W'(\sigma)(1 + 32\epsilon\lambda_0 W^2) \end{aligned} \quad (7)$$

For $\lambda_0 \neq 0$ it represents an adapted version of the Low-Zee superpotential [15] introduced in the context of DW’s solutions of $d = 5$ Gauss-Bonnet improved gravity. The statement is that for each function $W(\sigma)$ that reproduces our matter potential (2) the solutions of the first order system (7) are solutions of the eqs. (6) as well.

³in what follows we use the convention $c = \kappa = \hbar = 1$

We next consider a particular linear superpotential $W(\sigma) = B\sigma$, which leads to a family of σ^4 -like (Z_2 symmetric $\sigma \rightarrow -\sigma$) potentials (2) with

$$\begin{aligned} \gamma &= 128\lambda_0 B^4(1 - 64\lambda_0 B^2) \quad , \quad m_\sigma^2 = -8\epsilon B^2(1 - 64\lambda_0 B^2) \\ \sigma_\pm^2 &= -\frac{1}{32\epsilon\lambda_0 B^2}, \quad \Lambda = \frac{1}{64\lambda_0} = m^2 \quad , \quad \bar{m}_\sigma^2 = V''(0) = -\frac{1}{2}V''(\sigma_\pm) = -\frac{m_\sigma^2}{2} \end{aligned} \quad (8)$$

Hence depending on the $\epsilon\lambda_0$ sign (and whether $B^2 < m^2$ or $B^2 > m^2$) we have normal or inverted double-well potentials for $\epsilon\lambda_0 < 0$ or the standard σ^4 -potentials for $\epsilon\lambda_0 > 0$. The fact that the form of the matter potentials (encoded here in $\gamma = \gamma(\lambda_0)$ and $m_\sigma^2 = m_\sigma^2(\lambda_0)$) does depend on the gravitational mass scale $m^2 = \frac{1}{64\lambda_0}$ is an intrinsic property of the superpotential method (see eqs. (7)) when applied for models which actions contain K -type terms (1).

Note that for the particular choice of $W(\sigma)$ we consider, the parameters $\{\gamma, m_\sigma^2, m^2\}$ are not independent. As one can see from eqs.(8), they describe a special surface

$$m_\sigma^6 = 32\gamma m^2(m_\sigma^2 + 4\epsilon\gamma) \quad , \quad 1 + \epsilon\frac{m_\sigma^2}{2m^2} > 0 \quad (9)$$

in the parameter space of the BHT - matter model (4). Taking the effective masses m_σ^2 and m^2 as independent variables we realize that the coupling constant γ (or equivalently the ‘‘superpotential’’ parameter B^2) are given by:

$$\gamma_\pm = -\epsilon\frac{m_\sigma^2}{8}\left(1 \pm \sqrt{1 + \epsilon\frac{m_\sigma^2}{2m^2}}\right) \equiv -\epsilon\frac{m_\sigma^2 B_\pm^2}{4m^2} \quad (10)$$

i.e. they take two different values γ_\pm (and B_\pm^2) when $\frac{m_\sigma^2}{\epsilon m^2} < 0$ and only one γ_- (or B_-^2) in the case $\frac{m_\sigma^2}{\epsilon m^2} > 0$. Together with the signs of m_σ^2 and m^2 they characterize the particular shapes of the potential $V(\sigma|\gamma_\pm, m_\sigma^2)$ for which we are able to solve analytically eqs. (6) by the superpotential method. We will consider mainly the case $\bar{m}_\sigma^2\gamma < 0$ (*i.e.* $\epsilon\lambda_0 < 0$) which describes:

- (a) $\epsilon = -1, \lambda_0 > 0$ BHT - model with double-well (Higgs) potential $V_-(\sigma)$.
- (b) $\epsilon = 1, \lambda_0 < 0$ BHT - model with *inverted* double well potential $V_+(\sigma)$.

In both cases we have $\sigma_\pm^2 > 0$ and therefore $\sigma_\pm = \pm\sqrt{\frac{m_\sigma^2}{2\gamma}}$ are the minima ($\epsilon = -1$) or the maxima ($\epsilon = 1$) of the potentials V_\pm . For these ‘‘vacuum’’ values σ_\pm of the field $\sigma(t)$, the BHT - matter model (4) reduces to the pure BHT - gravity (1) for a particular value of the cosmological constant⁴ $\Lambda = m^2 = \frac{1}{64\lambda_0}$, *i.e.* we have $\Lambda < 0$ for $\lambda_0 < 0$ ($\epsilon = 1$) and $\Lambda > 0$ for $\lambda_0 > 0$ ($\epsilon = -1$). The effective cosmological constant Λ_{eff} is defined by the values of 3D curvature $R^{(3)}$ on the corresponding vacuum solutions $(g_{\mu\nu}^{vac, \pm}, \sigma_\pm)$:

$$R^{(3)}(g_{\mu\nu}^{vac}, \sigma_\pm) = 2\ddot{\varphi} + \frac{3}{2}\dot{\varphi}^2 \equiv -8\epsilon(W')^2(1 + 32\lambda_0\epsilon W^2) + 6W^2 \quad (11)$$

which gives $R^{(3)}(g_{\mu\nu}^{vac}, \sigma_\pm) = -\frac{3}{16\epsilon\lambda_0} \equiv 6\Lambda_{eff}^\pm$. Therefore in both cases when $\epsilon\lambda_0 < 0$, the $\Lambda_{eff}^\pm = 2|m^2| > 0$ is positive, and the scale factor $a_{vac}^2(t) = e^{2\sqrt{\Lambda_{eff}^\pm}t} \equiv a_{dS_3}^2$ represents dS_3 geometry in agreement with the generic relation between Λ_{eff} and Λ [4]:

$$\Lambda_{eff}^\pm = -2m^2\left(\epsilon \pm \sqrt{1 + \lambda_{BHT}}\right) \quad (12)$$

We have two options for the initial conditions of $a(t)$ and $\sigma(t)$:

⁴In the BHT notations [4] we have $\lambda_{BHT} = -\frac{\Lambda}{m^2}$ and therefore both $\epsilon\lambda_0 < 0$ cases correspond to the limiting value $\lambda_{BHT} = -1$.

- type (1): $\sigma^2(t) \leq \sigma_{\pm}^2$, *i.e.*

$$\sigma(t \rightarrow \pm\infty) = \sigma_{\pm}, \quad \sigma_- \leq \sigma(t) \leq \sigma_+$$

$$a_{vac}^{\pm} = a_{dS_3}(t) = e^{\sqrt{\Lambda_{eff}}t}$$

- type (2): $\sigma^2(t) \geq \sigma_{\pm}^2$

– (2a)

$$\sigma_- < \sigma(t) \leq \sigma(t = t_0) = \infty$$

$$a_{vac}^- = a_{dS_3} \text{ but } a(t_0) = 0 \text{ and } R^{(3)}(t = t_0) = \infty$$

– (2b)

$$\sigma_+ \leq \sigma(t) < \sigma(t = t_0) = \infty$$

Let us consider first the *non – singular* solutions corresponding to the initial conditions of the *type (1)* above, *i.e.* the ones interpolating between the two vacuum states: $|vac\rangle = |\sigma_{\pm}, dS_3(\Lambda_{eff})\rangle$. We find by direct integration of eqs. (7) two exact solutions:

$$\sigma_{\pm}(t|\epsilon) = \sqrt{\frac{m_{\sigma}^2}{2\gamma_{\pm}}} \tanh(b_{\pm}(t - t_0)) \quad , \quad a_{\pm}^2(t|\epsilon) = e^{\varphi_{\pm}(t|\epsilon)} = \left(\cosh b_{\pm}(t - t_0) \right)^{\delta_{\pm}} \quad (13)$$

where δ_{\pm} and b_{\pm} are given by:

$$b_{\pm} = \frac{m^2}{\sqrt{2|m^2|}} \left(1 \pm \sqrt{1 + \epsilon \frac{m_{\sigma}^2}{2m^2}} \right) \quad , \quad \delta_{\pm} = -\epsilon \frac{m_{\sigma}^2}{2\gamma_{\pm}} = -2\epsilon \frac{|m^2|}{B_{\pm}^2} \quad (14)$$

Observe that $\sigma_{\pm}(t)$ formally coincide with the *flat space* QM "instanton" solution⁵ for the double well potential $V_{\pm}(\sigma)$, but now γ and m_{σ}^2 are related by eq. (9). The scale factor $a_{\pm}^2(t)$ resembles the well known bounce solution for closed $k = 1$ dS_3 Universe ($\lambda_0 = 0$, $\sigma = const$) as one can see by comparing the behavior of the corresponding Hubble parameters:

$$H_{\pm}^{k=0}(t, \epsilon, \lambda_0) = \frac{\dot{a}_{\pm}}{a_{\pm}} = -\epsilon B_{\pm} \sigma_{\pm}(t) \quad , \quad H^{k=1}(t) \sim \tanh 2\sqrt{\Lambda}(t - t_0)$$

The explicit form of the curvature scalar (11), *i.e.*

$$R_{\pm}(t) = \frac{8B_{\pm}^2}{\cosh^2 b_{\pm}(t - t_0)} \left(-\epsilon + \frac{3}{4} \sigma_{\pm}^2 \sinh^2 b_{\pm}(t - t_0) \right) \quad (15)$$

confirms that the bounce solution (13) represents non-singular asymptotically dS_3 ($(a)dS_3$) geometry for all $t \in (-\infty, \infty)$, in particular

$$R^{(3)}(t \rightarrow \pm\infty) = 12|m^2| = 6\Lambda_{eff}^{\pm} = \frac{6}{L_{gr}^2}, \quad L_{gr} \gg l_{pl} \quad (16)$$

Another important feature of this $(a)dS_3$ Bounce solution (13) is encoded in the behavior of deceleration parameter:

$$q_{\pm}(t) \equiv -1 - 2 \frac{\ddot{\varphi}}{\dot{\varphi}^2} = -1 - \frac{s_{\pm}(\epsilon)}{\sinh^2(b_{\pm}(t - t_0))} \quad (17)$$

⁵note that in our curved space example the time is not euclidean (see eq. (5)) so that it just looks like an instanton

where s_{\pm} are real numbers

$$s_{\pm}(\epsilon = -1) = \frac{1}{2} \left(1 \pm \sqrt{1 - \frac{m_{\sigma}^2}{2m^2}} \right) = \frac{B_{\pm}^2}{m^2} \quad , \quad s_{-}(\epsilon = 1) = \frac{1}{2} \left(1 - \sqrt{1 + \frac{m_{\sigma}^2}{2m^2}} \right) = -\frac{B_{-}^2}{|m^2|}$$

They are determined by the *ratio* of the two scales

$$L_{gr}^2 = \frac{1}{|\Lambda_{eff}|} = \frac{1}{2|m^2|} \quad , \quad L_{\sigma}^2 = \frac{1}{|m_{\sigma}^2|}$$

$$l_{pl} \ll L_{gr} < L_{\sigma} \tag{18}$$

introduced by the graviton mass $M_{gr}^2 = 2|m^2|$ and by the scalar field mass⁶ $|m_{\sigma}^2|$.

Note that our Bouncing 3D Universe is *almost* invariant under time reflections $T : t \rightarrow 2t_0 - t$ ($t \rightarrow -t$ for $t_0 = 0$). The *exact* invariance is achieved by combining T with the Z_2 - charge conjugation of $\sigma(t)$:

$$C : C\sigma_{(k)}(t) = -\sigma_{(k)}(t) = \sigma_{(ak)}(t)$$

which interchanges “kink” $\sigma_{(k)}(t)$ with the “anti-kink” $\sigma_{(ak)}(t) = -\sigma_{(k)}(t) = \sigma_{(k)}(-t)$ (*i.e.* $\sigma_{+} \rightarrow \sigma_{-}$ in the vacuum sector). Hence the past of the “kink” coincides with the future of the “anti-kink”. This fact indeed reflects the CPT - invariance ($Px^i = -x^i$, $i = 1, 2$) of the BHT - matter model (4).

Although the forms of the Bounce solutions (13) in the two cases $\{\epsilon = -1, \lambda_0 > 0\}$ and $\{\epsilon = 1, \lambda_0 < 0\}$ considered above are *almost* identical⁷, the features of the Universe evolutions they describe are completely different. The origin of the qualitatively different physics behind is in the form of the matter interaction V_{\pm} : *stable* vacuum double well potential for $\epsilon = -1$ versus *unstable* “maxima” of the inverse double well potential for $\epsilon = 1$ case⁸. Consider first the $\{\epsilon = -1, \lambda_0 > 0\}$ case. Since $s_{\pm}(\epsilon = -1) > 0$ is always positive, the deceleration (17) $q_{\pm}(t|\epsilon = -1) < 0$ is *negative* for all $t \in (-\infty, \infty)$ and therefore we have an *eternally* accelerated Universe. It starts its evolution from well defined (stable) dS_3 vacuum state $(\sigma_{-}, dS_3(\Lambda_{eff}))$ at the left minima of the matter potential living a period of (accelerated) expansion ($\dot{\varphi} > 0$ for $B < 0$) for all $t < t_0$. At the moment $t = t_0$ (say $t_0 = 0$) the Universe enters in the contraction ($\dot{\varphi} < 0$) period towards the new (“final”) vacuum state $(\sigma_{+}, dS_3(\Lambda_{eff}^{+}))$ with $\Delta\sigma = \sigma_{+} - \sigma_{-} = m_{\sigma}\sqrt{\frac{2}{\gamma}}$ and $\Lambda_{eff}^{+} = \Lambda_{eff}^{-} = 2m^2$.

In the second case $\{\epsilon = 1, \lambda_0 < 0\}$, that corresponds to the inverted double well potential $V_{-}(\sigma)$, due to the negative values of $s_{-}(\epsilon = 1) = -\frac{B_{-}^2}{|m^2|}$, the deceleration $q_{-}(t|\epsilon = 1, \lambda_0 < 0)$ changes twice its sign. Therefore we have in this case two acceleration epochs (initial and final ones) and one period of deceleration. By studying the properties of scale factor $a^2(t)$ and its derivatives \dot{a} and \ddot{a} we can reconstruct the history of this non-singular Bounce - like 3D Universe. At the infinite past $t \rightarrow -\infty$ when: $\sigma(t) \rightarrow \sigma_{-}$, $a_{-}^2(t) \rightarrow e^{2|m|\sqrt{2}t}$ it starts from the unstable maximum $(\sigma_{-}, dS_3(\Lambda_{eff}^{-}))$ of $V_{-}(\sigma)$ its first (semi-infinite) period of acceleration and contraction with $R^{(3)}(t|\epsilon = 1, \lambda_0 < 0)$ decreasing from $6\Lambda_{eff}^{-} = 2|m^2|$ to $R^{(3)}(t_1) = 0$. At the moment $t_1 < 0$

$$t_1 = -\frac{1}{b_{-}} \ln \left(\sqrt{\frac{2|s_{-}|}{3}} + \sqrt{\frac{2|s_{-}|}{3} + 1} \right),$$

⁶or by $\bar{m}_{\sigma}^2 = -\frac{m_{\sigma}^2}{2}$ for the case $\epsilon = 1$, since in this case $m_{\sigma}^2 < 0$, *i.e.* it describes tachyons.

⁷the only difference is in the scale factors $a^2(\epsilon = -1) = a^{-2}(\epsilon = 1)$.

⁸intrinsically related to the wrong/right signs $\epsilon = \pm 1$ of the Einstein-Hilbert term ϵR in the action (4)

the curvature $R^{(3)}(t_1 < t < -t_1) < 0$ changes its sign entering in the epoch of negative curvature of finite duration $T_- = 2|t_1|$ up to the moment $\bar{t}_1 = -t_1 > 0$ when it changes its sign once more towards the finite *unstable* $\{\sigma_+, dS_3\}$ - state. Concerning the acceleration we observe that at the moment $|t_a| > |t_1|$:

$$|t_a| = \frac{1}{b_-} \ln \left(\sqrt{|s_-|} + \sqrt{|s_-| + 1} \right)$$

the Universe enters in deceleration period which takes place over a finite period of time $T_{dec} = 2|t_a|$. The second (final) epoch of acceleration and expansion starts at $t = |t_a|$ towards its “*future eternity*” $\{\sigma_+, dS_3\}$ at $t \rightarrow \infty$.

An equivalent (but more compact) description of the Universe evolution is based on the so called equations of state: $p_{eff} = p_{eff}(\rho_{eff})$, where the effective pressure p_{eff} and energy density ρ_{eff} are defined by the diagonal r.h.s. of eqs. (3):

$$\langle vac | T_\nu^\mu(\sigma, h_{\alpha\beta}) | vac \rangle = (-\rho_{eff}, p_{eff}, p_{eff})$$

and it appears to be a simple consequence of the eqs. (6), (7):

$$p_{eff}(t) = 8\epsilon m^2 s_\pm + (s_\pm - 1)\rho_{eff}(t), \quad \rho_{eff} > 0 \quad (19)$$

In the $\{\epsilon = -1, \lambda_0 > 0\}$ case we have

$$p_{eff}(t) = -8B_\pm^2 - \rho_{eff}(t) \left(1 - \frac{B_\pm^2}{m^2} \right), \quad B_\pm^2 \leq m^2 \quad (20)$$

i.e. $p_{eff} < 0$ is always negative and that is why the 3D Universe is eternally accelerated. In the second case $\{\epsilon = 1, \lambda_0 < 0\}$ the pressure changes twice its sign (at the moments $\pm t_a$), due to the existence of critical value of the energy density

$$\rho_{eff}^{cr}(\epsilon = 1, \lambda_0 < 0) = \frac{8B_-^2 |m^2|}{|m^2| + B_-^2}$$

where p_{eff} vanishes. This fact leads to the observation that there exist two periods of negative pressure and another one of positive pressure between them.

An important problem in the classical (and quantum) Cosmology concerns the initial and/or final conditions (i/f. c.'s). Namely, whether and how the particular features of the Universe evolution depend on the corresponding i/f. c.'s. The BHT - matter model (4) provides an example of two distinct families of such conditions (type (1) and (2) above) that lead to qualitatively different Universe histories: *non-singular* Big-Bounce for the type (1) ($\sigma^2 \leq \sigma_\pm^2$) and the *singular* Big - Bang for the type (2) ($\sigma^2 \geq \sigma_\pm^2$) for the *same values* of all the parameters $\{\kappa^2, \lambda_0, m_\sigma^2\}$ of the model. The corresponding *type* (2) solutions:

$$\tilde{\sigma}(t|\epsilon) = \sqrt{\frac{m_\sigma^2}{2\gamma_\pm}} \coth(b_\pm(t - t_0)) \quad , \quad \tilde{a}_\pm^2(t|\epsilon) = |\sinh b_\pm(t - t_0)|^{\delta_\pm} \quad (21)$$

are “singular” at $t = t_0$, *i.e.*

$$\tilde{\sigma}(t \rightarrow t_0) \rightarrow \infty, \quad \dot{\tilde{\varphi}}(t \rightarrow t_0) \rightarrow \infty, \quad \tilde{R}^{(3)}(t \rightarrow t_0) \rightarrow \infty \quad (22)$$

and they exist in two disconnected periods of time. For $t \in (-\infty, t_0)$ we have the same vacuum $\{\sigma_-, dS_3(\Lambda_{eff})\}$ (stable for $\epsilon = -1$; unstable for $\epsilon = 1$) as starting point. In the case $\epsilon = -1, \lambda_0 > 0$

these solutions describe accelerating and contracting to Big-Crunch 3D Universe, while for the unstable inverse double-well potential V_- , *i.e.* in $\{\epsilon = 1, \lambda_0 < 0\}$ case, the initial acceleration period ends at the moment t_a ($t_a < t_0$), when $\tilde{q}(t)$ changes sign and the new deceleration epoch begins. Similarly the eqs. (21) when considered in the interval $t_0 < t < \infty$ represent accelerating and expanding Universe created in the moment t_0 from the Big-Bang singularity ($R^{(3)}(t_0) \rightarrow \infty$) towards the infinite future ($t \rightarrow \infty$) dS_3 - vacuum $\{\sigma_+, dS_3(\Lambda_{eff})\}$ in the “stable” $\{\epsilon = -1, \lambda_0 < 0\}$ case. Again the $\{\epsilon = 1, \lambda_0 < 0\}$ - Universe exhibits periods of acceleration and deceleration.

The purpose of this short discussion about the effects of the choice of the i/f c.’s $\{(\sigma_{i,f}, \dot{\sigma}_{i,f})\}$ and $\{(\varphi_{i,f}, \dot{\varphi}_{i,f})\}$ on the properties of the solutions of eqs. (3) for the BHT - Matter model (4) is to demonstrate the *incompatibility* of the type (2) i/f c.’s, involving scales $L_2 \leq l_{pl}$ (due to $< \tilde{R}^{(3)} > \rightarrow \infty$, $< \dot{\tilde{\varphi}} > \rightarrow \infty$), with the range of validity $L_2 \gg l_{pl}$ of eqs. (3). On the other hand, the type (1) i/f. c.’s (*i.e.* $\sigma^2 \leq \sigma_{\pm}^2$, $\dot{\varphi}^2 \leq 4B^2\sigma_{\pm}^2$, etc.) provides an example of “semiclassical consistency”, when the parameters of the model $\{\epsilon = \pm 1, \Lambda_{eff} = M_{gr}^2 = \frac{1}{L_{gr}^2}, m_{\sigma}^2 = \frac{1}{L_{\sigma}^2}\}$ are chosen such that⁹

$$l_{pl} \ll L_{gr} < L_{\sigma}$$

(remember $L_{gr} \equiv L_{(1)}$ is the smallest scale in this approximation).

The non-perturbative cosmological solutions of BHT - massive gravity coupled to scalar QFT’s with σ^4 self-interaction (4) we have constructed within the restrictions (9), (10) imposed by the super-potential method, *do not allow* a complete discussion of the “unitarity consistency” of these models. However one can easily verify whether (and for which values of the parameters $\epsilon = \pm 1, \lambda_0$ and m_{σ}^2) the above mentioned restrictions are *compatible* with the pure BHT - gravity unitarity conditions [4]:

$$\begin{aligned} m^2(\Lambda_{eff} - 2\epsilon m^2) &> 0 \\ M_{gr} = -\epsilon m^2 + \frac{1}{2}\Lambda_{eff} &\geq \Lambda_{eff} \end{aligned} \quad (23)$$

together with the well known conditions for unitarity and tachyon-free “light” massive scalar σ^4 - QFT (2) in dS_3 classical background :

$$0 < m_{\sigma}^2 \leq \Lambda_{eff} \quad (24)$$

We have already realized that the Bounce solutions (13) in both cases $\epsilon\lambda_0 < 0$, *i.e.* of normal and inverted double well potentials $V_{\pm}(\sigma)$, correspond to the limiting value $\lambda_{BHT} = -\frac{\Lambda}{m^2} = -1$ of the BHT “cosmological” parameter λ_{BHT} . As a consequence the effective cosmological constant

$$\Lambda_{eff}^+ = \Lambda_{eff}^- = 2|m^2|$$

is equal to the “graviton (*mass*)²”, *i.e.* $M_{gr}^2 = \Lambda_{eff} = 2|m^2|$. In the case: $\epsilon = -1, \lambda_0 > 0$ the BHT - u.c.’s (23) are automatically satisfied and the restriction (9) now reads

$$4\gamma < m_{\sigma}^2 \leq 2m^2 \quad (25)$$

Therefore the BHT - matter model (4) with “wrong” sign $\epsilon = -1$ of the curvature term, $\lambda_0 > 0$ and $V_+(\sigma)$ representing double well potential (2) with $\tilde{m}_{\sigma}^2\gamma < 0$ and $\gamma > 0$ is *unitary* and *tachyons-free*. It corresponds to the lower bound $\lambda_{BHT} = -1$ of the dS_3 unitary window [4]:

$$-1 \leq \lambda_{BHT} < 0 \quad (26)$$

⁹in both cases the treatment of (sub) Planckian scales $L \leq l_{pl}$ requires an introduction of other degrees of freedom (than $\hat{h}_{\mu\nu}$) and other methods of quantization

This special point of the pure BHT - gravity model (1) is known to describe “*partially massless*” graviton of one propagating mode, instead of the two helicities ± 2 graviton massive modes for the generic $M_{gr}^2 > \Lambda_{eff}$ case. The observed reduction of the physical degrees of freedom of $\hat{h}_{\mu\nu}$ (for $\lambda_{BHT} = -1$) turns out to be a result of the enhancement of the gauge symmetries of the linearized (effectively Pauli-Fierz) form of the corresponding equations of motion [16],[4].

We have indeed similar non-singular solutions (13) for the case $\epsilon = 1$, $\lambda_0 < 0$ (*i.e.* $m^2 < 0$) representing the inverted double well potential $V_-(\sigma)$ ($\gamma < 0$, $\bar{m}_\sigma^2 > 0$). However the BHT - matter model for this range of the parameters is *neither* unitary *nor* tachyons-free. For $\epsilon = 1$, $m^2 < 0$ and $\Lambda_{eff} = 2|m^2|$ the first of the BHT - u.c.’s (23) does not hold and now $\sigma_\pm - dS_3$ “vacuum” represent the maxima of $V_-(\sigma)$. Hence they are unstable and the corresponding asymptotic ($t \rightarrow \pm\infty$) σ - particle states are tachyonic ($m_{\sigma_\pm}^2 < 0$). Due to the “right” sign $\epsilon = 1$ of the R - term one can try to interpret this model as an effective action for the scalar QFT with $V_-(\sigma)$ self-interaction in *classical* gravitational background, thus relaxing the BHT - quantum gravity unitarity conditions (23). In order to avoid tachyonic instabilities one should consider “semiclassical” states build on the metastable minima of $V_-(\sigma)$ (*i.e.* $\sigma_0 = 0$) and to further study their decay. Note that such state is qualitatively different from the “constant curvature” $|\sigma_\pm, dS_3(\Lambda_{eff}) >$ - states. As one can easily see from eqs. (6), now we have

$$\dot{\sigma}(0) = 2B, \quad \ddot{\varphi}(0) = -4B^2, \quad \dot{\varphi}(0) = 0$$

As a result this relative minima state describes linearly growing matter and (non-constant curvature) Gaussian scalar factor

$$a_0^2(t) = e^{-2\epsilon B^2 t^2}, \quad \sigma_0(t) = 2Bt \tag{27}$$

which makes the analytic study of the spectrum of the fluctuations around it rather complicated.

Let us briefly discuss the remaining two models when $\epsilon\lambda_0 > 0$ (or equivalently $\bar{m}_\sigma^2\gamma > 0$), *i.e.*

$$V_\pm^{cyc}(\sigma) = \pm \frac{|\gamma|}{4} \left(\sigma^2 + \frac{2|m^2|}{B^2} \right)^2 - 2\Lambda \tag{28}$$

The solutions of eqs. (7) are now periodic of period $T = \frac{\pi|m|}{B^2\sqrt{2}}$:

$$\sigma_{cyc}(t) = \frac{|m|\sqrt{2}}{B} \tan\left(\frac{B^2\sqrt{2}}{|m|}t\right) \quad , \quad a_{cyc}^2(t) = \left| \cos\left(\frac{B^2\sqrt{2}}{|m|}t\right) \right|^{\frac{2\epsilon|m^2|}{B^2}} \tag{29}$$

and are defined on the finite interval $t \in (-\frac{T}{2}, \frac{T}{2})$ only, where they represent initial (Big-Bang) and final (Big-Crunch) curvature singularities with $\sigma_{cyc}(\pm\frac{T}{2}) \rightarrow \pm\infty$ at both “ends”. The main problem with these models (and of their solutions (29)) is that they do *not* admit “vacuum” solutions of constant curvature. The absolute minima/maxima (for $\epsilon = \pm 1$) of $V_\pm^{cyc}(\sigma)$, $\sigma_0 = 0$ corresponds to singular non-constant curvature solution (27). Note that the BHT - analysis of the perturbative consistency of 3D gravity is based on the linearization $\hat{g}_{\mu\nu} = g_{\mu\nu}^{vac} + \kappa\hat{h}_{\mu\nu}$ around constant curvature stable vacuum states $|\sigma_\pm, g_{\mu\nu}^{vac} >$ (*i.e.* the ones corresponding to M_3 , AdS_3 , or dS_3). They are indeed present in the pure BHT - gravity (1) for all values of the parameters ϵ , λ_0 and Λ . When matter is added, the presence or absence of such type of vacuum solutions depend on the form of the potential $V(\sigma)$ and on the values of the parameters ϵ , λ_0 , m_σ^2 . As we have seen on the examples of the double-well potentials ($\epsilon\lambda_0 < 0$) the analysis of perturbative “semiclassical consistency” of the BHT - matter models follows the lines of the BHT - arguments [4], [5] with special attention to the unitarity and renormalizability of the matter sector. In the $\epsilon\lambda_0 > 0$ type of models characterized

by the potential (28) such an analysis requires new methods and their consistency remains an open problem.

We have avoided up to now the discussion of the important question about the possible manners to couple Matter QFT's to the New Massive Gravity. To begin with the natural questions to ask are: Should one also include higher derivatives terms like : $R|\nabla\sigma|^2$, $|\nabla\sigma|^4$, $R^{\mu\nu}\nabla_\mu\sigma\nabla_\nu\sigma$, $RV(\sigma)$ etc.? How such terms might influence the unitarity properties of the corresponding Matter-Gravity model?, etc. Our particular choice (4) was motivated by the renormalizability of the matter QFT and we have included in the effective action the UV counter-terms (all of type K for the minimal coupling $\xi = 0$) that appear in the lowest order in the couplings κ^2 and γ . The analysis of the structure of divergent diagrams of this graviton -scalar QFT (in constant curvature background) at orders κ^3 and κ^4 suggests that new counter-terms like those mentioned above could be present. This line of arguments however unavoidably leads to the introduction of large number of new parameters in the corresponding effective action. On the other hand, the original unitarity proof of Bergshoeff, Hohm and Townsend [3] that is based on the equivalence of the NMG action (1) to the specific “second order” derivatives (including one more auxiliary spin two massive field $f^{\mu\nu}$) action

$$S_{PF}^{eff} = \frac{1}{\kappa^2} \int dx^3 \sqrt{-g} \left[-R + f^{\mu\nu} G_{\mu\nu} - \frac{m^2}{4} (f^{\mu\nu} f_{\mu\nu} - f^2) \right] \quad (30)$$

(with $f = g^{\mu\nu} f_{\mu\nu}$) appoints an economic alternative without any new parameters involved. Namely to consider the following rather obvious generalization of the Pauli - Fierz inspired action (30) including matter fields:

$$S_{PFM}^{eff} = \frac{1}{\kappa^2} \int dx^3 \sqrt{-g} \left\{ \epsilon R - \kappa^2 \left(\frac{1}{2} |\nabla\sigma|^2 + V(\sigma) \right) + f^{\mu\nu} E_{\mu\nu} - \frac{m^2}{4} (f^{\mu\nu} f_{\mu\nu} - f^2) \right\} \quad (31)$$

where we denote

$$E_{\mu\nu} = G_{\mu\nu} - \frac{\kappa^2}{2} T_{\mu\nu} \quad , \quad T_{\mu\nu}(\sigma) = \nabla_\mu\sigma\nabla_\nu\sigma - \frac{1}{2} g_{\mu\nu} \nabla_\rho\sigma\nabla^\rho\sigma - g_{\mu\nu} V(\sigma) \quad (32)$$

The matter potential $V(\sigma)$ has to be chosen by renormalizability arguments. Similarly to the pure gravity case [3], [17] one can easily eliminate $f_{\mu\nu}$ in order to get the following suggestive form of the New Massive Gravity - Matter action:

$$S_{NMG M}^{eff} = \frac{1}{\kappa^2} \int dx^3 \sqrt{-g} \left\{ \epsilon R + \frac{1}{m^2} K - \kappa^2 \left(\frac{1}{2} |\nabla\sigma|^2 + V(\sigma) \right) - \frac{\kappa^2}{m^2} \left(R^{\mu\nu} T_{\mu\nu} - \frac{1}{4} R T_\mu^\mu \right) + \frac{\kappa^4}{4m^2} \left(T^{\mu\nu} T_{\mu\nu} - \frac{1}{2} (T_\mu^\mu)^2 \right) \right\} \quad (33)$$

Observe that the last two (new) terms:

$$\begin{aligned} U &= R_{\mu\nu} T^{\mu\nu} - \frac{1}{4} R T_\mu^\mu = R^{\mu\nu} \nabla_\mu\sigma\nabla_\nu\sigma - \frac{1}{4} R V - \frac{3}{8} R |\nabla\sigma|^2 \\ Y &= T^{\mu\nu} T_{\mu\nu} - \frac{1}{2} (T_\mu^\mu)^2 = \frac{5}{8} |\nabla\sigma|^4 - \frac{3}{2} V^2 - \frac{1}{2} V |\nabla\sigma|^2 \end{aligned} \quad (34)$$

are of order κ^3 and κ^4 correspondingly since the κ -expansion of both $R_{\mu\nu}$ and R starts by order κ terms. Hence the above action (33) when terms up to order κ^2 only are taken into account reduces to the action (4) we have studied. Although the symmetry principles that single out such higher derivatives Gravity-Matter actions among the many others (with much more free parameters) are

not completely clear their unitarity and renormalizability indeed represent an interesting research problem, which however is out of the scopes of the present paper. Let us just mention that the unitarity of the matter sector in the complete action (33) is rather problematic due to the presence of the both $|\nabla\sigma|^2$ and $|\nabla\sigma|^4$ terms. The propagator of such massive scalar field contains ghost contributions unless certain fine tuning of the coefficients takes place.

The main objectives of the investigations presented in this paper are the properties and the perturbative consistency of the BHT - matter models (4) with specific renormalizable scalar 3D QFT added. We have considered as an example the σ^4 interaction potential that admits degenerate adiabatic vacuum states. Our goal was the construction of the explicit non-vacuum solutions of the corresponding “effective” action’s equations that possess “maximal” 2D Poincare symmetry $O(2) \otimes T_2$ and that interpolate between two such vacua. The particular Bounce solutions of the 3D Massive BHT - Cosmology with scalar matter (2) we have found by superpotential method turns out to saturate the lower bound $\lambda_{BHT} = -1$ (*i.e.* $\Lambda_{eff} = 2|m^2|$) of the BHT unitarity window. There exist strong indications that these solutions are stable as well, due to the fact that they carry Z_2 topological charges ± 1 . The power of the superpotential method we have used in this paper allows the construction of analytic solutions for a vast variety of potentials. The simplest case of linear superpotential we have studied in detail represents just one example selected by the renormalizability of σ^4 interactions in three dimensions. The generalization of our results to the case of simplest periodic superpotential $W(\sigma) = A + B \cos \sigma$ that corresponds to the modified Double Sine-Gordon potential $V(\sigma)$ is straightforward. Let us also mention the special case of BHT- gravity coupled to free massless scalar field, *i.e.* $V(\sigma) = const = -2\Lambda$. In this simple model a family of unitarity consistent Bounce solutions can be easily found for arbitrary values of the cosmological constant $0 < |\Lambda| < m^2$ without the use of any superpotentials.

Another important line of investigations concerns the construction of domain wall solutions of the Massive 3D Gravity -Matter models (2) and their further applications for the reconstruction of the renormalization group flows in the perturbed 2D CFT dual to these BHT-matter models. The well known relation between the FRW cosmological solutions and the static DW’s permits an extension of our results to the case of Janus kink-like DW’s for the model (4) that interpolate between two stable AdS_3 vacua. It turns out that they correspond again to the special value $\lambda_{BHT} = -1$, but now with $\Lambda_{eff} = -2|m^2|$ [18].

In conclusion: the unitarity consistency of the simplest Massive 3D Gravity - Matter model (4) we have demonstrated on the particular class of cosmological Bounce solutions confirms the expectations that the BHT - model indeed admits physically interesting renormalizable matter extensions.

Acknowledgments. We are grateful to C.P.Constantinidis for discussions and for critical reading of the manuscript. This work has been partially supported by PRONEX project No.35885149/2006 from FAPES-CNPq (Brazil).

References

- [1] G.t’Hooft and M.Veltman, One loop divergencies in the theory of gravitation, Ann.Inst.H.Poincare**A20**(1974),69
- [2] K.S.Stelle, Renormalization of higher-derivative quantum gravity, Phys.Rev.**D16**953(1977)
- [3] E.A.Bergshoeff, O.Hohm and P.K.Townsend, Massive Gravity in Three Dimensions, Phys.Rev.Lett.**102**,201301(2009)

- [4] E.A.Bergshoeff, O.Hohm and P.K.Townsend, More on Massive 3D Gravity, *Phys.Rev.D* **79**,124042(2009)
- [5] I.Oda, Renormalizability of Massive Gravity in Three Dimensions,JHEP 0905:064(2009); M.Nakasone and I.Oda, On Unitarity of Massive Gravity in Three Dimensions, *Prog.Theor.Phys.* 121:389 (2009)
- [6] Y.Liu and Y.W.Sun, On the Generalized Massive Gravity in AdS(3), *Phys.Rev. D* **79**,126001(2009); Y.Liu and Y.W.Sun, Consistent Boundary Conditions for New Massive Gravity in AdS(3),JHEP 0905:039(2009); Y.Liu and Y.W.Sun, Note on New Massive Gravity in AdS(3), *JHEP* 0904:106(2009)
- [7] G.Clement, Warped AdS(3) black holes in new massive gravity, *Class.Quant.Grav.* **26** : 105015(2009)
- [8] E.Ayon-Beato, G.Giribet and M.Hassaine, Bending AdS Waves with New Massive Gravity, *JHEP* 0905:029 (2009); J.Oliva,D.Tempo and R.Troncoso, Three-dimensional black holes, gravitational solitons,kinks and wormholes for BHT gravity, *JHEP* 0907:011(2009)
- [9] I.Gullu adn B.Tekin, Massive Higher Derivative Gravity in D-dimensional Anti-de Sitter Spacetimes, *Phys.Rev.D* **80**:064 033(2009)
- [10] N.D.Birrell and P.C.W.Davis, *Quantum Fields in Curved Space*, Cambridge University Press, 1982
- [11] T.S.Bunch, P.Panangaden and L.Parker ,On Renormalization of $\lambda\phi^4$ Field Theory in Curved Space-Time.I.,*J.Phys.A :Gen.Phys.*13,901(1980); T.S.Bunch and P.Panangaden ,On Renormalization of $\lambda\phi^4$ Field Theory in Curved Space-Time.II.,*J.Phys.A :Gen.Phys.*13,919(1980)
- [12] J.F.Donoghue, Introduction to Effective Field Theory Description of Gravity, arXiv:gr-qc/9512024
- [13] M.Cvetic, S.Griffies and S.J.Rey, Static Domain Walls in $N = 1$ Supergravity, *Nucl. Phys.* **B381** (1992) 301; K.Behrndt, G.Lopes Cardoso and D.Lust, Curved BPS Domain Wall solutions in four-dimensional $N = 2$ supergravity, *Nucl. Phys.* **B607** (2001) 391
- [14] D.Z.Freedman, C.Nunez, M.Schnabl and K.Skenderis, Fake supergravity and domain walls stability, *Phys.Rev.D* **69**,104 027(2004); J.Sonner and P.K.Townsend ,Dilaton domain walls and dynamical systems, *Class.Quant.Grav.***23**,441 (2006); K.Skenderis and P.K.Townsend, Hidden supersymmetry of domain walls and cosmology, *Phys.Rev.Lett.* **96**:191 301 (2006)
- [15] I.Low and A.Zee, Naked Singularity and Gauss-Bonnet term in Brane world scenarios, *Nucl. Phys.* **B585** (2000) 395
- [16] S.Deser and R.I.Nepomechie, Gauge Invariance versus Masslessness in De Sitter Space, *Ann.Phys.***154**,369 (1984); B.Tekin, Partially masseless spin 2 fields in string generated models, arXiv:hep-th/0306 178
- [17] S.Deser, Ghost-free, finite, fourth order $D = 3$ (alas) gravity, *Phys.Rev.Lett.***103**:101 302(2009)
- [18] H.L.C.Louzada, U.Camara dS and G.M.Sotkov, Massive 3D Gravity-Matter Domain Walls (in preparation)