

Modified Entropic Force

Changjun Gao*

*The National Astronomical Observatories, Chinese Academy of Sciences
Key Laboratory of Optical Astronomy, NAOC, CAS, Beijing, 100012 and
Kavli Institute for Theoretical Physics China, CAS, Beijing 100190, China*

(Dated: March 1, 2010)

The theory of thermodynamics tells us the equipartition law of energy does not hold in the limit of very low temperature. It is found the Debye model is very successful in explaining the experimental results. Motivated by this fact, we modify the entropic force formula which is proposed very recently. Since the Unruh temperature is proportional to the strength of gravitational field, so the modified entropic force formula is an extension of the Newtonian gravity to weak field. On the contrary, General Relativity extends Newtonian gravity to strong field case. Corresponding to Debye temperature, there exists a Debye acceleration g_D . It is found the Debye acceleration is $g_D = 10^{-14} \text{N} \cdot \text{kg}^{-1}$. This acceleration is very much smaller than the gravitational acceleration $10^{-4} \text{N} \cdot \text{kg}^{-1}$ which is felt by the Neptune. Therefore, the modified entropic force can be very well approximated by the Newtonian gravity in solar system. With this Debye acceleration, we also find the current cosmic speeding up can be explained without invoking any kind of dark energy.

PACS numbers: 98.80.Cq, 98.65.Dx

I. INTRODUCTION

Recently, Verlinde [1] makes one interesting proposal that gravity may be explained as an entropic force caused by the changes in the information associated with the positions of material bodies. The proposal is very interesting in the following two aspects. In the first place, with the assumption of the entropic force together with the Unruh temperature [2], Verlinde is able to derive the second law of Newtonian mechanics. Secondly, the assumption of the entropic force together with the holographic principle and the equipartition law of energy can lead to the Newtonian law of gravitation. Similar discoveries are also made by Padmanabhan [3]. Padmanabhan finds that using the equipartition law of energy for the horizon and the thermodynamic relation $S = E/2T$, one can obtain the Newtonian law of gravity. Here S and T are thermodynamic entropy and the temperature associated with the horizon. E is the active gravitational mass which produces the gravitational field in the spacetime [4]. Actually, Jacobson has derived the Einstein's equations by employing the Clausius relation and the equivalence principle in 1995 [5].

Using the entropic force, Shu and Gong [6], Cai, Cao and Ohta [7] derived the Friedmann equation nearly simultaneously while with different methods. On the other hand, using this entropic force, Smolin [8] derived the Newtonian gravity in loop quantum gravity. Mäkelä [9] pointed out that Verlinde's entropic force is actually the consequence of a specific microscopic model of spacetime [10]. Similar ideas were applied to the construction of holographic actions from black hole entropy by Caravelli and Modesto [11]. Li and Wang [12] showed that the

holographic dark energy [13] can be derived from the entropic force formula.¹

In this paper, motivated by the Debye's model in thermodynamics, we extend the entropic force formula. With this modification, we find the resulting entropic force can speed up the expansion of the Universe. In this paper, we use the units of $c = G = \hbar = k = 1$.

II. MODIFIED ENTROPIC FORCE

Following [1], we consider a compact spatial region V with a compact boundary B which is a sphere. The compact boundary B plays the role of a holographic screen. The number N of bits for freedom on the screen is taken to be equal to the area A of the screen:

$$N = A. \quad (1)$$

Let g denotes the proper gravitational acceleration on the screen which is produced by the matter sources inside the screen B . Then the Unruh effect tells us the gravitational field would correspond to a system filled with thermal gas with the temperature of T for a proper observer:

$$T = \frac{g}{2\pi}. \quad (2)$$

So according to the equipartition law of energy, the total energy inside the screen is

$$E = \frac{1}{2}NT. \quad (3)$$

*Electronic address: gaojcj@bao.ac.cn

¹ Same observation was disclosed by Johannes Koelman in his blog. (see <http://www.scientificblogging.com-hammock> physicist-it bit how get rid dark energy)

Verlinde assumes the energy is equal to the gravitational mass M inside the screen:

$$E = M . \quad (4)$$

Combining above equations, we can obtain the spherically symmetric and static gravitational field equations:

$$M = \frac{1}{2} \cdot 4\pi r^2 \cdot \frac{g}{2\pi} , \quad (5)$$

namely,

$$g = \frac{M}{r^2} . \quad (6)$$

It is exactly the Newtonian law of gravity. Thermodynamics tells us the equipartition law of energy fails in the limit of temperature approaching zero. It is found the Debye model is very successful. Motivated by this fact, we modify the equipartition law of energy as follows:

$$E = \frac{1}{2} NT \mathfrak{D}(x) , \quad (7)$$

where the Debye function is defined by

$$\mathfrak{D}(x) \equiv \frac{3}{x^3} \int_0^x \frac{y^3}{e^y - 1} dy . \quad (8)$$

x is related to the temperature T as follows:

$$x \equiv \frac{T_D}{T} . \quad (9)$$

T_D is the Debye critical temperature. For a gravitational system, T_D is related to the critical gravitational acceleration. So we obtain the modified entropic force from Eq. (7) as follows

$$g = \frac{M}{r^2} \cdot \frac{1}{\mathfrak{D}(x)} , \quad (10)$$

with

$$x \equiv \frac{g_D}{g} . \quad (11)$$

The Debye acceleration is related to the Debye temperature

$$g_D = 2\pi T_D . \quad (12)$$

We note that when $x \ll 1$, i.e. in the limit of strong gravitational field, we have

$$\mathfrak{D}(x) = 1 . \quad (13)$$

So the Newtonian gravity is recovered. On the other hand, in the limit of weak gravitational field, i.e. $x \gg 1$, we have

$$\mathfrak{D}(x) = \frac{\pi^4}{5x^3} , \quad (14)$$

and

$$g = \left(\frac{5Mg_D^3}{\pi^4} \right)^{\frac{1}{4}} \cdot \frac{1}{\sqrt{r}} . \quad (15)$$

In this limit, the gravitational field is very different from Newtonian gravity. In the next, let's look for the value of g_D from the considerations of cosmological physics. Using the method of Shu and Gong [6], we obtain the following equation

$$4\pi(\rho + p) = - \left(\dot{H} - \frac{K}{a^2} \right) \left[-2\mathfrak{D}(x) + \frac{3x}{e^x - 1} \right] . \quad (16)$$

Here ρ , p are the total energy density and total pressure of cosmic fluids, respectively. a , H are the scale factor and Hubble parameter, respectively. $K = 0, +1, -1$ describe the three kind of geometry of the Universe. Dot denotes the derivative with respect to the comoving time t . Taking into account the Hawking temperature T for the Universe [14, 15]

$$T = \frac{H}{2\pi} , \quad (17)$$

we find x can be rewritten as

$$x \equiv \frac{H_D}{H} . \quad (18)$$

Thus the critical Hubble parameter H_D is exactly the Debye acceleration g_D :

$$H_D = g_D . \quad (19)$$

Eq. (16) together with the energy conservation equation

$$\dot{\rho} + 3H(\rho + p) = 0 , \quad (20)$$

govern the evolution of the Universe. Using these two equations, we find the current cosmic acceleration can be interpreted while without invoking any kind of dark energy. To show this point, we may investigate the cosmic evolution sourced by dark matter and baryon matter. Observations reveal that the Universe is highly flat in space. So in the following we shall put $K = 0$. Define

$$u \equiv \ln a . \quad (21)$$

Combining Eq. (16) and Eq. (20) we obtain the Friedmann equation:

$$8\pi\rho = \int 3 \left[-2\mathfrak{D}(x) + \frac{3x}{e^x - 1} \right] dH^2 . \quad (22)$$

Let

$$h \equiv \frac{H}{H_0} , \quad \Omega_{m0} \equiv \frac{\rho_{m0}}{\rho_0} \quad H_D = \zeta H_0 , \quad (23)$$

then we have

$$\Omega_{m0} e^{-3u} = \int_{\infty}^x \left[4\mathfrak{D}(x) - \frac{6x}{e^x - 1} \right] \cdot \frac{\zeta^2}{x^3} dx , \quad (24)$$




FIG. 1: The dimensionless Hubble parameter h with redshift z . The solid line is for the standard Λ CDM model. The circled line is for our model.




FIG. 2: The deceleration parameter q with redshift z . The solid line is for the standard Λ CDM model. The circled line is for our model.

and

$$x = \frac{\zeta}{h}. \quad (25)$$

Here H_0 , ρ_0 , ρ_{m0} are the Hubble parameter, total energy density and matter energy density ratio for the present-day Universe. ζ is a dimensionless free parameter. The standard cosmological model, e.g. as in Komatsu et al. [16], predicts a present matter density ratio $\Omega_{m0} = 0.25$. Using this result and taking $\zeta = 10^{-5}$, we plot the dimensionless Hubble parameter h via redshifts z in Fig. 1 for our model and the standard Λ CDM model. We find they have nearly the same behavior.

In order to show our model can lead to the cosmic acceleration, we plot the evolution of deceleration parameter q :

$$q \equiv - \left(1 + \frac{d \ln h}{du} \right), \quad (26)$$

in Fig. 2. We find the model predicts nearly the same transition redshift $z_T = 0.8$ as the standard Λ CDM model.

At higher redshifts (large H), we have $x \ll 1$. Then Eq. (16) restores to the standard equation in General Relativity. So the physics at and before the radiation dominated epoch is not modified. Finally, let's calculate the Debye acceleration g_D :

$$g_D = \zeta H_0 = 10^{-14} \text{N} \cdot \text{kg}^{-1}. \quad (27)$$

It is the order of 10^{-10} of the gravitational field strength in the solar system. So Eq. (10) can be perfectly written as the Newtonian law of gravity in solar system.

III. CONCLUSION AND DISCUSSION

Thermodynamics tells us the equipartition law of energy does not hold in the limit of temperature approaching zero. It is found that the Debye model is very successful in interpreting the physics of very low temperature system. Since a gravitational system can also be taken as a thermodynamical system, we expect the equipartition law of energy for gravitational system shall break down in the limit of very low temperature (or very low gravitational field strength). Motivated by Debye model, we modified the entropic force. We find the modification can interpret the current acceleration of the Universe while without invoking any kind of dark energy.

The model is closely related the Debye acceleration g_D . We find $g_D = 10^{-14} \text{N} \cdot \text{kg}^{-1}$ is consistent with cosmic observations very well. This gravitational strength is very smaller than the gravitational field strength in solar system by orders of 10^{10} . Therefore, the modified entropic force is not inconsistent with solar system tests. How to derive the Debye model from the specific microscopic model of spacetime? We leave this problem to future study.

Acknowledgments

We sincerely thank Dr. Savvas Nesseris for stimulating and illuminating discussions which help me find two typographical errors in Eq. (22) and Eq. (24). This work is supported by the National Science Foundation of China under the Distinguished Young Scholar Grant 10525314, the Key Project Grant 10533010, Grant 10575004, Grant 10973014 and the 973 Project (No. 2010CB833004).

[1] E. P. Verlinde, "On the Origin of Gravity and the Laws of Newton," arXiv:1001.0785 [hep-th].

[2] W. G. Unruh, "Notes on black hole evaporation," Phys.

- Rev. D **14**, 870 (1976).
- [3] T. Padmanabhan, “Equipartition of energy in the horizon degrees of freedom and the emergence of gravity,” arXiv:0912.3165 [gr-qc].
 - [4] T. Padmanabhan, “Gravitational entropy of static spacetimes and microscopic density of states,” *Class. Quant. Grav.* **21**, 4485 (2004) [arXiv:gr-qc/0308070].
 - [5] T. Jacobson, “Thermodynamics of space-time: The Einstein equation of state,” *Phys. Rev. Lett.* **75**, 1260 (1995) [arXiv:gr-qc/9504004].
 - [6] F-W, Shu and Y. Gong, “Equipartition of energy and the first law of thermodynamics at the apparent horizon,” arXiv:1001.3237 [gr-qc].
 - [7] R-G, Cai, L-M cao and N. Ohta, “Friedmann Equations from Entropic Force,” arXiv:1001.3470 [hep-th].
 - [8] L. Smolin, “Newtonian gravity in loop quantum gravity,” arXiv:1001.3668 [gr-qc].
 - [9] J. Mäkelä, “Notes Concerning ‘On the Origin of Gravity and the Laws of Newton’ by E. Verlinde,” arXiv:1001.3808 [gr-qc]
 - [10] J. Mäkelä, arXiv: 0805.3952; arXiv: 0805.3955; arXiv: 0810.4910.
 - [11] F. Caravelli and L. Modesto, “Holographic actions from black hole entropy” arXiv:1001.4364 [gr-qc].
 - [12] M, Li and Y. Wang, “Quantum UV/IR Relations and Holographic Dark Energy from Entropic Force” arXiv:1001.4466 [hep-th].
 - [13] M. Li, *Phys. Lett. B* **603**, 1 (2004).
 - [14] R. G. Cai and S. P. Kim, “First law of thermodynamics and Friedmann equations of Friedmann-Robertson-Walker universe,” *JHEP.* **0502**, 050 (2005) [arXiv:hep-th/0501055];
 - [15] R. G. Cai, L. M. Cao and Y. P. Hu, “Hawking Radiation of Apparent Horizon in a FRW Universe,” *Class. Quant. Grav.* **26**, 155018 (2009) [arXiv:0809.1554 [hep-th]].
 - [16] E. Komatsu *et al*, *Astrophys. J. Suppl.* **180**, 330 (2009).