

Revised checklist of Turkish Agromyzidae (Diptera) fauna of Turkey

Hasan Sungur CİVELEK^{1,*}, Emine ÇIKMAN², Oktay DURSUN¹

¹Muğla University, Faculty of Arts and Science, Biology Department 48170 Kötekli, Muğla - TURKEY

²Harran University, Agriculture Faculty, Plant Protection Department, Şanlıurfa - TURKEY

Received: 28.08.2008

Abstract: This study was carried out in the years of 2007 and 2008 in some provinces of Turkey. Specimens of leafminers were collected on cultured and non-cultured plants during 1 year. In this study, 29 new records belonging to 7 genera were found for the Turkish fauna. With this study, the number of leafminer species, which was 136 previously, has been updated to 165 species.

Key words: Checklist, new records, Agromyzidae, Turkey

Türkiye Agromyzidae (Diptera) faunasının gözden geçirilmiş tür listesi

Özet: Bu çalışma 2007–2008 yılları arasında Türkiye’de bazı yörelerde yürütülmüştür. Galerisineği örnekleri bir yıl boyunca kültürü yapılan ve kültürü yapılmayan bitkiler üzerinden toplanmıştır. Bu çalışmada Türkiye faunası için yedi cinsde ait 29 yeni kayıt bulunmuştur. Bu çalışmaya birlikte önceden 136 türü bilinen galerisineği tür sayısı 165 türü yükselmiştir.

Anahtar sözcükler: Kontrol listesi, yeni kayıt, Agromyzidae, Türkiye

Introduction

Agromyzidae (leafminer flies) is one of the largest fly families, with more than 2790 valid species belonging to 27 genera worldwide (Spencer, 1989; Gu et al., 1991; Pakalniskis, 1992, 1994; Woodley and Janzen, 1995; Pakalniskis, 1996; Sasakawa, 1997; Pakalniskis, 2000; Cerny, 2001, 2005a, 2005b, Sasakawa, 2005; Cerny, 2007a, 2007b; Çıkman and

Sasakawa, 2008). From this family, about 1171 species have been identified in the Palearctic region (Scheirs et al., 1999; Pakalniskis, 2000; Cerny, 2005b; Çıkman and Sasakawa, 2008). Many species were described in the period from the 19th century to the 60s of the last century but only a few isolated cases of damage by Agromyzidae were recorded during this period. Hering (1957) made a special study of all leaf-mining insects in various orders, and produced keys to

* E-mail: chasan@mu.edu.tr

European leafminers arranged by plant genera. However, in his study, Hering (1957) attached too great an importance to the food plant and trivial differences in the leafminers as diagnostic characters, and many new species that he described on these bases have been reduced to synonyms since then. An even more significant advance in clarifying the systematics of the family has been made during the past 45 years with the study of male genitalia. Spencer (1964, 1972, 1973, 1976, 1989, 1990) and Tschirnhaus (1991, 1994, 1999) contributed to the Agromyzidae both in terms of number of taxa and systematics with many studies from 1960 to the present. Adults can be minute, with a wing length of little more than 1 mm. The maximum size known is 6.5 mm. The majority of species are in the wing length range of 2 to 3 mm. Agromyzids are typically phytophagous, with larvae living in the tissues of living plants. Larvae of all the agromyzid species are internal plant feeders with a family specific morphology. Larvae of the most leafminers feed within the leaf parenchyma. Some species are stemborers or develop in roots, seeds or galls. One genus develops exclusively in the cambium of the young and old trees.

The Agromyzidae are one of the most important fly families in Turkey, because of their pest status, especially on vegetable and ornamental plants in the greenhouses. The Turkish agromyzid fauna is poorly known. One of the reasons is that only few entomologists identified Agromyzidae captured in the Turkish territory (e.g. Giray, 1980; Uygun et al., 1995;

Yabaş et al., 1995; Civelek et al., 2000a, 200b). Another important reason is the difficulties in identification of the Agromyzidae in general. As a result, in the past 15 years, many studies have been carried out. Over the past 70 years some 136 agromyzid species have been identified and reported in Turkey in various publications (Civelek, 2003; Cerny and Merz, 2006; Civelek et al., 2008). It is estimated that the actual Agromyzidae fauna of Turkey is in the range of at least 200–300 species.

The goal of this study was to contribute to the knowledge of the leafminer fauna of Turkey and to clarify which valid leafminer species are in Turkey. Also this study will provide a base for future researches regarding the fauna of leafminer species.

Materials and methods

Seven geographical regions of Turkey divided into latitudes and longitudes stated by Yiğit and Çolak (1998) were chosen to make the survey economic and practical. The regions are shown in the Figure. The field studies carried out in 6 regions formed 5 longitudes every month between May and October in 2007 and between May and July in 2008. The leafminer species were collected from the selected provinces stated in the Table for each region.

The leafminer specimens were collected from both cultured and non-cultured plants. A GPS (Global Positioning System) was used while the samples were being collected. The adults were obtained by




Figure. Six regions of Turkey formed 5 longitudes as stated by Yiğit and Çolak (1998).

Table. Degree of the longitude, number of the region and name of the selected cities.

Degree of the longitude	Number of the region	Name of the selected cities
24- 28	1	Edirne, Tekirdağ, Çanakkale, İzmir
28- 32	2	İzmit, Manisa, İsparta, Antalya
32- 36	3	Kastamonu, Yozgat, Aksaray, Mersin
36- 40	4	Ordu, Kayseri, Malatya, Hatay
40- 44	5	Trabzon, Artvin, Bingöl, Mardin
44- 48	6	Ardahan, Ağrı, Van, Hakkari

sweeping. Since the male genitalia are important characters for identification of the leafminers, slide preparations were made. The following general procedures were applied: the abdomen of each male was boiled in 10% KOH, then transferred into 5% glacial acetic acid for 5 min and subsequently transferred to 96% alcohol for 5 min. Afterwards the abdomen was further dissected under a stereoscopic microscope. The male genitalia were transferred into euparal on a micromount pinned under the individual specimen in order to preserve the material perpetuity (Civelek et al., 2000a). Identifications of the species were made using Spencer (1972, 1973, 1976, 1989, 1990) and Cerny (2007a, 2007b).

Results

In this study, 29 newly recorded species were added to the Turkish leafminer fauna. Representative specimens were deposited at the laboratories in the Science and Arts Faculty, Biology Department, Entomology Laboratory, Muğla, and Harran University, Agriculture Faculty, Plant Protection Department, Şanlıurfa, Turkey.

The taxa are presented alphabetically with their distributions in Turkey.

Subfamily 1. Agromyzinae

Genus 1. *Agromyza* Fallén, 1810

Agromyza abiens Zetterstedt, 1848

Mardin (Çıkman and Civelek, 2005)

Agromyza albipennis Meigen, 1830

Şanlıurfa (Çıkman and Uygun, 2003)

- Agromyza albatarsis* Meigen, 1830
İzmir (Giray, 1980); Mardin (Çıkman and Civelek, 2005)
- Agromyza ambigua* Fallén, 1823
Aydın (new record)
- Agromyza anderssoni* Spencer, 1976
Kırklareli (European part of Turkey, new record)
- Agromyza anthracina* Meigen, 1830
Erzurum (Civelek et al., 2007)
- Agromyza apfelbecki* Strobl, 1902
İzmir (Giray, 1980)
- Agromyza bromi* Spencer, 1966
Erzurum (Civelek et al., 2007)
- Agromyza felleri* Hering, 1941
Diyarbakır, Mardin (Çıkman and Sasakawa, 2005)
- Agromyza flavipennis* Hendel, 1920
İsparta (new record)
- Agromyza frontella* (Rondani, 1875)
İzmir (Civelek et al., 2000a).
- Agromyza graminicola* Hendel, 1931
Erzurum (Civelek et al., 2007)
- Agromyza hiemalis* Becker, 1908
İzmir (Giray, 1980); Adana and Hatay (Uygun et al., 1995)
- Agromyza idaeiana* Hardy, 1853
Artvin (Civelek et al., 2007)
- Agromyza intermittens* Becker, 1907

- İzmir (Deeming and Civelek, 1997)
Agromyza marionae Griffiths, 1963
Hatay (new record)
Agromyza megalopsis Hering, 1933
Turkey (Cerny and Merz, 2006)
Agromyza myosotidis Kaltenbach, 1864
Isparta (new record)
Agromyza nana Meigen, 1830
İzmir (Deeming and Civelek, 1997)
Agromyza nigrescens Hendel, 1920
Turkey (Cerny and Merz, 2006)
Agromyza nigripes Meigen, 1830
Turkey (Cerny and Merz, 2006)
Agromyza phylloposthia Sasakawa, 2008
Adiyaman (Çıkman and Sasakawa, 2008)
Agromyza pittodes Hendel, 1931
Muğla (new record)
Agromyza prespana Spencer, 1957
Turkey (Cerny and Merz, 2006)
Agromyza reptans Fallén, 1823
Turkey (Hering, 1931-1936); Şanlıurfa (Çıkman and Uygın, 2003)
Agromyza rondensis Strobl, 1900
İzmir (Deeming and Civelek, 1997); Şanlıurfa (Çıkman and Uygın, 2003)
Agromyza trebinjensis Strobl, 1900
İzmir (Giray, 1980)
Agromyza vicifoliae Hering, 1932
Mardin (Çıkman and Sasakawa, 2005).

Genus 2. ***Hexomyza*** Enderlein, 1936
Hexomyza simplex (Loew 1869)
Kırklareli (European part of Turkey, new record)

Genus 3. ***Japanagromyza*** Sasakawa, 1958
Japanagromyza salicifolii (Collin, 1911)
Turkey (Hering, 1957); Izmir (Giray, 1980)
Genus 4. ***Melanagromyza*** Hendel, 1920

Melanagromyza aeneoventris (Fallén, 1823)
Adiyaman (Çıkman and Sasakawa, 2006)
Melanagromyza cunctans (Meigen, 1830)
Muğla (Civelek, 2003)
Melanagromyza pubescens Hendel, 1923
Adiyaman (Çıkman and Sasakawa, 2006)
Melanagromyza tripolii Spencer, 1957
Elazığ (Civelek et al., 2007)
Melanagromyza vignalis Spencer, 1959
Şanlıurfa (Çıkman and Uygın, 2003)

Genus 5. ***Ophiomyia*** Braschnikov, 1897
Ophiomyia beckeri (Hendel, 1923)
Mardin (Çıkman and Sasakawa, 2005)
Ophiomyia cunctata (Hendel, 1920)
İzmir (Civelek et al., 2000a); Şanlıurfa (Çıkman and Uygın, 2003)
Ophiomyia curvipalpis (Zetterstedt, 1848)
İzmir (Civelek et al., 2000a)
Ophiomyia delphinii Hendel, 1928
Muğla (Cerny and Merz, 2007)
Ophiomyia galii Hering, 1937
Muğla (Cerny and Merz, 2007)
Ophiomyia inaequabilis (Hendel, 1931)
Turkey (Hendel, 1931-1936); Izmir (Civelek et al., 2000a)
Ophiomyia labiatarum Hering, 1937
Turkey (Cerny and Merz, 2006)
Ophiomyia maura (Meigen, 1838)
Çanakkale (Hepdurgun et al., 2007)
Ophiomyia memorabilis Spencer, 1974
Turkey (Cerny and Merz, 2006)
Ophiomyia orbiculata (Hendel, 1931)
İzmir (Deeming and Civelek, 1997)
Ophiomyia phaseoli (Tryon, 1895)
Adana, Hatay (Civelek and Ulusoy, 2000)
Ophiomyia pinguis (Fallén, 1820)
Turkey (Hendel, 1931-1936); Şanlıurfa (Çıkman and Uygın, 2003)

<i>Ophiomyia pulicaria</i> (Meigen, 1830)	Genus 4. <i>Cerodontha</i> Rondani, 1861
Muğla (Civelek, 2002)	Subgenus 1. <i>Butomomyza</i> Nowakowski, 1967
<i>Ophiomyia subaura</i> Hering, 1926	<i>Cerodontha (Butomomyza) angulata</i> (Loew, 1869)
Turkey (Hendel, 1931-1936).	Turkey: (Hendel, 1931-1936)
Subfamily 2. Phytomyzinae	<i>Cerodontha (Butomomyza) pseuderrans</i> (Hendel, 1931)
Genus 1. <i>Amauromyza</i> Hendel, 1931	Rize (new record)
Subgenus 1. <i>Amauromyza</i> Hendel, 1931	<i>Cerodontha (Butomomyza) scutellaris</i> (von Roser, 1840)
<i>Amauromyza</i> (<i>Amauromyza</i>) <i>morionella</i> (Zetterstedt, 1848)	Şanlıurfa (new record)
İzmir (Giray, 1980).	
Subgenus 2. <i>Cephalomyza</i> Hendel, 1931	Subgenus 2. <i>Cerodontha</i> Rondani, 1861
<i>Amauromyza</i> (<i>Cephalomyza</i>) <i>flavifrons</i> (Meigen, 1830)	<i>Cerodontha (Cerodontha) denticornis</i> (Panzer, 1806)
İzmir (Civelek et al., 2000b).	Turkey (Hendel, 1931-1936); İzmir (Civelek, 1998); Şanlıurfa (Çıkman and Uygun, 2003); Diyarbakır and Mardin (Çıkman and Civelek, 2005)
<i>Amauromyza</i> (<i>Cephalomyza</i>) <i>luteiceps</i> (Hendel, 1920)	<i>Cerodontha (Cerodontha) fulvipes</i> (Meigen, 1830)
Muğla (Civelek et al., 2000b)	Artvin, Rize, Trabzon (new record)
Genus 2. <i>Aulagromyza</i> Enderlein, 1936	Subgenus 3. <i>Dizygomyza</i> Hendel, 1920
<i>Aulagromyza buhri</i> (de Meijere, 1938)	<i>Cerodontha (Dizygomyza) bimaculata</i> (Meigen, 1830)
Muğla (Civelek, 2002).	Isparta (new record)
<i>Aulagromyza hamata</i> (Hendel, 1932)	
İzmir (Giray, 1980).	Subgenus 4. <i>Poemyza</i> Hendel, 1931
<i>Aulagromyza heringii</i> (Hendel, 1920)	<i>Cerodontha (Poemyza) imbuta</i> (Meigen, 1838)
Muğla (Giray, 1980)	Rize (new record)
<i>Aulagromyza orphana</i> (Hendel, 1920)	<i>Cerodontha (Poemyza) incisa</i> (Meigen, 1830)
İzmir (Civelek et al., 2000).	Mersin (new record)
<i>Aulagromyza populi</i> (Kaltenbach, 1864)	<i>Cerodontha (Poemyza) lateralis</i> (Macquart, 1835)
Isparta, İzmir (Giray, 1980)	Turkey (Cerny and Merz, 2006)
<i>Aulagromyza trivittata</i> (Loew, 1873)	<i>Cerodontha (Poemyza) pygmaea</i> (Meigen, 1830)
Balıkesir (Hepdurgun et al., 2007)	Turkey (Cerny and Merz, 2006)
Genus 3. <i>Calycomyza</i> Hendel, 1931	Genus 5. <i>Chromatomyia</i> Hardy, 1849
<i>Calycomyza humeralis</i> (von Roser, 1840)	<i>Chromatomyia ciliata</i> (Hendel, 1935)
İzmir (Civelek et al., 2000a); Şanlıurfa (Çıkman and Uygun, 2003); Mardin (Çıkman and Civelek, 2005)	Erzurum (Civelek et al., 2007)
	<i>Chromatomyia fuscula</i> (Zetterstedt, 1838)

- Kahramanmaraş (Mart et al., 2005)
- Chromatomyia horticola* (Goureau, 1851)
Adana, İstanbul, İzmir, Muğla (Giray, 1980);
Adana and Hatay (Uygun et al., 1995); Şanlıurfa
(Çıkman and Uygun, 2003); Ankara (Gençer, 2004);
Diyarbakır and Mardin (Çıkman and Civelek, 2005)
- Chromatomyia luzulae* (Hering, 1924)
Rize (Civelek et al., 2007)
- Chromatomyia milii* (Kaltenbach, 1864)
İzmir (Civelek et al., 2000a)
- Chromatomyia nigra* (Meigen, 1830)
Elazığ (Civelek et al., 2007)
- Chromatomyia scolopendri* (Robineau-Desvoidy,
1851)
Muğla (Civelek, 2002).
- Chromatomyia succisae* (Hering, 1922)
Turkey (Cerny and Merz, 2006).
- Genus 6. *Galiomyza* Spencer, 1981
Galiomyza violiphaga (Hendel, 1932)
Çanakkale (Hepdurgun et al., 2007)
- Genus 7. *Liriomyza* Mik, 1894
Liriomyza alyssi (Hering, 1960)
Turkey (Cerny and Merz, 2006)
Liriomyza amarella Hering, 1963
Çanakkale (Hepdurgun et al., 2007)
Liriomyza amoena (Meigen, 1830)
İzmir (Giray, 1980)
Liriomyza balcanica (Strobl, 1900)
İzmir (Civelek et al., 2000a).
Liriomyza brassicae (Riley, 1884)
Antalya, İzmir (Giray, 1980).
Liriomyza bryoniae (Kaltenbach, 1858)
İzmir (Giray, 1980); Adana and Hatay (Uygun et
al., 1995).
Liriomyza buhri Hering, 1937
Antalya (new record)
- Liriomyza cannabis* Hendel, 1931
İzmir, Kastamonu (Giray, 1980)
- Liriomyza cardariae* Sasakawa, 2008
Adiyaman (Çıkman and Sasakawa, 2008)
- Liriomyza centaureae* Hering, 1927
İzmir (Civelek et al., 2000a)
- Liriomyza cepae* (Hering, 1927)
İzmir (Giray, 1980)
- Liriomyza cicerina* (Rondani, 1875)
İzmir (Lodos, 1962); Adana and Hatay (Uygun et
al., 1995); Şanlıurfa (Çıkman and Uygun, 2003).
- Liriomyza congesta* (Becker, 1903)
İzmir, Manisa (Giray, 1980) Şanlıurfa (Çıkman
and Uygun, 2003); Mardin (Çıkman and Civelek,
2005)
- Liriomyza dracunculi* Hering, 1932
Rize (Civelek et al., 2007)
- Liriomyza eupatorii* (Kaltenbach, 1873)
Mersin (new record)
- Liriomyza euphorbiae* Martinez, 1999
Isparta (Martinez and Sobhian, 1999)
- Liriomyza flaveola* (Fallén 1823)
Muğla (Civelek, 2002)
- Liriomyza freyella* Spencer, 1976
Kırklareli (European part of Turkey, new record)
- Liriomyza hieracivora* Spencer, 1971
Şanlıurfa (Çıkman and Uygun, 2003)
- Liriomyza huidobrensis* (Blanchard, 1926)
Adana, İzmir (Yabaş et al., 1995); Aydın (Yıldırım,
2002)
- Liriomyza orbona* (Meigen, 1830)
İzmir (Deeming and Civelek, 1997)
- Liriomyza pascuum* (Meigen, 1838)
Muğla (Civelek, 2004)
- Liriomyza pedestris* Hendel, 1931
Turkey (Cerny and Merz, 2006)
- Liriomyza pusilla* (Meigen, 1830)

- Turkey (Hendel, 1931–1936)
- Liriomyza sativae* Blanchard, 1938
- Muğla (Civelek, 2002); Diyarbakır (Çıkman and Civelek, 2005)
- Liriomyza strigata* (Meigen, 1830)
- Denizli, İzmir (Giray, 1980); Adana and Hatay (Uygun et al., 1995); Şanlıurfa (Çıkman and Uygun, 2003); Mardin (Çıkman and Civelek, 2005)
- Liriomyza trifolii* (Burgess in Comstock, 1880)
- Adana, Hatay (Uygun et al., 1995); Şanlıurfa (Çıkman and Uygun, 2003); Aydın (Yıldırım, 2002); Muğla (Civelek and Weintraub, 2003); Ankara (Gençer, 2004); Diyarbakır and Mardin (Çıkman and Civelek, 2005)
- Liriomyza virgula* Frey, 1946
- Mersin (new record)
- Genus 8. *Metopomyza* Enderlein, 1936
- Metopomyza scutellata* (Fallén, 1823)
- Mersin (new record)
- Metopomyza xanthaspis* (Loew, 1858)
- Mersin (new record)
- Genus 9. *Napomyza* Westwood, 1840
- Napomyza bellidis* Griffiths, 1967
- Hatay (new record)
- Napomyza carotae* Spencer, 1966
- Erzurum (Civelek et al., 2007).
- Napomyza cichorii* Spencer, 1966
- Isparta (new record)
- Napomyza elegans* (Meigen, 1830)
- Mardin (Çıkman and Civelek, 2005).
- Napomyza lateralis* (Fallén, 1823)
- İzmir (Civelek et al., 2000a); Diyarbakır (Çıkman and Civelek, 2005)
- Genus 10. *Phytoliriomyza* Hendel, 1931
- Phytoliriomyza arctica* (Lundbeck, 1901)
- Turkey (Spencer, 1964); İzmir (Civelek et al., 2000a)
- Phytoliriomyza dorsata* (Siebke, 1863)
- Diyarbakır (Çıkman and Civelek, 2005)
- Phytoliriomyza oasis* (Becker, 1907)
- Turkey (Cerny and Merz, 2006)
- Phytoliriomyza perpusilla* (Meigen, 1830)
- Kahramanmaraş (Mart et al., 2005)
- Genus 11. *Phytomyza* Fallén, 1810
- Phytomyza aconitophila* Hendel, 1927
- Muğla (Civelek, 2004)
- Phytomyza affinis* Fallén, 1823
- Turkey (Bodenheimer, 1958); Şanlıurfa (Çıkman and Uygun, 2003)
- Phytomyza albipennis* (Fallén, 1823)
- Muğla (Civelek et al., 2007)
- Phytomyza angelicae* Kaltenbach, 1872
- Muğla (Civelek, 2002)
- Phytomyza astrantiae* Hendel, 1924
- Trabzon (Civelek et al., 2007)
- Phytomyza aquilonia* Frey, 1946
- Diyarbakır (Çıkman and Civelek, 2005)
- Phytomyza aurei* Hering, 1931
- Kırklareli (European part of Turkey, new record)
- Phytomyza brevifacies* Hendel, 1934
- Rize (Civelek et al., 2007)
- Phytomyza cecidonomia* Hering, 1937
- Muğla (new record)
- Phytomyza chelonei* Spencer, 1969
- Şanlıurfa (Çıkman and Uygun, 2003)
- Phytomyza cirsii* Hendel, 1923
- Turkey (Hering, 1957)
- Phytomyza clematidis* Kaltenbach, 1859
- İzmir (Civelek et al., 2000a).
- Phytomyza continua* Hendel, 1920
- Turkey (Cerny and Merz, 2006).
- Phytomyza conyzae* Hering, 1920

- Muğla (Civelek, 2002)
Phytomyza crassiseta Zetterstedt, 1860
Bursa, Çanakkale, İzmit, Muğla (new record)
Phytomyza fallaciosa Brischke, 1880
Kahramanmaraş (Mart et al., 2005).
Phytomyza ferulae Hering, 1927
İzmir (Giray, 1980)
Phytomyza flavigornis Fallén, 1823
Turkey (Cerny and Merz, 2006).
Phytomyza geminata Sasakawa, 2008
Adiyaman (Çıkman and Sasakawa, 2008)
Phytomyza glabra Hendel, 1935
Erzurum (Civelek et al., 2007).
Phytomyza gymnostoma Loew, 1858
İzmir (Civelek et al., 2000a)
Phytomyza kaltenbachi Hendel, 1922
Balikesir, İzmir, Manisa (Giray, 1980)
Phytomyza lappae Goureau, 1851
Ankara (Gençer, 2004)
Phytomyza nigritula Zetterstedt, 1838
Burdur (Civelek et al., 2007)
Phytomyza origani Hering, 1931
Çanakkale (new record)
Phytomyza orobanchia Kaltenbach, 1864
Balikesir, Bursa, Manisa (Civelek and Demirkhan, 1998); Şanlıurfa (Çıkman and Uygun, 2003); Diyarbakır, Mardin (Çıkman and Civelek, 2005)
Phytomyza petoei Hering, 1924
İzmir (Civelek et al., 2000a)
Phytomyza phillyreae Hering in Buhr, 1930
Turkey (Cerny and Merz, 2006)
Phytomyza plantaginis Robineau-Desvoidy, 1851
İzmir (Civelek et al., 2000a); Mardin (Çıkman and Civelek, 2005)
Phytomyza podagrariae Hering, 1954
Çanakkale (new record)
Phytomyza pubicornis Hendel, 1920
Isparta (new record)
- Phytomyza ranunculi* (Schrank, 1803)
İzmir (Deeming and Civelek, 1997)
Phytomyza rostrata Hering, 1934
İzmit (new record)
Phytomyza rufescens von Roser, 1840
İzmir (Civelek et al., 2000a)
Phytomyza rufipes Meigen, 1830
İzmir (Civelek et al., 2000a); Mardin (Çıkman and Civelek, 2005)
Phytomyza sii Hering, 1930
Kırklareli (European part of Turkey, new record)
Phytomyza spoliata Strobl, 1906
Turkey (Hering, 1957)
Phytomyza socia Brischke, 1880
Diyarbakır (Çıkman and Sasakawa, 2005)
Phytomyza tenella Meigen, 1830
İzmir (Civelek et al., 2000a)
Phytomyza tetrasticha Hendel, 1927
İzmir (Giray, 1980)
Phytomyza thysselinivora Hering, 1924
Muğla (Civelek, 2002)
- Genus 12. **Pseudonapomyza** Hendel, 1920
Pseudonapomyza atra (Meigen, 1830)
Şanlıurfa (Çıkman and Uygun, 2003)
Pseudonapomyza europaea Spencer, 1973
Turkey (Cerny, 2005b)
Pseudonapomyza hispanica Spencer, 1973
İzmir (Civelek et al., 2000a)
Pseudonapomyza pyriformis Sasakawa, 2008
Adiyaman (Çıkman and Sasakawa, 2008)
Pseudonapomyza spicata (Malloch, 1914)
Şanlıurfa (Çıkman and Uygun, 2003)
Pseudonapomyza spinosa Spencer, 1973
Şanlıurfa (Çıkman and Uygun, 2003)
Pseudonapomyza strobliana Spencer, 1973
Turkey (Cerny, 2005b)

Conclusion

With this study, the number of leafminer species, which was 136 previously, has been updated to 165 species.

References

- Bodenheimer, F.S. 1958. Türkiye'de Ziraate ve Agaçlara Zararı olan Böcekler ve Bunlarla Savaş Hakkında Bir Etud. Bayur Matbaası, Ankara, 346 s.
- Cerny, M. 2001. *Phytobia bohemica* sp. n. from the Czech Republic (Diptera: Agromyzidae). Heyrovskyaná 9: 53–67.
- Cerny, M. 2005a. *Amauromyza (Amauromyza) maltensis* sp. nov. (Diptera: Agromyzidae) with account of agromyzid mining flies from the Republic of Malta. Folia Heyrovskyaná 12: 85–104.
- Cerny, M. 2005b. A new species of *Pseudonapomyza* from Egypt. With notes on distribution of some other Palaearctic species of the genus (Diptera: Agromyzidae) (pp. 95–100). In: Kubík, Š. and Barták, M. (eds). Dipterologica bohemoslovaka 11. Folia Fac. Sci. Nat. Univ. Masaryk. Brun., Biol. 109 (2004): 95–100.
- Cerny, M. 2007a. Description of eight new species of Agromyzidae (Diptera) from North Korea, including new records. Studia dipterologica 14: 209–229.
- Cerny, M. 2007b. New faunistic records of Agromyzidae (Diptera) from Andorra including descriptions of three new species. Boletín Sociedad Entomológica Aragonesa 41: 43–51.
- Cerny, M. and Merz, B. 2006. New records of Agromyzidae (Diptera) from the Palaearctic Region. Mitteilungen Der Schweizerischen Entomologischen Gesellschaft 79: 77–106.
- Cerny, M. and Merz, B. 2007. New records of Agromyzidae (Diptera) from the West Palaearctic Region, with an updated checklist for Switzerland. Mitteilungen Der Schweizerischen Entomologischen Gesellschaft Bulletin De La Société Entomologique Suisse 80: 107–121.
- Cerny, M., Vala, M. and Barták, M. 2001. Agromyzidae. In: Barták M. and Vaňhara J. (eds.): Diptera in an Industrially Affected Region (North-Western Bohemia, Bílina and Duchcov Environs), II. Folia Fac. Sci. Nat. Univ. Masaryk. Brun., Biol., 105: 349–364.
- Civelek, H. S. 1998. Systematic Studies on the Species of The Family Agromyzidae (Diptera) In Izmir Province. Ege University, Science Institute, PhD Thesis, Plant Protection Department, Bornova, Izmir, 187 pp. (Turkish, with an English summary).
- Civelek, H.S. 2002. New records for the Turkish leafminer fauna from Western Turkey. Insecta Mundi 16: 49–55.
- Civelek, H.S. 2003. Checklist of Agromyzidae (Diptera) of Turkey, with a new record. Phytoparasitica 31: 132–138.
- Civelek, H.S. 2004. Two new records for the Turkish Agromyzidae (Diptera) fauna. Turk. J. Entomol. 28: 15–19.
- Civelek, H. S. and Demirkhan, H. 1998. Investigations on infestation with *Phytomyza orobanchia* of *Orobanche* that is the problem in process tomatoes fields in Marmara Region. – Proceedings of the 2nd Turkish Herbology Congress, (1997, Izmir, Turkey), pp.45–51 (Turkish, with an English summary).
- Civelek, H.S. and Ulusoy, M.R. 2000. A new record for the Turkish leafminer fauna: *Ophiomyia phaseoli* (Tryon, 1895). Turk. J. Entomol. 24: 163–166 (Turkish, with English summary).
- Civelek, H.S. and Weintraub, P.G. 2003. Effects of Bensulfat on larval serpentine leafminers, *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae), in tomatoes. Crop Protection, 22: 479–483.
- Civelek, H.S., Deeming, J.C. and Önder, F. 2000a. Some new records for Turkish leafminers (Diptera: Agromyzidae) fauna from Izmir province. Turk. J. Entomol. 24: 17–26.
- Civelek, H.S., Önder, F. and Deeming, J.C. 2000b. Two new records for the Turkish *Amauromyza* Hendel (Diptera: Agromyzidae) fauna from Aegean region Turkey. Turk. J. Entomol. 24: 83–86.
- Civelek, H.S., Tonguç, A., Özgül O. and Dursun, O. 2007. Contributions to The Turkish Agromyzidae (Diptera) Fauna from Anatolian Part of Turkey, with sixteen New Records. Mitt. Internat. Entomol. 30: 21–28.
- Civelek, H.S., Dursun, O., Eskin, A., and Taç, G. 2008. A study on Agromyzidae (Diptera) fauna of Turkey and species list. Anadolu University, Journal of Science and Technology 9: 1–16 (Turkish, with English summary).
- Çikman, E. and Uygun, N. 2003. The determination of Agromyzidae (Diptera) and their parasitoids in cultivated and non-cultivated areas in Şanlıurfa province. Turk. J. Entomol. 27: 305–318 (Turkish, with English summary).
- Çikman, E. and Civelek, H.S. 2005. Contributions to the leafminer fauna (Diptera: Agromyzidae) from Turkey, with four new records. Phytoparasitica 33: 391–396.
- Çikman, E. and Sasakawa, M. 2005. The new records of Agromyzidae (Diptera) from southern Turkey. Int. J. Dipteral. Res. 16: 169–173.
- Çikman, E. and Sasakawa, M. 2006. Two new records for the Turkish Agromyzidae (Diptera) fauna. Turk. J. Zool. 30: 255–260.
- Çikman, E. and Sasakawa, M. 2008. The Turkish Agromyzidae (Diptera), with descriptions of four new species. Entomol. Sci. 11: 81–86.
- Deeming, J.C. and Civelek, H.S. 1997. New records for the Turkish Agromyzidae (Diptera) family. In: Proceeding 3rd Turkish Entomology Congr., Ankara, Turkey, pp. 526–533 (Turkish, with English summary).

Acknowledgements

Many thanks to The Scientific and Technological Research Council of Turkey (TÜBİTAK, 106T122), Sıtkı Koçman Vakfı, and Tükkiye İş Bankası for supporting this study.

- Gençer, L. 2004. A Study on the Chalcidoid (Hymenoptera: Chalcidoidea) Parasitoids of Leafminers (Diptera: Agromyzidae) in Ankara Province. *Turk. J. Zool.* 28: 119-122.
- Giray, H. 1980. Notes on the Primary List of Leafminers in Turkey: Their Hosts and Distributions. *Publ. Ege Univ. Agric. Fac.* 374: 1-106 (Turkish, with English summary).
- Gu, X., Fan, Z. and Sasakawa, M. 1991. Descriptions of Seven New Species of Agromyzidae (Diptera) from China. *Japanese Journal of Entomology* 59: 331-342.
- Hendel, F. 1923. Neue europäische *Melanagromyza*-Arten (Dipt.). *Konowia*. 2: 142-145.
- Hendel, F. 1931-1936. 59. Agromyzidae. In: Lindner, E. [Ed.] *Die Fliegen Palaearktischen Region*. Vol. 6. E. Schweizerbart'sche Verlagsbuch-Handlung, Stuttgart, Germany, XII + 570 pp. Taf. I-XVI.
- Hepdurgun, B., Civelek, H.S., Turanlı, T. and Dursun, O. 2007. Contributions to Turkish Agromyzidae (Diptera) fauna. *Turk. J. Entomol.* 31: 153-159.
- Hering, E.M., 1957. Bestimmungstabellen der Blattminen von Europa einschließlich des Mittelmeerbeckens und der Kanarischen Inseln. Vol. 1-3. Dr. W. Junk, The Hague, The Netherlands.
- Lodos, N. 1962. Two harmful flies on chickpea in Aegean region: *Liriomyza cicerina* (Rondani) and *Phytomyza atricornis* Meigen. *Plant Protection Bulletin* 2: 44-48. (Turkish with English summary).
- Mart, C., Tursun, A.Ö. and Civelek, H.S. 2005. Contributions to the Agromyzidae (Diptera) fauna of Turkey. *Turk. J. Zool.* 29: 357-359.
- Martinez, M. and Sobhian, R. 1999. A new palaearctic species of leafminer on *Euphorbia* spp.: *Liriomyza euphorbiae* Martinez, n. sp. (Diptera, Agromyzidae). *Nouvelle Revue d'Entomologie* (N.S.) 15 (1998): 273-277.
- Pakalniskis, S. 1992. Notes on Lithuanian Agromyzidae (Diptera) with the description of three species new to science. In *New and Rare for Lithuania Insect Species. Records and descriptions of 1992*. Vilnius pp. 47-55.
- Pakalniskis, S. 1994. The Lithuanian Agromyzidae (Diptera). Descriptions of 6 new species and other notes. *Acta entomologica Lituanica* 12: 5-34.
- Pakalniskis, S. 1996. The Lithuanian Agromyzidae (Diptera). Descriptions of 4 new species and other notes. In *Lietuvos entomologo darbai*, Vilnius, pp. 17-34.
- Pakalniskis, S. 2000. New data on the bionomics and distribution of Agromyzidae (Diptera) with the description of a new species. *Acta Zoologica Lituanica*, 10: 59-62.
- Sasakawa, M. 1997. Lauxaniidae and Agromyzidae (Diptera) of the Ryukyus. *ESAKIA* 37: 141-148.
- Sasakawa, M. 2005. The Neotropical Agromyzidae (Insecta: Diptera). Part 6. New or Little-known Species from El Salvador and Guatemala. *Species Diversity* 10: 151-169.
- Scheirs, J., Bruyn, L. de and Tschirnhaus, M. von 1999. Agromyzidae (Diptera) of the nature reserve "Etang de Virelles": faunistics and life-history aspects. *Bulletin S.R.B.E./K.B.V.E.* 135: 152-158.
- Spencer, K.A., 1964. The genus *Phytoliriomyza* Hendel (Agromyzidae, Diptera): a clarification of the four European species, with a list of eight other species now known in the genus. *Annals and Magazine of Natural History*, Ser. 13, vol. VII, 657 pp.
- Spencer, K.A. 1972. Diptera Agromyzidae. *Handbooks for the Identification of British Insects*, 10. 1-136. Royal entomological society of London.
- Spencer, K.A. 1973. Agromyzidae (Diptera) of Economic Importance. *Series entomologica*, 9. 1-405. Dr. W. Junk, The Hague, The Netharlans.
- Spencer, K.A. 1976. The Agromyzidae (Diptera) of Fennoscandia and Denmark. *Fauna Entomol. Scand.* 5: 1-606.
- Spencer, K. A. 1989. Family Agromyzidae. In: Evenhuis, N.L. [Ed]. *Catalog of the Diptera of the Australasian and Oceanian Regions*. Bishop Museum Spec. Publ. 86: 538-547.
- Spencer, K.A. 1990. Host Specialization in the world Agromyzidae (Diptera). *Series Entomologica* 45. 1-444. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Tschirnhaus, M. von. 1991. New results on the ecology, morphology and systematics of Agromyzidae (Diptera). In: Weisman, L., Orszagh, I. and Pont, A.C. [Eds.] *Proc. Second Int. Congr. Of Dipterology, VEDA* (Bratislava, Slovakia) pp. 285-313.
- Tschirnhaus, M. von. 1994. Minierfliegen (Diptera: Agromyzidae) aus Malaise-Fallén in spezifischen Pflanzengesellschaften: Ein Weinberg der Ahr-Eifel in Entwicklung zu einem Felsenbirnen-Gebüsch (*Cotoneastro-Amelanchieretum*). *Beitr. Landespfllege Rheinland-Pfaz* 16: 481-534.
- Tschirnhaus, M. von. 1999. Agromyzidae. In: Schumann, H., Bährmann, R. and Stark, A. [Eds.] *Entomofauna Germanica. Checkliste der Diptera Deutschland. Studia Dipterologica Suppl.* 2: 118-130. Ampyx-Verlag, Halle/Saale, Germany.
- Uygun, N., Polatöz, Z. and Başpinar, H. 1995. Faunistic studies on Agromyzidae (Diptera) in the south east Mediterranean region of Turkey. *Turk. J. Entomol.* 19: 123-136 (Turkish with English summary).
- Woodley, N.E. and Janzen, D.H. 1995. A new species of *Melanagromyza* (Diptera: Agromyzidae) mining leaves of *Bromelia pinguin* (Bromeliaceae) in a dry forest in Costa Rica. *Journal of Natural History* 29: 1329-1337.
- Yabaş, C., Civelek, H.S. and Ulubilir, A. 1995. A new leafminer *Liriomyza huidobrensis* (Blanchard, 1926) on vegetables for Turkey. *Turk. J. Entomol.* 19: 117-122 (Turkish with English summary).
- Yıldırım, E.M. 2002. Investigation on control possibilities of some botanical pesticides against *Liriomyza huidobrensis* and its population density on cucumber in greenhouse. Adnan Menderes University, Science Institute, Master Science Thesis, Aydın, 39 pp. (Turkish with English summary).
- Yiğit, N. and Çolak, E. 1998. Contribution to the Geographic Distribution of Rodent Species and Ecological Analyses of Their Habitats in Asiatic Turkey. *Tr. J. of Biology* 22: 435-446.