

The Karyotypes of Some Rodent Species (Mammalia: Rodentia) from Eastern Turkey and Northern Iran with a New Record, *Microtus schidlovskii* Argyropulo, 1933, from Eastern Turkey

Nuri YİĞİT¹, Mohammed Moradi GHARKHELOO², Ercüment ÇOLAK¹, Şakir ÖZKURT³,
Şafak BULUT¹, Tolga KANKILIÇ¹, Reyhan ÇOLAK¹

¹Ankara University, Faculty of Science, Department of Biology, Beşevler, Ankara - TURKEY

²Department of Biology, Faculty of Science, Zanjan University, Zanjan - IRAN

³Gazi University, Kirşehir Education Faculty, Kirşehir - TURKEY

Received: 24.01.2006

Abstract: During a field excursion in eastern Turkey and the Zanjan province of Iran, 9 rodent species were captured, and the diploid number of chromosomes ($2n$), the number of autosomal arms (FNa) and the fundamental number (FN) were determined for these species. *Microtus schidlovskii* with $2n = 60$ and FNa = 58 was recorded for the first time from Özalp (Van) and Yüksekova (Hakkari). The karyological record of *Meriones vinogradovi* in Turkey was also given for the first time. The karyological results were determined as follows: *Spermophilus xanthophrymnus* ($2n = 42$, FNa = 78, FN = 82 in Turkey), *Mesocricetus brandti* ($2n = 42$, FNa = 78, FN = 82 in Iran), *Microtus socialis* ($2n = 62$, FNa = 60 FN = 62 in Iran, $2n = 60$, FNa = 58 FN = 60 in Turkey), *Meriones tristrami* ($2n = 72$, FNa = 78, FN = 80 in Iran, $2n = 72$, FNa = 82, FN = 84 in Turkey), *M. vinogradovi* ($2n = 44$, FNa = 74, FN = 77 in Turkey), *Meriones libycus* ($2n = 44$, FNa = 76, FN = 80 in Iran), *Apodemus iconicus* ($2n = 48$, FNa = 46, FN = 48 in Iran), and *Mus domesticus* ($2n = 40$, FNa = 38, FN = 40 in Iran and Turkey).

Key Words: Rodent, karyotype, Turkey, Iran

Türkiye'nin Doğusu ve İran'ın Kuzeyinden Bazı Rodent (Mammalian: Rodentia) Türlerinin Karyotipi; Türkiye'nin Doğusundan Yeni Bir Kayıt *Microtus schidlovskii* Argyropulo, 1933

Özet: Türkiye'nin doğusunda ve İran'ın Zanjan şehri civarında yapılan arazi çalışmasında, 9 kemirici türü yakalandı ve bu türlere ait diploid kromozom sayıları ($2n$), otozomal kromozomların kol sayıları (FNa) ve kromozom kol sayıları (FN) belirlendi. $2n = 60$, FNa = 58 karyolojik değerlere sahip *Microtus schidlovskii* Özalp (Van) ve Yüksekova'dan (Hakkari) ilk kez kaydedildi. Çalışılan türlerden *Meriones vinogradovi*'nın karyolojik bulguları Türkiye'den ilk kez bu çalışmaya verildi. Karyolojik çalışmalarda; *Spermophilus xanthophrymnus* (Türkiye; $2n = 42$, FNa = 78, FN = 82), *Mesocricetus brandti* (İran; $2n = 42$, FNa = 78, FN = 82), *Microtus socialis* (İran; $2n = 62$, FNa = 60 FN = 62, Türkiye; $2n = 60$, FNa = 58 FN = 60), *Meriones tristrami* (İran; $2n = 72$, FNa = 78, FN = 80, Türkiye; $2n = 72$, FNa = 82, FN = 84), *M. vinogradovi* (Türkiye; $2n = 44$, FNa = 74, FN = 77), *Meriones libycus* (İran; $2n = 44$, FNa = 76, FN = 80), *Apodemus iconicus* (İran; $2n = 48$, FNa = 46, FN = 48), *Mus domesticus* (İran ve Türkiye; $2n = 40$, FNa = 38, FN = 40) karyolojik değerlere sahip olduğu saptandı.

Anahtar Sözcükler: Kemirici, karyotip, Türkiye, Iran

Introduction

To date, more than 65 rodent species have been recorded from Turkey, and the karyotypes of Turkish rodents have been intensively investigated (Soldatovic and Savic, 1978; Gülkaç and Yüksel, 1989; Doğramacı and Kefelioğlu, 1991; Doğramacı et al., 1994; Çolak et al., 1994; Yiğit et al., 1997, 1998a, 1998b, 1998c, 2000a, 2000b, 2001; Ivanitskaya et al., 1997; Çolak et al., 1997a, 1997b, 1997c, 1998, 1999, 2005; Kivanç et al.,

1997a, 1997b; Çolak and Yiğit, 1998; Yiğit and Çolak, 1998, 1999; Kefelioğlu and Kryštufek, 1999; Öz Kurt, et al., 1999, 2002; Çoşkun, 1999, 2003; Sözen et al., 2000; Özkan et al., 2003; Çolak 2003; Gözcelioğlu et al., 2005). However, karyotypes of Iranian rodents are poorly known (Mathey, 1954; Nadler and Lay, 1967; Benazzou et al., 1982; Gharkhelo and Kivanç, 2003). Furthermore, karyotypes of certain species show geographical variation in the fundamental number of

chromosomes as well as the diploid number of chromosomes as in the genera *Nannospalax* and *Meriones* (Nevo et al., 1994; Yiğit et al., 1998a, 1998b, 1998c; Sözen et al., 2000). The present study focused on the karyotypes of rodent species from eastern Turkey and northern Iran due to the very rare karyological records from those areas, and aimed to reveal the geographical variation in the fundamental number and the diploid number of chromosomes of these species.

Materials and Methods

During a field excursion in eastern Turkey and the Zanjan province of Iran, we caught 8 rodent species (Figure). Measurements of 4 external characters (total length, tail length, ear length and hindfoot length in mm) and weight (g) were taken from specimens, which were karyotyped in accordance with the conventional bone marrow method. A total of 20 slides were prepared for each specimen, and well-spread metaphase cells were analyzed. The diploid number of chromosomes ($2n$), the number of autosomal arms (FNa) and the fundamental number (FN) were determined by examining photographs of the slides. The zygomatic index ($ZI = \text{width of malar process} / \text{width of upper part of zygomatic arch}$) and tail length percentage to head and body length ($\text{tail length} \times 100 / \text{head body length} = T / HB$) were determined to identify *Mus* species. Voucher specimens were skinned and stuffed in the standard museum manner. The skin, skulls and karyotype preparations were deposited in

Ankara University, Faculty of Science, Department of Biology.

Specimens karyotyped: *Spermophilus xanthophrymnus* (Coll. Nr. 4518, 4595, 4600), *Mesocricetus brandti* (Coll. Nr. 4865, 4431, 4432, 4867), *Microtus socialis* (Coll. Nr. 4375, 4616), *Microtus schidlovskii* (Coll. Nr. 4607, 4608, 4616, 4617, 4618, 4619), *Meriones tristrami* (Coll. Nr. 4385, 4396), *Meriones vinogradovi* (Coll. Nr. 4404), *Meriones libycus* (Coll. Nr. 4381), *Apodemus iconicus* (Coll. Nr. 4606), and *Mus domesticus* (Coll. Nr. 4381, 4421).

Results and Discussion

Spermophilus xanthophrymnus (Bennet, 1835)

We karyotyped 2 female specimens and 1 male specimen from 3 different localities, namely Doğubeyazıt (İlgdır), Digor (Kars) and Başkale (Van). The karyotype is composed of $2n = 42$, $FNa = 78$ and $FN = 82$ with 2 pairs of metacentric, 17 pairs of submetacentric and a pair of acrocentric chromosomes. The X chromosome is medium-sized and metacentric, and the Y chromosome is the smallest acrocentric (Table). Our findings are consistent with those given by Özku et al. (2002).

Mesocricetus brandti Nehring, 1898

Two female specimens and 1 male specimen of *Mesocricetus brandti* were examined from Zanjan




Figure. Map showing the sampling locations in eastern Turkey (1 = Digor, 2 = Doğubeyazıt, 3 = Özalp, 4 = Başkale, 5 = Yüksekova) and northern Iran (6 = Zanjan).

Table. A comparison of karyological records of the given rodent species.

Species	Location / Country	2n	NFa	NF	References
<i>S. xanthophrymnus</i>	Doğubeyazıt / Turkey	42	78	82	Present study
<i>S. xanthophrymnus</i>	Western Turkey	42	78	82	Özkurt et al. (2002)
<i>M. brandti</i>	Zanjan / Iran	42	78	82	Present study
<i>M. brandti</i>	Turkey	42	78,80	80,84	Yiğit et al. (2000a)
<i>M. socialis</i>	Zanjan / Iran	62	60	62	Present study
<i>M. schidlovskii</i>	Özalp & Yüksekova / Turkey	60	58	60	Present study
<i>M. schidlovskii</i>	Caucasia, Georgia	60	58	60	Golenishchev et al. (2002)
<i>M. socialis</i>	Turkey	62	60	62	Kefelioğlu (1995)
<i>M. socialis</i>	Turkey	62	60	62	Kefelioğlu and Kryštufek (1999)
<i>M. tristrami</i>	Doğubeyazıt / Turkey	72	82	84	Present study
<i>M. tristrami</i>	Zanjan / Iran	72	78	80	Present study
<i>M. tristrami</i>	Iran	72	82	84	Benazzou et al. (1982)
<i>M. tristrami</i>	Iran and Palestine	72	70	74	Matthey (1957)
<i>M. tristrami blackieri</i>	Manisa / Turkey	72	72	76	Yiğit et al. (1998a)
<i>M. tristrami lycaon</i>	Karaman / Turkey	72	78	82	Yiğit et al. (1998a)
<i>M. tristrami kilensis</i>	Kilis / Turkey	72	74	78	Yiğit and Çolak (1998)
<i>M. vinogradovi</i>	Doğubeyazıt / Turkey	44	74	77	Present study
<i>M. vinogradovi</i>	Iran	44	78	-	Nadler and Lay (1967)
<i>M. libycus</i>	Zanjan / Iran	44	80	76	Present study
<i>M. libycus</i>	Iran	44	77	74	Benazzou et al. (1982)
<i>M. libycus</i>	Jordan	44	77	74	Qumsiyesh (1986)
<i>A. iconicus</i>	Zanjan / Iran	48	46	48	Present study
<i>A. iconicus</i>	Turkey	48	46	48	Çolak (2003)
<i>M. domesticus</i>	Zanjan / Iran	40	38	40	Present study
<i>M. domesticus</i>	Northwest Turkey	40	38	40	Gözelioğlu et al. (2005)

province (Iran). Their karyotypes have $2n = 42$, $FNa = 78$ and $FN = 82$. The karyotype consists of 8 metacentric chromosomes, 32 submetacentric chromosomes and 2 acrocentric chromosomes. The X chromosomes are middle-sized metacentric and the Y chromosome is smaller than the X chromosome and metacentric (Table). Yiğit et al. (2000a) reported $FNa = 80$, $FN = 84$ for Eastern Anatolia (Ardahan and Van specimens), and $FNa = 78$, $FN = 80$ for Central Anatolia. FNa and FN of Iranian specimens were similar to those of the specimens from Central Anatolia.

Microtus socialis (Pallas, 1771)

The diploid number of chromosomes ($2n$) is 62 in a specimen from Zanjan (Iran), and the karyotype has 62

acrocentric chromosomes with $FNa = 60$ and $FN = 62$. The X and the Y chromosomes are both acrocentric and the X chromosome is larger than the Y chromosome (Table). The karyological values of Iranian specimens were similar to those reported for Turkish specimens by Kefelioğlu (1995) and Kefelioğlu and Kryštufek (1999).

Microtus schidlovskii Argyropulo, 1933

In the specimens from Özalp (Van) and Yüksekova (Hakkari), the karyotype has $2n = 60$, $FNa = 58$ and $FN = 60$. The autosomal set contains 29 pairs of acrocentric chromosomes. Both sex chromosomes are acrocentric and the Y chromosome is smaller than the X (Table). Kefelioğlu and Kryštufek (1999) reported a new diploid number of chromosomes ($2n = 60$) in Central Anatolia

for social voles. They also stated that these specimens have very enlarged bullae anteriorly, and then described these specimens as a new species, *Microtus anatolicus*. However, the specimens of Özalp and Yüksekovalı have smaller mastoid parts of tympanic bulla than *M. anatolicus* and show the typical cranial features of *M. socialis*. Golenishchev et al. (2002) recorded *Microtus schidlovskii* from Caucasia. They also stated that this species has $2n = 60$, and belongs to the *socialis* group of voles. When the geographical proximity and karyotype are taken into consideration, it can be concluded that Turkish specimens with $2n = 60$, $FNa = 58$ are *Microtus schidlovskii*.

Meriones tristrami Thomas, 1892

The specimen from Doğubeyazıt has $2n = 72$, $FNa = 82$ and $FN = 84$. Autosomes consist of 12 meta/submetacentric chromosomes and 58 acrocentric chromosomes. The X is large acrocentric and the Y is small acrocentric (Table).

The karyotype of Iranian specimens is composed of $2n = 72$, $FNa = 78$ and $FN = 80$. There are 4 metacentric and 4 submetacentric chromosomes in the autosomal chromosome set. The others are acrocentric. Sexual chromosomes are both acrocentric and X is larger than Y (Table). Three different FN values for *M. tristrami* were formerly reported from Turkey: FN = 76 for *Meriones tristrami blackleri*, FN = 78 for *Meriones tristrami kilensis* and FN = 82 for *Meriones tristrami lycaon* (Yiğit et al., 1998a; Yiğit and Çolak, 1998). These findings support the idea that there are variations in the chromosome arms in this geographical race of *M. tristrami*.

Meriones vinogradovi Heptner, 1931

The karyotype of the specimen from Doğubeyazıt (Coll. No. 4404) showed $2n = 44$, $FNa = 74$ and $FN = 77$. The autosomal complement is composed of 16 pairs of biarmed chromosomes (5 large, 8 varying in size from medium to small, and 3 small) and 5 pairs of acrocentric chromosomes varying in size from medium to small. The X chromosome is large and metacentric and the Y chromosome is small and acrocentric (Table). According to Matthey (1954), Nadler and Lay (1967) and Vorontsov and Korobitsina (1970), this species has similar $2n = 44$

but different FN (FN = 74 and 78). Their findings are different from the FN of Turkish specimens, and support the idea that there is a chromosomal variation in the FN of this species.

Meriones libycus (Lichtenstein, 1823)

The karyotype of specimens from Zanjan, Iran (Coll. No. 4381) has $2n = 44$, $FNa = 76$ and $FN = 80$. The karyotype consists of 7 large, 7 middle-sized and 3 small pairs of metacentric or submetacentric autosomes, 4 pairs of acrocentric autosomes, a large metacentric X chromosome and a relatively small acrocentric Y chromosome (Table). Matthey (1957) and Qumsiyeh et al. (1986) stated that FN is 74 in Iranian and Jordanian specimens. Benazzou et al. (1982) also reported $2n = 44$ for Iranian specimens with different FN. These findings supported the idea that there are karyological variations in the chromosomal arms of this species.

Apodemus iconicus (Heptner, 1948)

Based on the criteria given by Filippucci et al. (1996) and Krystufek (2002), the specimen was assigned to *A. iconicus*. The diploid number of chromosomes is $2n = 48$, $FNa = 46$ and $FN = 48$. The karyotype is composed of 46 acrocentric chromosomes in *A. iconicus* from Zanjan, Iran. The X and Y chromosomes are acrocentric. The X chromosome is larger than the Y chromosome (Table). The karyological values of Iranian specimens are consistent with Turkish specimens (Çolak, 2003).

Mus domesticus (Rutty, 1772)

According to Orsini et al. (1983), Auffray et al. (1990) and Cucchi et al. (2005), ZI value is a distinguishing characteristic among *Mus* species. ZI (0.25 to 0.46) and $T \times 100 / HB$ (0.73 to 1.0) were reported in the northwest population of *Mus domesticus* in Turkey (Gözcelioğlu et al., 2005). The *Mus* specimens from Doğubeyazıt and Zanjan were assigned to *M. domesticus* by means of the values of ZI and T / HB . The karyotypes of *M. domesticus* from Doğubeyazıt and Zanjan are $2n = 40$, $FNa = 38$ and $FN = 40$. The chromosomal set has 20 pairs of acrocentric chromosomes. The Y chromosome is smaller than the X chromosome (Table).

References

- Auffray, J.C., Tchernov, E., Bonhomme, F., Heth, G., Simson, S. and Nevo, E. 1990. Presence and ecological distribution of *Mus "spretoides"* and *Mus musculus domesticus* in Israel. Circum-Mediterranean vicariance in the genus *Mus*. Mammalian Biology, 55: 1-10.
- Benazzou, T., Viegas-Pequignot, E., Petter, F. and Dutrillax, B. 1982. Phylogenie chromosomique de quatre espèces de *Meriones* (Rongeur, Gerbillidae). Ann. Genet. 25, 19-24.
- Coşkun, Y. 1999. New karyotype of the mole rat *Nannospalax ehrenbergi* from Turkey. Folia Zool. 48: 313-314.
- Coşkun, Y. 2003. A study on the Morphology and Karyology of *Nannospalax nehringi* (Satunin, 1898) (Rodentia: Spalacidae) from Northeast Anatolia, Turkey. Turk J Zool. 27: 171-176.
- Cucchi, T., Vigne, J.D. and Auffray, J.C. 2005. First occurrence of the house mouse (*Mus musculus domesticus* Schwarz & Schwarz, 1943) in the Western Mediterranean: a zooarchaeological revision of subfossil occurrences. Biol. J. Linn. Soc. 84: 429-445.
- Çolak, E., Kivanç, E. and Yiğit, N. 1994. A study on taxonomic status of *Allactaga euphratica* Thomas, 1881 and *Allactaga williamsi* Thomas, 1897 (Rodentia: Dipodidae) in Turkey. Mammalia 58: 591-600.
- Çolak, E., Kivanç, E. and Yiğit, N., 1997a. *Allactaga elater aralyicensis* (Satunin, 1901)' in Taksonomik Durumu ve Yayılışı. Tr. J. of Zoology 21: 355-360.
- Çolak, E., Yiğit, N., Sözen, M. and Özkurt, Ş. 1997b. Distribution and taxonomic status of the genus *Microtus* (Mammalia: Rodentia) in south-eastern Turkey. Israel J. Zoology 43: 391-396.
- Çolak, E., Yiğit, N., Özku̇rt, Ş. and Sözen, M. 1997c. Karyotype of *Clethrionomys glareolus* (Schreber, 1780) in Turkey. Tr. J. of Zoology 21: 123-125.
- Çolak, E., Yiğit, N., Sözen, M. and Özkurt, Ş. 1998. A study on taxonomic status of *Microtus subterraneus* (de Selys & Longchamps, 1836) and *Microtus majori* Thomas, 1906 (Mammalia: Rodentia), in Turkey. Tr. J. of Zoology 22: 119-129.
- Çolak, E. and Yiğit, N. 1998. A new subspecies of Jerboa from Turkey *Allactaga euphratica kivanci* subsp. n. Tr. J. of Zoology 22: 93-98.
- Çolak, E. 2003. A study of *Apodemus iconicus* Heptner, 1948 (Mammalia: Rodentia) in Turkey. Turk. J. Zool. 27: 61-63.
- Çolak, E., Yiğit, N. and Verimli, R. 1999. On the karyotype of the long-clawed mole rat, *Prometheomys schapochnikovi* Satunin, 1901 (Mammalia: Rodentia) in Turkey.- Z. Säugetierkunde. 64: 1-2.
- Çolak, R., Çolak, E. and Yiğit, N., 2005. Morphometric, karyotypic and electrophoretic analysis of the genus *Apodemus* Kaup, 1826 (Mammalia: Rodentia) in Thrace, Turk. J. Zool., 29: 147-153.
- Doğramacı, S. Kefelioğlu, H. and Gündüz, İ. 1994. Anadolu *Mesocricetus* (Mammalia: Rodentia) Türlerinin Karyolojik Analizi. Tr. J. of Zoology 18: 41-45.
- Doğramacı, S. and Kefelioğlu, H. 1991. Karyotype of *Apodemus* spp. (Mammalia: Rodentia) in Anatolia. Doğa - Tr. J. of Zoology 15: 46-52.
- Filippucci, M.G., Storch, G. and Macholán, M. 1996. Taxonomy of genus *Sylvaemus* (Mammalia: Rodentia) in western Anatolia: morphological and electropheretical evidence. Senckenbergiana Biologica 75: 1-14.
- Gharkhloo, M. and Kivanç, E. 2003. A study on the morphology, karyology and distribution of *Ellobius* Fisher, 1814 (Mammalia: Rodentia) in Iran. Turk. J. Zool. 27: 281-292.
- Golenishchev, F.N., Sablina, O.V., Borodin, P.M. and Gerasimov, S. 2002. Taxonomy of voles of the subgenus *Sumeriomys* Argyropula, 1933 (Rodentia, Arvicolinae, Microtus), Russian Journal of Theriology. 1: 43-55.
- Gözcelioglu, B., Çolak, R., Çolak, E. and Yiğit, N. 2005. A study on *Mus domesticus* Rutty, 1772 and *Mus macedonicus* Petrov and Ruzic, 1983 (Mammalia: Rodentia) distributed along the line of Ankara, Bolu and Zonguldak. Turk. J. Zool. 29: 133-140.
- Gülkaç, M.D. and Yüksel, E. 1989. A cytogenetical study on blind mole rats around Malatya province. DOĞA TU. Biyol. D. 13: 63-71 (in Turkish).
- Ivanitskaya, E., Coşkun, Y. and Nevo, E. 1997. Banded karyotypes of mole rats (Spalax, Spalacidae, Rodentia) from Turkey: a comparative analysis. J Zool. Syst. Evol. Res. 35: 171-177.
- Kefelioğlu, H. 1995. The taxonomy of the genus of *Microtus* (Mammalia: Rodentia) and its distribution in Turkey. Tr. J. of Zoology 19: 35-63.
- Kefelioğlu, H. and Kryštufek, B. 1999. The taxonomy of *Microtus socialis* group (Rodentia: Microtinae) in Turkey, with the description of a new species. Journal Nat. Hist., 33: 289-303.
- Kivanç, E., Sözen, M., Çolak, E. and Yiğit, N. 1997a. Karyologic and phallic aspects of spiny mouse, *Acomys cahiricus* Spitzemberger, 1978 (Rodentia: Muridae) in Turkey. Tr. J. of Zoology 21: 167-169.
- Kivanç, E., Sözen, M., Çolak, E. and Yiğit, N. 1997b. Karyologic and phallic characteristics of *Dryomys laniger* Feiten and Storch, 1968 (Rodentia: Gliridae) in Turkey. Isr. J. Zool. 43: 401-403.
- Kryštufek, B. 2002. Identity of four *Apodemus* (*Sylvaemus*) types from the eastern Mediterranean and the Middle East. Mammalia 66: 43-51.
- Matthey, R. 1954. Recherches sur les chromosomes des Muridae. Caryologia Pisa 6: 1-44.
- Matthey, R. 1957. Cytologie et taxonomie du genre *Meriones* Illiger (Rodentia - Muridae - Gerbillinae). Säugetierkd. Mitt. 5, 145-150.
- Nadler, C.F. and Lay, D.M. 1967. Chromosomes of some species of *Meriones* (Mammalia: Rodentia). Z. Säugetierkunde, 32: 285-291.
- Nevo, E., Filippucci, M.G., Redi, C., Korol, A. and Beiles A. 1994. Chromosomal speciation and adaptive radiation of mole rats in Asia Minor correlated with increased ecological stress. Proc. Natl. Acad. Sci. USA 91: 8160-8164.

- Orsini, P., Bonhomme, F., Britton-Davidian, J., Croset, H., Gerassimov, S. and Thaler, L. 1983. Le complexe d'espèces du genre *Mus* en Europe Centrale et Orientale. Z. Säugetierkunde, 48: 86-95.
- Özkan, B., Yiğit, N. and Colak, E. 2003. A study on *Micromys minutus* (Pallas, 1771) (Mammalia: Rodentia) in Turkish Thrace. Turk. J. Zool. 27: 55-60.
- Özkurt, Ş., Çolak, E., Yiğit, N., Sözen, M. and Verimli, R. 1999. Contribution to karyology and morphology of *Arvicola terrestris* (Lin., 1758) in Central Anatolia. Tr. J. of Zoology, 23: 253-257.
- Özkurt, Ş., Yiğit, N. and Çolak, E. 2002. Karyotype variation in Turkish populations of *Spermophilus* (Mammalia: Rodentia). Z. Säugetierkunde, 67: 117-119.
- Qumsiyeh, M.B., Schlitter, D.A. and Disi, A.M. 1986. New records and karyotypes of small mammals from Jordan. Z. Säugetierkunde 51: 139-146.
- Soldatovic, B. and Savic, I. 1978. Karyotypes in some populations of the genus *Spalax* (Mesospalax) in Bulgaria and Thrkey. Säugetierkdl. Mitt. 26: 252-256.
- Sözen, M., Çolak, E. and Yiğit, N. 2000. Contributions to the karyology and taxonomy of *Spalax leucodon nehringi* Satunin, 1898 and *Spalax leucodon armeniacus* Mehely, 1909 (Mammalia: Rodentia) in Turkey. Z. Säugetierkunde 65: 309-312.
- Vorontsov, N.N. and Korobitsina, K.V. 1970. Materials on a comparative karyology of Gerbillinae. Tziologiya, 12: 152-157.
- Yiğit, N., Çolak, E., Kivanç, E. and Sözen, S. 1997. A new gerbil from Turkey *Gerbillus dasyurus* Wagner, 1836 (Rodentia: Gerbillinae). Isr. J. Zool. 43: 13-18.
- Yiğit, N., Kivanç, E., and Çolak, E. 1998a. On the taxonomic status of *Meriones tristrami* Thomas, 1892 (Rodentia: Gerbillinae) in Turkey. Zoology in the Middle East 16: 19-30.
- Yiğit, N., Kivanç, E. and Çolak, E. 1998b. Contribution to taxonomy and karyology of *Meriones meridianus* (Pallas, 1773) and *Meriones crassus* Sundevall, 1842 (Rodentia: Gerbillinae) from Turkey. Z. Säugetierkunde 63: 311-314.
- Yiğit, N., Çolak, E., Sözen, M. and Özkurt, Ş. 1998c. The taxonomy and karyology of *Rattus norvegicus* (Berkenhout, 1769) and *Rattus rattus* (Linnaeus, 1758), in Turkey. Tr. J. of Zoology 22: 203-212.
- Yiğit, N. and Çolak, E. 1998. A new subspecies of *Meriones tristrami* Thomas, 1892 (Rodentia: Gerbillinae) from Kilis (Southeastern Turkey): *Meriones tristrami kilensis* subsp. n. Tr. J. of Zoology 22: 99-104.
- Yiğit, N. and Çolak, E. 1999. A study of the taxonomy and karyology of *Meriones persicus* (Blanford, 1875) (Mammalia: Rodentia) in Turkey. Tr. J. of Zoology 23: 269-274.
- Yiğit, N., Çolak, E., Sözen, M., Özkurt, Ş. and Verimli, R. 2000a. The distribution, morphology, and karyology of the genus *Mesocricetus* (Mammalia: Rodentia) in Turkey. Folia Zool. 49: 167-174.
- Yiğit, N., Verimli, R., Sözen, M., Çolak, E. and Özkurt, Ş. 2000b. The karyotype of *Apodemus agrarius* (Pallas, 1771) (Mammalia: Rodentia) in Turkey. Zoology in the Middle East 20: 21-23.
- Yiğit, N., Çolak, E., Verimli, R., Özkurt, Ş. and Sözen, M. 2001. A study on the distribution, morphology and karyology of *Tatera indica* (Hardwicke, 1807) (Mammalia: Rodentia) in Turkey. Turk. J. Zool. 25: 67-70.