

## Community Structure of Zooplanktonic Organisms in Lake Akşehir\*

Nuray (EMİR) AKBULUT

Hacettepe University, Faculty of Science, Department of Biology - Hydrobiology 06532-Beytepe, Ankara - TURKEY

Received: 13.01.1998

**Abstract:** In this study, the abundance and seasonal distribution of rotifer and crustacean zooplankton were examined. A negative correlation between rotifers and crustaceans in terms of their abundance rates was observed. Copepods were most abundant and comprised 43.3 %; rotifers comprised 34 % and cladocerans comprised 22.6% of the total zooplanktonic organisms. Copepods were abundant in autumn whereas rotifers were abundant in spring and autumn but cladocerans were abundant in summer periods.

Some zooplanktonic organisms recorded in all samples throughout the study period were *Filinia*, *Hexarthra*, *Brachionus*, *Asplanchna*, *Keratella*, *Arctodiaptomus spinosus* and *Diaphanosoma lacustris*.

**Key Words:** Rotifera, Cladocera, Copepoda, Biomass, Seasonal distribution

### Akşehir Gölü Zooplanktonik Organizmalarının Kommunite Yapısı

**Özet:** Bu çalışmada, rotiferlerin ve kabuklulara ait zooplanktonik organizmaların mevsimsel dağılımları ve bollukları incelenmiş ve aralarında negatif bir korelasyon olduğu saptanmıştır.

Toplam zooplanktonun sayısal olarak % 43.3 ünü Copepoda, % 34 üünü Rotifera ve % 26 sini Cladocera oluşturmaktadır. Copepodlar genellikle sonbaharda, rotiferler İlkbahar ve sonbaharda yoğunken, kabuklular yaz aylarında yoğun olarak gözlenmiştir.

Tüm örneklemde dönemi boyunca gölde tespit edilen zooplanktonik organizmalardan bazıları şunlardır. *Filinia*, *Hexarthra*, *Brachionus*, *Asplanchna*, *Keratella*; *Arctodiaptomus spinosus* ve *Diaphanosoma lacustris*.

**Anahtar Sözcükler:** Rotifera, Cladocera, Copepoda, Biyomas, Mevsimsel Dağılım

### Introduction

Although many recent studies have appeared concerning zooplankton taxonomy in Turkey (1-8), there is a great need for ecological studies of lake ecosystems.

Lake Akşehir is shallow and large, and is an "A" class wetland of central Anatolia. The lake is very important for ornithology, fisheries and the native crayfish. The lake is connected to lake Eber by Taşköprü stream. Recently the water color and structure of the lake have been affected both by industrial pollution from the alkohol factory located near lake Eber and sewage effluent of Akşehir.

The aim of this paper is to report the short term changes in the species composition and seasonal distribution of zooplanktonic organisms between 1991 and 1992.

### Study area

Coordinates: 38° 31'N-31° 28'E

Altitude: 960 m

Surface Area: 33500 ha

Lake Akşehir is located between the Sultan and Emir Mountains in central Anatolia. The lake is fed mainly by Taşköprü stream, lake Eber, Adıyan and Engelli streams and rain as well snow water (9-12). (Figure 1).

During the study period, the following macrophytes species were observed: *Ceratophyllum demersum*, *Phragmites australis*, *Potamogeton pectinatus*, *Bulboschoenus maritimus*, *Tamarix* sp., *Cirsium alatum*, *Pseudocretecum* sp., *Galega officinalis*, *Plantago lanceolata*, *Najas marina*, and *Ranunculus saniculifolius*.

\* This study is a part of PhD dissertation




Figure 1. Sampling area

The following phytoplanktonic organisms appear most during the surveys: *Achnanthes*, *Cocconeis*, *Cyclotella*, *Cymbella*, *Diatoma*, *Epithemia*, *Fragillaria*, *Gomphonema*, *Gyrosigma*, *Caloneis*, *Asterionella*, *Amphora* (Bacillariophyta); *Anabaena*, *Chrococcus*, *Lygbya*, *Oscillatoria*, *Merismopedia* (Cyanophyta); *Closterium*, *Cosmarium*, *Oocystis*, *Euastrum*, *Chlamydomonas* (Chlorophyta); *Ceratium*, *Peridinium* (Dinophyta); *Euglena*, *Phacus*, *Trachelomonas* (Euglenophyta); *Cryptomonas* (Cryptophyta)

The fishes observed in the lake in 1991 were as follows *Cyprinus carpio*, *Esox lucius*, *Leuciscus cephalus*, *Leuciscus lepidus*, *Alburnus orontis*, *Gobio gobio*, *Cobitis taenia*, *Neomachilus angorae* and *Cyprinus carpio* (11). However during the sampling period the most abundant organisms were *Cyprinus carpio*, *Esox lucius* and *Leuciscus cephalus*.

## Materials and Methods

During this study, samples were collected monthly from three stations using a plankton net of 44 µm mesh

size and a Patalas sampler with a volume of 3.5 L and hauling through the water column. The samples were fixed with formaldehyde in 4 % and then, they were concentrated. One to ten ml of each sample were counted until 200 individuals at least or 10 % of the sample volume were enumerated using an inverted microscope (13) and 95 % confidence limits were fitted to calculate the mean population estimates. Rotifera and Crustacean species were identified from the literature (14-20).

During the sampling period, dissolved oxygen, conductivity, salinity, pH, temperature and Secchi depth were measured in the pelagic zone of the lake using portable instruments (YSI 33 SCT meter, YSI 51 B oxygen meter and pH meter).

## Results

The rotifera species recorded from the lake, are given below. The crustacean species and genera are given in Table 2. The zooplanktonic community of the lake is composed of mainly Copepoda, Rotifera and Cladocera. A total of 36 Rotifera, 2 Copepoda and 6 Cladocera species were

identified. Copepoda comprised 43 %, Rotifera 34 % and Cladocera 22.6 % of the total zooplankton density.

The measured physical parameters are given in Table 1, the seasonal distribution of the zooplankton of Lake Akşehir is given in Table 2 and also in Figures 2-3.

#### Identified Rotifera species

##### 1. Fam.: Brachionidae

1. *Brachionus quadridentatus* HERMANN, 1783
2. *Brachionus plicatilis* (O. F. M., 1986)
3. *Brachionus urceolaris* (O. F. M., 1773)
4. *Brachionus rubens* EHRENBERG, 1838
5. *Brachionus calyciflorus* PALLAS, 1766
6. *Brachionus angularis* GOSSE, 1851
7. *Keratella quadrata* (O. F. M., 1786)
8. *Keratella cochlearis* (GOSSE, 1851)
9. *Anuraeopsis fissa* (GOSSE, 1851)

##### 2. Fam.: Euchlanidae

10. *Euchlanis dilatata* EHRENBERG, 1832

##### 3 Fam.: Trichotriidae

11. *Trichotria pocillum* (O. F. M., 1773)

##### 4 Fam.: Colurellidae

12. *Colurella colurus* (EHRENBERG, 1830)
13. *Colurella uncinata* (O. F. M., 1773)

Table 1. Measured physical parameters in Lake Akşehir

Date	T(°C)	O <sub>2</sub> (mg/L)	EC µS/cm	Salinity (‰)	pH	Depth (m)	Secchi D.(cm)
5/9/90	20.5	7.8	597	0.2	9	3	35
30/10/90	13	8.5	3000	0.2	8	2	30
25/12/90	3	13.2	1600	2	9	2	35
4/5/91	26	7.4	4780	3.25	9.5	3.25	50
8/6/91	25	7.8	4500	3	9.5	2.4	30
24/8/91	21.5	7.8	4200	2.5	9.5	2.25	30
16/11/91	23	7.8	3800	2.5	9	2.2	30
30/11/91	6.5	7.3	2850	2.3	8.7	2	30
18/4/92	15	8.3	3100	2.5	8.35	2	25
26/6/92	15	8.3	4200	4	9.5	2.8	40
12/7/92	22	8.9	3880	2.5	7.68	3	30
1/8/92	19.7	8.7	3900	2.5	9.56	2.8	25
6/8/92	13	8.9	3800	2.3	10	3	33
29/8/92	22.7	9	4350	2.5	8.72	2.8	30
10/10/92	15.5	9.6	3900	2.9	8.6	2.5	35

##### 14. *Lepadella patella* (O. F. M., 1786)

##### 5 Fam.: Lecanidae

15. *Lecane luna* (O. F. M., 1776)
16. *Lecane (M.) lamellata* (DADAY, 1893)
17. *Lecane nana* (MURRAY, 1913)
18. *Cephalodella gibba* (EHRENBERG., 1838)

##### 6. Fam.: Gastropodidae

19. *Ascomorpha saltans* (BARTSCH, 1870)
20. *Ascomorpha ovalis* (BERGENDAHL, 1892)

##### 7. Fam.: Synchaetidae

21. *Synchaetae pectinata* EHRENBERG, 1832
22. *Synchaeta oblonga* EHRENBERG, 1831
23. *Polyarthra vulgaris* CARLIN, 1943
24. *Polyarthra dolichoptera* IDELSON, 1925

##### 8. Fam.: Asplanchnidae

25. *Asplanchna priodonta* GOSSE, 1850
26. *Asplanchna (A). girodi* (DE GUERNE, 1888)
27. *Asplanchna (A) sieboldi* (LEYDIG, 1854)

##### 9. Fam.: Testudinellidae

28. *Testudinella patina* (HERMANN, 1783)
29. *Pompholyx complanata* GOSSE, 1851

Community Structure of Zooplanktonic Organisms in Lake Akşehir

Table 2. Counted Zooplanktonic Organisms of Lake Akşehir in Different Sampling Dates (ind/m<sup>3</sup>)

	5/9/90	25/12/90	4/5/91	8/6/91	24/8/91	12/10/91	16/11/91
<b>COPEPODA</b>							
Calanoid Copepoda	186(131-259)	2290(929-5632)	3801(762-18927)	6456(1573-26468)	75(24-146)	2382(2246-2523)	23547(8611-69957)
Copepodit	316(126-779)	19860(8205-48060)	9332(2696-32287)	2041(852-4876)	145(135-152)		11220(6130-20531)
Nauplius	309(73-1275)	191(33-1048)	22908(20093-26114)	4786(778-29385)	537(493-654)	6280(5868-6718)	17021(10637-27232)
Cyclopoid Copepoda		76(5-956)				4415(4012-4855)	12023(212-678096)
<b>CLADOCERA</b>							
Diaphanosoma lacustris		234(116-467)		698(390-1190)	1788(1040-3056)		6166(581-65285)
Other Cladocera species	1348(1035-1751)	234(64-832)	36507(547-427714)	6918(3997-11967)		10964(10245-11730)	
<b>ROTIFERA</b>							
<i>Brachionus calyciflorus</i>	56(7-389)					56(96-127)	
<i>B. urceolaris</i>	112 (96-127)				46(14-133)		
<i>B. angularis</i>			38018(10472-138004)	4897(693-34522)			
<i>B. quadridentatus</i>			1999(337-11793)	1809(103-29305)	1590(453-5564)		
<i>B. plicatilis</i>		33(19-51)	2884(1601-5190)		235(62-707)		
<i>Keratella quadrata</i>	76 (19-280)		314(61-1561)	954(301-2305)	24(11-42)	105(26-420)	
<i>K. cochlearis</i>	4 (0-32)						
<i>Anuraeopsis fissa</i>			758(226-2523)				
<i>Lecane luna</i>	118 (110-124)	30(4-155)	867(273-2338)				
<i>Hexarthra</i> spp.	73 (20-237)		95(30-325)	118(28-495)	75(25-225)	3530(1176-10589)	
<i>Filinia</i> spp.			11841(4320-32443)	9388(825-13899)	3100(3096-10850)	1540(549-4311)	
<i>Ascomorpha</i> spp.			225(179-280)				
<i>Lecane</i> nana			227(89-568)				
<i>L. lamellata</i>			83(17-380)				7586(2605-22073)
<i>Lepadella patella</i>			225(179-280)				21887(13678-35237)
<i>Asplanchna</i> spp.				152(58-385)		158(30-837)	324(65-1577)
<i>Euchlanis</i> sp.				45(12-145)			
<i>Polyarthra</i> sp.				27(6-98)			
	30/11/91	18/4/92	26/6/92	12/7/92	1/8/92	29/8/92	10/10/92
<b>COPEPODA</b>							
Calanoid Copepoda (Female)	11749(916-150384)	1072(770-1489)	1318(539-3214)	1698(811-3547)	73(47-108)	776(83-7310)	380(120-1188)
Calanoid Copepoda (Male)	11482(6413-20550)	243(156-347)		275(67-1101)	22(9-43)	200(80-478)	138(114-163)
Copepodit	18543(1243-306824)	1905(1189-3047)	2754(586-12915)	8128(4299-1536)	53(26-103)	2138(135-33565)	288(235-350)
Nauplius	19634(12270-31412)	10715(10401-11035)	1120(639-1939)	5248(3915-7031)	457(327-634)	1517(229-9965)	1585(953-2629)
<b>CLADOCERA</b>							
Diaphanosoma lacustris	10750(8268-13974)	5210(1487-18234)		5012(2397-10474)	134(116-151)	2137(89-50218)	331(157-690)
Other Cladocera species					2310(1540-3464)		
<b>ROTIFERA</b>							
<i>Brachionus calyciflorus</i>		43(8-190)					
<i>Keratella quadrata</i>		570(90-3010)				6(0-99)	
<i>Hexarthra</i> spp.	2495(795-7806)	1412(1044-1905)		724(424-1229)	145(102-199)	660(240-1670)	2570(1316-5010)
<i>Filinia</i> spp.	6025(3089-11748)	680(140-3263)	1385(553-3461)	15136(6333-36174)	31(17-51)	3890(2945-5133)	660(634-686)
<i>Asplanchna</i> spp.		3236(2327-4497)	76(11-463)	275(224-334)		224(202-245)	62(37-97)
<i>Brachionus angularis</i>				38(8-190)	78(62-93)		

95% Confidence limits of counted zooplanktonic organisms are shown in parentheses

COPEPODA	7/11/92
Calanoid Copepoda (Female)	1445(1362-1530)
Calanoid Copepoda (Male)	426(420-429)
Copepodit	1412(1014-1961)
Nauplius	1023(810-1287)
<b>CLADOCERA</b>	
Diaphanosoma lacustris	660(256-1688)
Other Cladocera species	
<b>ROTIFERA</b>	
<i>Brachionus calyciflorus</i>	
<i>Keratella quadrata</i>	
<i>K. cochlearis</i>	
<i>Hexarthra</i> spp.	
<i>Hexarthra</i> spp.	1690(712-4004)
<i>Filinia</i> spp.	580(85-3885)
<i>Asplanchna</i> spp.	368(16-7579)
<i>Brachionus angularis</i>	




Figure 2. Seasonal distribution of zooplankton during the sampling period




Figure 3. Seasonal distribution of rotifers during the sampling period

10. Fam.: Conochilidae

30. *Conochilus natans* (SELIGO, 1900)

11. Fam.: Hexarthridae

31. *Hexarthra mira* (HUDSON, 1871)

32. *Hexarthra fennica* (LEVANDER, 1892)

12. Fam.: Filiniidae

33. *Filinia longiseta* (EHRENBURG, 1834)

34. *Filinia terminalis* (PLATE, 1886)

35. *Filinia pejleri* HUTCHINSON, 1964

13. Fam.: Philodinidae

36. *Philodina megalotrocha* EHRENBURG, 1832

Identified Cladocera and Copepoda Species

*Moina macrura*

*Daphnia longispina*

*Diaphanosoma lacustris*

*Alona rectangula*

*Chydorus sphaericus*

*Bosmina longirostris*

*Scaphaloberis mucronata*

*Arctodiaptomus spinosus*

*Acanthodiaptomus denticornis*

*Cyclops* sp.

Among crustacean species *Arctodiaptomus spinosus* and *Acanthodiaptomus denticornis* had high density within the total zooplanktonic community. These species were found at all the stations throughout the year. Especially in autumn periods, they reached maximum abundance. Among cladoceran, *Diaphanosoma lacustris* was dominant. This species reached maximum density in summer and autumn periods (14454 ind./m<sup>3</sup>). The total number of copepods varied from 148 ind./m<sup>3</sup> in August 1992 to 51000 ind./m<sup>3</sup> in October 1991. Whereas the total number of Cladocera varied from 331 ind./m<sup>3</sup> in October 1992 to 36507 ind./m<sup>3</sup> in May 1991.

Among rotifers, *Brachionus angularis*, *B. quadridentatus*, *B. plicatilis*, *Keratella quadrata*, *Hexarthra*, *Filinia* and *Asplanchna* were abundant organisms. The highest percentages of rotifers was found in May, June and November 1991, July 1992 and July 1993. The total number of

rotifers varied from 63 ind./m<sup>3</sup> in December 1990 to 57536 ind./m<sup>3</sup> in May 1991 (Figure 3).

Although the horizontal distribution of the zooplanktonic organisms was nearly uniform, usually copepods and rotifers were dominant at the first and second stations, while cladocerans were abundant at the third station.

During the spring period, some rotifers such as *Brachionus angularis* were the most abundant species at the first station, *Brachionus quadridentatus* at the first and second stations, *B. plicatilis* at the second and third stations, *Anuraeopsis fissa* at the first, *Filinia longiseta* at the second and third stations, *Lecane luna* at the second, *L. nana* at the first, *Keratella quadrata* at the first and *Diaphanosoma lacustris* at the second and third stations. In this period Copepoda had 1850 individuals per m<sup>3</sup> at the first station, 11418 individuals per m<sup>3</sup> at the second station and 2753 ind./m<sup>3</sup> at the third station (Figures 2-3).

In June 1991, phytoplanktonic organisms were dominant at the first station in parallel to the high number of the rotifers of *Brachionus angularis* (18890), *B. quadridentatus* (4934), *Filinia* (9163) and *Keratella quadrata* (1410) individuals per m<sup>3</sup>. At the same time, Copepoda at the first station had 16916 individuals, at the second station 3392 individuals and at the third station 4832 individuals per m<sup>3</sup> respectively. It was observed that larger cladocerans were abundant at the second and third stations. Among rotifers, *Filinia* and *Asplanchna* were abundant at the second and third stations also. In summer periods cladocerans were more abundant than copepods and rotifers respectively.

In Lake Akşehir the clear water period was observed in May (21). The clear water period is caused by grazing herbivorous zooplankton (22,23). In Lake Akşehir an early peak of small, rapidly growing algae was followed by a short period of clear water and transparency in mid May. During this period Secci depth was measured as 50 cm, which was the highest value for the lake (Table 1). In the summer period, large algae and blue green algae developed after the clear water phase (8). It was observed that in Elmacı, (21) the same results were obtained during the period between June 1992 and November 1993. The decline in algal populations could be a consequence of nutrient limitation, sedimentation and the grazing effect of large bodied grazers.

During the study period, among phytoplankton, Bacillariophyta were dominant in terms of the species diversity and frequency; Cyanophyta and Chlorophyta were also dominant but the members of other divisions were not numerous in Lake Akşehir.

## Discussion

Most of the studies dealing with this topic have shown that the increase in nutrients and the corresponding higher phytoplankton biomass result in higher zooplankton biomass (24-29). In Lake Akşehir zooplankton biomass was higher in the spring and early summer periods when the nanoplanktonic algae were dominant.

In the eutrophic lakes, herbivorous zooplanktonic organisms can effectively reduce the phytoplankton biomass vulnerable to grazing, at the beginning of the growing season (24,26). After an early spring pulse a period of low phytoplankton standing crop is usually observed, caused most likely by high grazing pressure exerted by large bodied filter feeding herbivorous species. However, community grazing rates can vary seasonally and among sites. Community grazing rates were positively related to zooplankton biomass and the size of organisms in a community (30-35).

Also, interactions between invertebrate predators and their preys are reflected in quantitative changes in community dynamics (32) and planktivorous fish control the zooplankton populations especially in the summer months (33). In the presence of planktivorous fish, the proportion of cladocerans decreases and phytoplankton as well as bacteria, flagellates and rotifers are able to proliferate (34-35). Hurlbert (35) observed that lakes with fish usually have a sparse zooplankton dominated by cyclopoid copepods and chydorid cladocerans; the opposite conditions tend to have an abundant zooplankton dominated by calanoid copepods and daphnids. In Lake Akşehir the same result was observed as the calanoid copepods were higher in number than cyclopoids.

Herzig (28) noted that strong wind conditions on the lake and the resulting turbulences cause a mechanical stress on the soft-bodied Cladocera. Among rotifers, *Hexarthra* is also sensitive to temperature and wind

action. Herzig and Koste (36) also observed that low temperature and high amount of wind generated suspended particles which can negatively influence the development of *Hexarthra* population and of several cladocerans (36). In Lake Akşehir, strong wind action might affect the horizontal distribution of the organisms as is the case in other lakes in the area (8).

In deep and stratified lakes, more phytoplankton cause higher densities of zooplankton. The changes in zooplankton abundance, fecundity and size might have been closely correlated with the amount of planktonic algea (37). In such lakes phytoplankton is the basic and frequent food source of the zooplankton community and higher phytoplankton biomass results in higher zooplankton biomass. However, many scientists have found that zooplankton feed on detritus (38,39). In Neusiedlersee it was shown that *Arctodiaptomus spinosus* and *Diaphanosoma brachyrum* may be found without an algal food source. In such shallow lakes the direct grazing food chain seems to be increasingly replaced by the indirect detritus food chain. The role of phytoplankton as a main food source is taken over by bacteria and detritus originating from the bottom (25,39).

In Lake Akşehir the salinity degree was between 0.2 and 4 %. In these lakes, such as Neusiedlersee, *Arctodiaptomus spinosus* was more abundant than the others. Species such as *Brachionus* spp., *Filinia* spp., *Keratella* spp., which can be regarded as the indicators of eutrophy, were present in the lake (8). Among phytoplankton, especially Diatomea such as *Camphylodiscus*, *Surirella*, *Cyclotella* and *Diatoma* were dominant and they can also be regarded as indicators of eutrophy and they occur in highly alkaline waters (40).

## Acknowledgements

The present study was made possible by TUBITAK-TBAG grant no: 933 'Çavuşcu Gölü Rotifera faunasının taxonomik ve ekolojik açıdan incelenmesi'

I am very grateful to Prof. Dr. Ali Demirsoy for guiding me in my work and supplying all my needs and Dr. hc. Walter Koste, Doz. Dr. Alois Herzig and Aydin Akbulut for their kind help with the phytoplankton investigations.

## References

1. Dumont, H. J., Rotifer from Turkey. *Hydrobiologia* 147: 65-73. 1987.
2. Emir, N., A note on four Rotifer species new to Turkey. *Biol Sb. Donaea* 57, 78-80, 1990a.
3. Emir, N., Samsun Bafra Gölü Rotatoria faunasının taksonomik yön-den incelenmesi Doğa Tr. *J. of Zoology C* 14, S1, 89-106, 1990b.
4. Segers, H., and Emir, N., Rotifera From North and Northeast Anatolia (Turkey). *Hydrobiologia* 245, 179-189, 1992.
5. Ustaoğlu, R., Zooplankton (Metazoa) of the Karagöl (Yamanlar, Izmir-TURKEY). *Biologia Callo helenica*, vol. 12, 273-281, 1986.
6. Ustaoğlu, R., Akgöl'ün (Selçuk-Izmir) Rotifer Faunası, VIII Ulusal Biyoloji Kongresi, cilt II, 614-626, 1987.
7. Emir, N., Samsun Bafra Gölü Rotatoria Türlerinin Mevsimsel Değişim Üzerine Ekolojik bir çalışma, Doğa.Tu. *Zooloji D.C.13.S.3.220-227*, 1989.
8. Emir, N., Zooplankton community structure of Çavuşcu and Eber lakes in Central Anatolia, *Acta Hydrochim. Hydrobiol.* 22,6, 280-288, 1994.
9. Altinayar, G., Eber, Akşehir sulama projesinin Eber ve Akşehir Göl-leri bitkisel ve hayvansal su ürünleri üretimi üzerindeki etkisi konusunda ön çalışmalar, 58s., 1993.
10. Anonymous, Akşehir Gölü'nün bazı limnolojik özellikleri üzerine bir araştırma, su ürünlerini yayıni 4. 40s., 1994.
11. Anonymous, Eber, Akşehir projesi çevresel etki değerlendirmesi raporu, içme suyu ve kanalizasyon Derneği Başkanlığı, Çevre sorunları ve suları kirlenmesini kontrol şube müdürlüğü 19 s., 1991.
12. Anonymous, (1992) Doğal Hayatı Koruma Derneği yayını 156 s.
13. Edmondson, W. A., Manual on methods for the assessment of secondary productivity in freshwater IBP Handbook no 17, 358 p. 1971.
14. Koste, W., Die Radetiere Mitteleuropas 1. texband Berlin Stuttgart 670 pp. 1978a.
15. Koste, W., Die Radetiere Mitteleuropas II. Tafelband Berlin Stuttgart, 235 pp., 1978b.
16. Edmondson, W. T., Freshwater Biology, University of Washington Seattle, 1248 pp., 1959.
17. Kuttikova, A., Rotatoria (The Rotifer fauna of SSCB), 670 pp., 1970.
18. Kolisko, R. A., Plankton rotifers biology and taxonomy, Biological station Lunz of the Austrian Academy of science, Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 974 p., 1971.
19. Kiefer, F., Das Zooplankton der Binnengewässer E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 380 p., 1978.
20. Herbst, V. H., Blattfusskrebsse, Gesellschaft der naturfreunde Franckh'sche verlagshandlung, Stuttgart, 120 p., 1962.
21. Stewart, J. A and Wetzel G. R., Cryptophytes and other microflagellates as couples in planktonic community dynamics. *Arch. Hydrobiol* 106, 1-19, 1986.
22. Sommer, U., Plankton Ecology Succession in plankton communities, Springer Verlag 369 p., 1989.
23. Elmaci, A., Akşehir (Konya) Gölü fitoplanktonu ve kıyı bölgesi (litoral bölge) alglerinin ekolojik ve floristik olarak incelenmesi, A. U. Doktora tezi, 122 s., 1995.
24. Lampert, W., Fleckner, W., Rai, H., Taylor B. E., Phytoplankton control by grazing zooplankton: A study on the spring clear water phase. *Limnol. Oceanogr.* 31 (3) 478-490, 1986.
25. Löffler, H., Neusiedlersee: The limnology of a shallow lake in Central Europe Dr. W. Junk publishers 280-335p., 1979.
26. Langeland, A., Reinersten, H., Interactions between phytoplankton and zooplankton in a fertilized lake. *Holoartic ecology* 5, 3, 253-271, 1982.
27. Havens, E. K., Zooplankton dynamics in a freshwater estuary. *Arch. Hydrobiol* 123, 1, 69-97, X. 1982.
28. Herzig, A., Predator prey relationship within the pelagic community of Neusiedlersee. *Hydrobiologia* 275/276, 81-96, 1994.
29. Cry, H. and Pace Michael L., Grazing by zooplankton and its relationship to community structure, *Can. J. Fish. Aquat. Sci.*, vol 49, 1455-1465, 1992.
30. Gliwicz, M. Z., Why do cladocerans fail to control algal blooms. *Hydrobiologia* 200/201, 83-97, 1990.
31. Gliwicz, M. Z., Relative significance of direct and indirect effects of predation by planktivorous fish on zooplankton. *Hydrobiologia* 272, 201-210, 1994.
32. Lair, N., Effects of Invertabrate predation on the seasonal succession of a zooplankton community: a two year study in Lake Aydat, France. *Hydrobiologia* 198, 1-12, 1990.
33. Kalikhman, I., Waline, P. and Gophen, M., Similtaneous patterns of temperature, oxygen, zooplankton and fish distribution in Lake Kinneet, Israel. *Freshwater Biology* 28, 337-347, 1992.
34. Cristofersen, K., Potential role of fish predation and natural populations of zooplankton in structuring a plankton community in eutrophic lake water, *Limnol. Oceanogr.* 38 (3), 561-573, 1993.
35. Hurlbert, H. Stuart., Fish flamingo plankton interactions in the peruvian Andes. *Limnology and Oceanography*, Volume 31, 3-7, 1986.
36. Herzig, A. and Koste, W., The development of *Hexarthra* spp. in a shallow alkaline lake. *Hydrobiologia* 186/187, 129-136, 1989.
37. Edmondson, W. T., Reproductive rate of planktonic rotifers as related to food and temperature in nature, *Ecol. Monogr.* 35, 61-111, 1965.
38. Nauwerck, A., Die beziehung zwischen zooplankton and phytoplankton im See Erken. *Symb. bot. upsal.* 17, 5, 1-163, 1963.
39. Laszlo, G. T., Katalin V. Balogh and Zankai, P. N., The significance and degree of abioeston consumption in the filter feeder *Daphnia galeata*, Sars am Richard (cladocera) in lake Balaton, *Arch. Hydrobiol* 106, 1, 45-60, 1986.
40. Yıldız, K., Şen, B., Atıcı, T., Akbulut, A., Akşehir gölü (Konya) fitoplanktonundaki diatomeler, XII. Ulusal Biyoloji Kongresi, Hidrobiyoloji Seksiyonu, Cilt IV, sf. 173-179, 1994.