

The Black Sea Zooplankton: Composition, Spatial/Temporal Distribution and History of Investigations

A. V. KOVALEV, V. A. SKRYABIN, Yu. A. ZAGORODNYAYA
Institute of Biology of Southern Seas, Nakhimov Ave., 2, Sevastopol-UKRAINE

F. BINGEL, A. E. KIDEYŞ, U. NIERMANN, Z. UYSAL
Institute of Marine Sciences, P. O. Box 28, TR 33731 Erdemli, İçel-TURKEY

Received: 09.06.1998

Abstract: Investigations into the composition, spatial and temporal distribution of the Black Sea zooplankton were reviewed in context with its historical development. The review comprises the main zooplankton studies published by the riparian countries of the Black Sea beginning with the earliest faunistic publications in the 1860s and 1870s up to the recent multinational ecosystem investigations carried out within the framework of the NATO-TU-Black Sea Project. The reviewed material covers a wide subject range of zooplankton investigations from descriptive studies to distribution and the production mechanisms in the ecosystem.

The main findings of recently performed large scale investigations into the horizontal and vertical distribution of zooplankton were also summarized and short and long term fluctuations in the composition and biomass were also analysed. Additionally, the influences of natural and anthropogenic factors on these changes were discussed and problem areas to be investigated in future were underlined.

Key Words: Black Sea, zooplankton, history, composition, distribution, fluctuation.

Karadeniz Zooplanktonu: Kompozisyonu, Yer ve Zamana göre Dağılımı ve Araştırmaların Tarihi

Özet: Karadeniz zooplanktonunun kompozisyonu, yer ve zamana göre dağılımı araştırmaların tarihi gelişmesi çerçevesinde derlendi. Bu derleme, Karadeniz'e kıyısı olan ülkelerde 1860'lı ve 1870'li yıllarda fauna tespitine yönelik yapılan ilk belli başlı yayınlardan başlayarak günümüzde NATO-TU-Karadeniz projesi çerçevesinde yapılan çok-uluslu eko-sistem araştırmalarını kapsamaktadır. Derlenen materyal, tanıtmacı çalışmalardan zooplanktonların dağılımına ve eko-sistemdeki üretim mekanizmalarına kadar uzanan çok geniş konuları içermektedir.

Zooplanktonların yatay ve dikey dağılımlarına ilişkin son yılların geniş ölçekli araştırmaları ile kompozisyon ve biyokitlede görülen kısa ve uzun süreli artma ve azalmalar da analiz edildi. Ek olarak, bu değişimlere doğal ve insan etkisinden kaynaklanan faktörlerin etkileri tartışıldı ve gelecekte yapılması gereken araştırmaların altı çizildi.

Anahtar Sözcükler: Karadeniz , zooplankton, tarihçe, kompozisyon, dağılım, artma ve azalma.

Introduction

The Black Sea is one of the most interesting seas of the world both in scientific and non-scientific aspects. Its history is full of romantic and dramatic events. Its diverse marine fauna has been influenced by the long and short term (geological, climatic, hydrological) natural and anthropogenic processes of the last three decades. Approximately 200 million people live around the Black Sea basin. However, its catchment area is much larger and numerous (waste) products from this area accumulate in this relatively small basin. Along with other human activities, they influence the Black Sea ecosystem in general and the zooplankton communities in particular.

In recent years some zooplankton species have either disappeared or become rare whilst others have flourished

and reached huge numbers (e.g., *Mnemiopsis*; Shushkina, et al., 1997). The study of qualitative and quantitative changes occurring in this ecosystem is necessary in order to understand and preserve the biodiversity.

Zooplankton, being the major consumer of the primary production, constitute the food source of organisms at higher trophic levels, some of which are of high economic value. Additionally, since the majority are filter feeders, this serves to cleanse the water column of suspended matter and hence contributes significantly to the improvement of the water quality, which is especially important in coastal zone management for recreational purposes. More than 1000 publications and the continuing interest in zooplankton studies in the Black Sea acknowledge their important role.

Of course not all publications (even, if they are thematically grouped) can be analysed in one article. However a significant number of them are taken into account for the present review, mainly dealing with the composition and distribution of zooplankton, its evolution, history of investigations and the main results of these investigations as well as future tasks to be undertaken in the Black Sea. In fact, no such review is available up to now, which covers research results of the last 150 years. Some available reviews consider only short time periods and evaluations are limited to the activities of one institution (1-4). The present review evaluates investigations starting from the first faunistic proceeding of the Sevastopol Biological Station up to the present day interdisciplinary and multi-institutional analysis of combined long-term data series of riparian countries. This latter development (establishment of a common database and joint evaluation and analysis) could be realised with the support of the NATO-Science for Stability Programme through the Institute of Marine Sciences of the Middle East Technical University of Turkey (5).

History of investigations

Investigations of the planktonic fauna in the Black Sea began in the middle of the 19th century carried out by Ukrainian scientists and were soon followed by others. The rising interest in marine life was the main reason for the subsequent foundation of marine biological stations and institutes. The Sevastopol Biological Station which was founded in 1871 was the first of its kind around the Black Sea basin. The founding and early managers (V. N. Ul'yanin; S. M. Pereyaslavtseva and A. A. Ostroumov) focused their studies on the planktonic fauna occurring in inshore areas. In 1891, A. A. Ostroumov and V. A. Karavaev (zoologists from Kiev) pioneered the offshore collection and description of dominant zooplanktonic species (6-8). Among others, the following scientists of the Sevastopol Biological Station (since 1963: Institute of Biology of the Southern Seas of the National Academy of Sciences of Ukraine) studied the zooplankton: S. A. Zernov, V. N. Nikitin, M. A. Galadzhiev, M. A. Dolgopolskaya, T. S. Petipa and V. N. Greze.

Many stations and institutes which carried out important investigations into the Black Sea zooplankton appeared in the 20th century. These were:

Ukraine-Karadag Biological Station, now Karadag Reservation. -Odessa Biological Station now Odessa branch of IBSS (Institute of Biology of Southern Seas), --

-Southern Scientific Research Institute of Marine Fishery and Oceanography, Kerch.

Russia-All-Russian Institute of Marine Fishery and Oceanography, Moscow. -Shirshov Institute of Oceanology in Moscow and its Southern Department, in Gelendzhik. -Novorossiysk Biological Station (now Biological Station of Krasnodar Univ, Novorossiysk.

Georgia-Batumi Fisheries Station.

Turkey-Hydrobiological Research Institute (now Institute of Marine Sciences and Geography)-Istanbul University. -Institute of Marine Sciences (IMS-METU), Erdemli/Içel -Institute of Marine Sciences and Technology, Izmir.

Bulgaria-Research Institute of Fisheries, Varna. - Institute of Oceanography, Varna. -Laboratory of Marine chemistry in Bourgas.

Romania-Romanian Marine Research Institute, Constanza.

In the past studies were carried out by individual scientists for personal scientific interest, however, nowadays scientists of different countries develop programs in which they combine their efforts in international programs such as the NATO-TU-Black Sea Project or the EROS 2000 program.

The subjects, contents and techniques of investigations naturally changed with time (2, 4, 9, 10). During the early studies various new species were described [6, 7, 11-14, (Chernyavskiy, 1868 and Krischagin, 1873; c.f., 15), and many others].

Zoogeographical analysis of the faunistic investigations were completed at the end the 19th century (15). Although some species and even higher taxons were revised later (e.g., 16-19). These basic taxonomic works were followed by wider and more detailed studies of the zooplankton in the Black Sea. For example, zooplankton including meroplanktonic larvae were investigated by Kiseleva, (20), Petran, (21), Murina, (22) and others, and the holoplanktonic larvae of copepoda were studied by Sazhina, (23, 24). Investigations carried out in the 1960s resulted in a series of 3 volumes concerning the "Identification of the Black Sea nad Azov Sea fauna", the main contributors being Braiko et al. (25-27). However, faunistic research is still being carried out even today (28-32). One direction within such research is the Mediterraneanization of the Black Sea fauna.

Parallel to the faunistic investigations, during nearly the same time period, ecological studies were started by

Zernov (33, 34). He investigated the quantitative distribution of zooplankton and its seasonal changes off Sevastopol. Following this, intensified research activities took place in several riparian countries of the Black Sea (e.g., 35-53 and many others). Meanwhile some authors characterised zooplankton communities of different regions including open waters and various depths (e.g., 54-57). V. N. Nikitin (35, 36) was the pioneering scientist who first described the vertical distribution of zooplankton in the open waters of the Black Sea. He also investigated features of diurnal and seasonal vertical migrations of the dominant planktonic species and demonstrated the relationship between diurnal migrations and water temperature. Investigations by V. N. Nikitin created the basis for further intensive studies of species ecology and peculiarities in the zooplankton dynamics in the Black Sea. The earlier findings of Nikitin on the vertical distribution of plankton organisms, were later verified using large volume water samples of 100-150 liters by Vinogradov et al., (56).

The comparative analysis of the spatial distribution of the dominant zooplankton species in the Black Sea and other seas of the Mediterranean basin was carried out by Kovalev (58). In recent decades quantitative studies have also been carried out, to describe features of the horizontal and vertical distribution of zooplankton in regions with different pollution and eutrophication levels (30, 31, 56, 59-63).

Routine monitoring of the zooplankton composition in different regions of the sea began in the 1950s. The impacts of natural and anthropogenic factors on the fluctuation of the zooplankton communities were studied by various authors (59, 64-74).

By the 1940s physiological investigations of single species were already included in plankton studies. Respiration, reproduction and feeding habits were studied both in the field and laboratory (75-82). The knowledge gained from these investigations formed the basis for the analysis of the structure, productivity, condition (well-being) and functioning of the zooplankton communities and populations (e.g., 3, 64, 79, 80, 84-87 and several others).

Since the 1970s studies on zooplankton dynamics have gained a multidisciplinary approach by the modeling of complex interactions of the biota (e.g., 3, 88-94).

Qualitative and quantitative field observations are the prerequisites for the understanding of the dynamics of the zooplankton in particular and the ecosystem in general.

Sampling methods and treatment

In the first decades of investigations the Nansen and Genze Net were used for the collection of samples. Later Ukrainian, Russian, Romanian and Bulgarian investigators used the Juday Net. This net, having an opening of 0.1 m² is still in use in these countries as the basic device for zooplankton sampling. Long-term usage of the Juday net has been important in the comparison and analysis of results obtained in different countries. However the mesh sizes of the nets varied between 125 and 160 µm, which is insignificant for mesozooplankton sampling. Either a Hensen net of 300 µm mesh size or Nansen net of 112 µm mesh size have been used by Turkish zooplanktologists.

In the later zooplankton studies, size spectra were taken into account when sampling. Microzooplankton (<0.5 mm), mesozooplankton (0.5-10 mm) and macrozooplankton (> 10 mm) were studied by Zaika et al., (95). The material was sampled by either Juday or Bogorov-Rass nets. Meantime special trawls were developed for the collection of macrozooplankton (96). During recent studies large volume water samplers of 100-150 liters capacity were used for mesozooplankton sampling (97 & others). The advantage of the latter sampler is that these collectors can also be used at greater depths with a higher accuracy of depth recording and they allow the study of the vertical distribution of organisms in relatively thin layers. Furthermore with the development of neuston nets, specially designed for surface or near surface sampling, dense populations of zooplankton were discovered in the surface layers (55). The application of various sampling devices allowed the characterisation of species composition, abundance and biomass of all size groups of zooplankton as well as the description of trophic structures (98).

During the majority of field studies, standard depths (0, 10, 25, 50, 75, 100, 150, 200 m) or depth layers (10-0, 25-10, 50-25, 100-50, 150-100, 200-150) were sampled. However, in recent years the sampling was performed with respect to the physical and chemical characteristics of the water column, most notably the density (99). The simplets type of this sampling procedure considered the upper mixed layer, thermocline layer and from the lower boundary of the thermocline to the hydrogen sulfide layer. The boundary of the oxic/anoxic layer was identified according to the sigma theta depth of 16.2, indicating the start of the H₂S-zone (100). In recent international surveys the whole oxic zone was sampled by vertical hauls, to collect all migrating species during both day and night (101). The results were

expressed per m⁻².

For the quantification of organisms in the samples the same procedures were used by all countries. Small and abundant species were counted applying sub-sampling techniques, large animals were enumerated in the whole sample. Biomasses were determined using respective conversion factors for each stage/animal, obtained from the literature (102). Volumetric measurements were also utilised for certain zooplankton groups, [for example the ctenophores; (101)].

Main results

In the following, the overall results of the Black Sea zooplankton research concerning composition, spatial and seasonal distribution and temporal changes are presented.

Zooplankton composition

Earlier investigations carried out demonstrated that the planktonic fauna of the Black Sea is sparse compared with the Mediterranean Sea (14 & others). This fact was attributed mainly to salinity. Typical marine stenohaline organisms (Radiolaria, Siphonophora, Pteropoda, Salpae), inhabiting the Mediterranean Sea (salinity: S=36-38‰), are not able to survive in the Black Sea (salinity: S=17-20‰). Medusae, Ctenophora, Copepoda and other groups, that have a high species number in the Mediterranean Sea are also represented by only one or few species in the Black Sea (9). About 80 holoplanktonic species, mainly coming from the Mediterranean Sea, occur in the Black Sea. Koval (60) discovered about 150 species of planktonic animals in different regions of the northwestern shelf. Almost half of these species are brackish-water and fresh water species inhabiting estuarine regions and gulfs. Besides these, meroplankton which is made up of numerous larvae of benthic invertebrates were an important part of zooplankton (>30 species of Polychaeta, >50 species of Gastropoda, >20 species of Bivalvia; (22).

Zooplankton inhabiting the Black Sea can be divided into three ecogeographical groups. These are Mediterranean invaders, Pontic relicts and fresh-water species (9). Although the species number of the Mediterranean invaders in the Black Sea is not high, they are the main component of the zooplankton communities in this sea.

Despite the fact that the Black Sea zooplankton has been studied relatively well, taxonomic studies carried out

in the Black Sea can not be considered as complete. For example detailed morphologic investigations of copepoda species of the Black Sea led to changes in the taxonomic status of some species. The sole species of the genus *Centropages* occurring in the Black Sea, *Centropages ponticus* KARAV, is not a variation of the Mediterranean species *Centropages kröyeri* GIESBR. (16). The predominant species of *Calanus* of the Black Sea was previously identified as *Calanus helgolandicus* CLAUS. This was then reclassified as *Calanus helgolandicus ponticus* (17), and later as *Calanus ponticus* KARAV., (18). Recently this species was renamed as *Calanus euxinus* KRICH., (19). Regarding the taxonomic status of large and small forms of *Acartia clausi* GIESBR., it was supposed that they were sibling species (103). Furthermore, the Black Sea species of *Oithona* is *O. nana* (104) but not *Oithona minuta* as previously thought by some authors (102, 105 & others). It can be assumed that taxonomic investigations on some other zooplankton species may reveal similar results.

Mediterraneanization is an important process for the enrichment of the Black Sea zooplankton. Studies of the planktonic species transported by the lower Bosphorus flow through the Bosphorus Strait first began in the 1960s (106) and resulted in interesting findings: Bogdanova & Shmeleva, (107) have shown that few individuals of some Mediterranean species were distributed not only off the Bosphorus or in the central regions of the Black Sea but also as far north as Sevastopol Bay. Porumb (108) and Kovalev et al., (109) have recorded similar findings for the area around the Danube delta. The total number of Mediterranean species in the Black Sea was estimated to be around 50, the majority being concentrated in the Bosphorus region (110, 111). Some of the individuals sampled far from the Bosphorus area bear well developed gonads indicating possible adaptation and reproduction at the lower salinity conditions of this sea. With increasing salinity, they may become a permanent member of the Black Sea fauna as stressed by Puzanov (112). For example Gordina et al., (113) have observed recently eggs of the Sea of Marmara anchovy in the southern Black Sea coasts implying adaptation of this stock to the Black Sea. Another good example for this process, i.e. the enrichment of Black Sea flora and fauna by organisms originating in the Mediterranean Sea, is the copepod species *Acartia tonsa* which has been found in large quantities in recent years (111, 114). Apparently, this species adapts well to the Black Sea conditions. It is worth noting that *Mnemiopsis leidyi* (accidentally introduced into the Black Sea from the north Atlantic via ballast water of vessels) which formed

a huge biomass in the whole basin must also have adapted well to the Black Sea conditions (115-117).

However, particularly during the last two decades some species have either vanished or became rare in the pelagic ecosystem especially in the shelf waters (63, 118-120). The former mass occurring species, *Oithona nana* has not been found in samples collected by the IBSS or by other institutes since 1989. Now this species has again appeared in zooplankton samples, though very seldom. Many investigators (63, 70, 118, 120,) suppose that these changes are connected with anthropogenic impacts (i.e. pollution, eutrophication, predation pressure of *Mnemiopsis leidyi*) on the Black Sea ecosystem, because they are the most significant in coastal areas.

Distribution of zooplankton (abundance and biomass)

The large-scale horizontal and vertical distributions of zooplankton organisms have been the subject of several publications (37, 39, 44-46, 52, 60, 67, 121, 122). Earlier data (1960s and 1970s) were reviewed by Zenkevich (9), and Greze et al., (66), respectively. These authors discussed the natural factors determining the distribution patterns. It was shown that zooplankton abundance and biomass were quite variable in time and space. Nevertheless, the distribution patterns of zooplankton display some common features.

Temperature and wind stress were important for the distribution of the zooplankton (65): during warmer periods the abundance of zooplankton was generally high at the northwestern shelf zone (66). An important factor influencing the zooplankton abundance is certainly the riverine flow. A positive linear correlation between the volume of water discharged and the concentration of zooplankton was found by various authors (e.g., 42, 60, 62, 66).

High numbers of zooplankton occur usually in coastal and shelf areas, e.g. high patches occur southwestern Crimean coast and the Bulgarian, Romanian, Turkish and Caucasian coasts. Zooplankton numbers increase as well at convergence zones of the western and eastern gyres off the Crimean and Anatolian coast. Due to the dynamic characteristics of these regions a high productivity is ascertained (66).

The difference between the average biomass in central and coastal areas of the Black Sea is small, (excluding the northwestern shelf) in comparison with many other seas, including the neighbouring Mediterranean Sea (58). This is due to a fairly intensive vertical-exchange in central

areas of this sea (123) and horizontal water-exchange between central and coastal areas (58, 124, 125).

The spatial variability in the zooplankton biomass differs between central and coastal areas. In the same season the maximum value could be two to three fold that of the minimum biomass value found in the central Black Sea. However, in the coastal areas this variability is about ten fold (49, 58, 97, 121).

In the Black Sea the majority of zooplanktonic organisms are concentrated in the upper 50 meter layer. In the northwestern Black Sea zooplankton seem to be more evenly distributed vertically compared with in the centres of the main gyres. At the centers of the main gyres, the upper boundary of hydrogen sulfide is high and vertical water exchange is less intensive, hence causing considerable patchiness.

Investigations carried out using the large volume water sampler with accurate depth recording revealed that high concentrations of *Pleurobrachia rhodopsis*, *Calanus euxinus*, *Pseudocalanus elongatus*, *Sagitta setosa* occur at a depth of 60 m down to the lower boundary of the oxic-anoxic interface at around 120 m, where these species occur in a small band just above the anoxic layer during the day time (97, 126). Later it was shown that living and non-living organisms were accumulated at a certain layer of density gradient (between sigma theta 15.4 and 16.2) corresponding to low oxygen concentrations (82, 127).

Temporal changes

In the past and also in recent years the diurnal vertical migration of zooplankton has been studied by a number of authors (35, 39, 82, 128-131). As mentioned above, high concentrations of *Calanus euxinus*, *Pseudocalanus elongatus*, *Pleurobrachia rhodopsis* and *Sagitta setosa* have been observed at depths just above the H₂S-zone during the day time. During night the concentration of organisms at this depth decreases considerably due to upward migration (97). In an experiment, zooplankton were collected at 3 hourly intervals, at the center of the western (anti-cyclonic) gyre at two stations visited in April and August 1976. A two fold change in zooplankton numbers within 24 hours was observed. During the night many species migrate to surface layers and during the day they are concentrated in deep layers (132).

This phenomenon could be explained by diurnal feeding and reproduction rhythms of planktonic animals. Laboratory experiments on reproduction, hatching and moulting of three Black Sea copepod species have

confirmed the existence of a diurnal rhythm during these processes (133, 134). However contradictory observations were also made at different regions of the sea.

Numerous authors investigated seasonal changes in zooplankton abundance and biomass in different regions of the Black Sea (40, 48, 57, 60, 61, 67, 108, 121, 135-140). It was noted that seasonal differences in migration patterns occur in the central and coastal areas (58). Usually two peaks for maximum zooplankton abundance were marked for coastal areas coinciding with spring and autumn periods. Sometimes a third peak was observed in summer. In central areas however only one peak was observed, during summer. This is due to some differences in hydrological and hydro-chemical conditions between shallow coastal and deep-water central areas. These conditions, in particular seasonal changes of temperature and nutrient input are important for reproduction; zooplankton respond to such variation faster in coastal areas and bays than in the open sea. In the shallow Azov Sea the maximum temperature of the coastal water is reached a month before that of the open sea (141) and thus the spring bloom of zooplankton starts earlier (58).

Unlike the open sea, coastal waters display more clearly defined regeneration cycles of biogenic substances in the summer-autumn period. Because of intensive bacterial degradation of dead phyto- and zooplankton after the spring bloom and of the faster regeneration of nutrients the bacterio- and phytoplankton may again increase to peak concentrations in coastal waters. These processes provide an autumn peak and sometimes a summer peak of zooplankton abundance. In the open sea these processes are slow and generally start much later (in summer). The phytoplankton production is low and does not merit a second zooplankton peak in autumn (58).

Long-term changes in the zooplankton abundances of the Black Sea (45, 46, 64, 65) have been reviewed by Greze et al., (66). Recent investigations have revealed some changes in the long-term variability of zooplankton since the 1970s. Systematic observations during 1960-1981 in the open eastern sea area showed an increase in biomass of the main zooplankton species. During the 1960-1969 period the average biomass of the main zooplanktonic organisms in the upper 100 m was 308 mg/m³, this increased to 471 between 1970 and 1975 and again rose to 527 mg/m³ between 1976 and 1981. However, the increase was entirely due to two gelatinous species; the dinoflagellate *Noctiluca scintillans* and the

ctenophore *Pleurobrachia rhodopsis* (58, 67).

A similar increase in the zooplankton was observed in the western Black Sea between 1978 and 1980. In the upper 100 m the zooplankton biomass increased in offshore areas (water depth >200m) by 1.7-2 fold, at the northwestern shelf the increase was 3-4 fold, in shallow coastal areas (0-25m) this rose to 10 fold. These changes were caused exclusively by the increase in *Noctiluca scintillans* (59). This corresponds with a significant increase in the phytoplankton biomass in the northwestern Black Sea (142).

In the area off the Danube estuary (in the upper 100 m) the average zooplankton abundance and biomass increased by factors of 3 and 5 respectively from the 1970s to the 1990s (63). As for other areas the proportions of *Noctiluca* and *Acartia clausi* also increased significantly. Similar changes were noted by other authors (31, 62, 70). In the late 1970s a sharp increase in the medusa *Aurelia aurita* was discovered (143, 144 & others).

Then, during the 1980s drastic changes in the plankton communities occurred. Most markedly were severe changes in the phytoplankton and mesozooplankton composition and a sharp decrease in the total zooplankton biomass including *Aurelia aurita* at the end of the 1980s and beginning of the 1990s (32, 71, 111, 122, 145-147). Changes in the dominant species had already been observed at the north western shelf and in Sevastopol Bay over the last two decades (30, 63, 118-120). During 1960-1964 the share of *Acartia clausi* in the total copepod community was 17%. This increased to over 30 % during 1981-1985, and during 1991-1994 it rose to 75 %. Copepod species such as *Oithona nana* and *O. similis* and the cold water species *Pseudocalanus elongatus* and *Calanus euxinus*, which were dominant before the 1980s, decreased during the 1990s (32). The species *O. nana* was not observed from 1989-1996. In offshore areas the dominant species *Pseudocalanus elongatus*, *Calanus euxinus* and *Sagitta setosa* decreased 10 to 100 fold (145, 146).

Most changes began at the beginning of the 1980s, when the northwestern Atlantic ctenophore *Mnemiopsis leidyi*, originating from eutrophic lagoons in North America, was accidentally introduced into the Black Sea (105, 116, 148). *M. leidyi* reproduced intensively and bloomed during 1988-1990 reaching a remarkable total biomass of 800 million tons wet weight in the Black Sea during the summer of 1989 (116). *Mnemiopsis* displayed the typical pattern of a new coloniser. After its mass development in the years 1989 and 1990 with 2 kg m⁻²

in the open sea and average numbers of $4.7 \text{ kg} \cdot \text{m}^{-2}$ at the northwestern shelf (145), its number and biomass had dropped to moderate levels by 1991 and afterwards fluctuated until 1996 in a range of $0.2 - 0.5 \text{ kg} \cdot \text{m}^{-2}$ (146, 149, 150).

After the peak of *Mnemiopsis* the biomass of *Aurelia* increased again and since the summer of 1991 until 1994 the biomass of both species remained more or less at the same level (149). After 1993 the mesozooplankton biomass recovered and displayed an increasing trend (71, 111, 122, 145, 146, 147, 151). And during 1997 the species *Oithona nana* was present again in some zooplankton samples.

Numerous investigators consider anthropogenic impacts as the main reason for the changes in the zooplankton of the Black Sea in recent years especially stressing eutrophication in estuaries (60, 63, 70, 105, 118, 120, 152). Additionally, the negative effect of pollution (e.g. oil pollution) could be the reason for the observed decrease in the zooplankton in the upper water layers (59, 61). The early stages in the life cycle of zooplankton and fishes in particular, which are very sensitive to pollution, live in the near surface layer.

Another important factor for the decrease in the zooplankton after 1988 was assumed to be the mass development of *Mnemiopsis leidyi* being both a voracious predator and a strong food competitor (105, 111, 117, 122, 127, 145, 147, 153, 154). The arguments are: zooplankton which inhabit the same depth range (0-30m) as *Mnemiopsis* seem to have suffered most. For example in the northwestern Black Sea the total biomass of fodder zooplankton (discounting *Noctiluca*, *Pleurobrachia*, *Aureila* and *Mnemiopsis*) was 50 times less in 1990 than in the 1975-1980 period. Furthermore, the biomass of *Aurelia aurita* competing with *Mnemiopsis* at the same trophic level decreased 100 fold (120, 145). At the same time the change in the biomass of fodder zooplankton living mainly below the thermocline, where *Mnemiopsis* was only occasionally found, was significantly less compared with those found above the thermocline (151). In the last few years, since the *M. leidyi* biomass has remained at moderate levels, the total biomass (wet weight) of *Aurelia* has shown an increasing tendency along with other zooplankton (58, 145, 151).

The impact of *M. leidyi* on the Black Sea zooplankton fits the general view, that possible reasons for the fluctuation of the zooplankton biomass and the changes observed in the species composition could be due to long term changes in prey/predator relationships (3, 67). According to Bogdanova and Vodyanitskiy (155), and

Zhorov and Boguslavskiy (156) long-term fluctuations in zooplankton could also be explained by interactions between biomass cycles and long-term climatic cycles (river flows and other hydrological parameters).

It is known that global atmospheric changes, such as the Southern Oscillation (SO), El Niño-Southern Oscillation (ENSO), and North Atlantic Oscillation (NAO) influence pelagic communities (157-161). Although the short and long term atmospheric variabilities in the Eastern Mediterranean/Black Sea regions are well known, hypotheses on tele-connections with global atmospheric events have recently been put forward (162). A good correlation between the Black Sea hydrology, the North Atlantic Sea Surface Temperature (SST) variability and the ENSO-type variability of the tropical Pacific Ocean, which indicates a global ocean-atmosphere coupling. Both types of variability lead to changes in the cyclone trajectories, precipitation over the river drainage basins, changes in river discharges, resulting in alterations in the northwestern Black Sea hydrography as shown by Polonsky et al., (163), which may finally cause changes in the Black Sea plankton community (74).

Comparison of the weather regime of the 1980s with previous decades showed indeed that the climate of the 1980s was unique. Especially the 1983-1990 period was dominated by a positive phase of the NAO, which increased strongly after 1988 (164). The oscillation of the global air temperature, which has a cyclic occurrence of 65-70 years gave its lowest amplitude during the mid eighties (165) which had again risen by the end of the 1980s. It may be interesting to note that the sun activity, which has a cycle of 11-12 years, was very low during the mid eighties and again increased to high values during 1988 (164). The 1980s displayed the largest annual variability on a global scale, of the 20th century (166). The Southern Oscillation experienced the strongest warm (El Niño) episode of the century (1982/83) and the strongest cold (La Niña) episode in 50 years (1988). According to Trenberth & Hurrell (167), the very strong La Niña event of 1988 apparently terminated the climate regime that was established in 1977.

Niermann et al., (74) compared the fluctuation of the plankton in different seas of the northern hemisphere and concluded hypothetically, that changes in the weather regime during the 1980s could have triggered the changes in the phyto- and mesozooplankton communities of the Black Sea, which caused the conditions for the outburst of *M. leidyi* and the subsequent decline in the anchovy stock during 1988-1990 (168).

Summarized are the significant changes in the

zooplankton community of Black Sea which took place during recent decades. The man-made eutrophication and pollution, especially at the shelf regions and in the estuaries of large rivers, seem to have had a strong impact on the zooplankton fluctuation. But less attention should not be given to natural factors which represent the first driving forces of the zooplankton community and which are modified or intensified by anthropogenic impact. While analysing the long-term changes in the zooplankton during the NATO TU-Black Sea program a connection between climatic changes (water temperature in particular) and zooplankton biomass (71, 111) could be established. Similar developments were found for other regions, including areas with less important anthropogenic influence (74, 160, 169). Overall it could be supposed that climatic changes are the fundamental causes determining the processes leading to changes in zooplankton composition (in qualitative and quantitative terms), but anthropogenic factors may considerably influence and change the shape of these processes.

Future investigations needed

The experience gained during the taxonomic and faunistic studies carried out in recent years proves the necessity of their continuation. Revision of some groups such as Protozoa is necessary. A basic task is to standardize the scientific names of the species. This will minimize the confusion which occurs when dealing with different names thus enabling compilation of data from numerous investigations into an international database.

References

1. Vinogradov, K. A., Articles on history of domestic hydrobiological investigations of the Black Sea. Kiev. Pub. of Ac. Sci. UkrSSR, 155 p. 1958, (in Russian).
2. Vodyanitskiy, V. A., Burdak, V. D., Kiseleva, M. I., Kovalev, A. V., Zoological investigations of A. O. Kovalevskiy Institute of Biology of Southern Seas. Ac. Sci. UkrSSR (100. Anniversary of Sevastopol Biological Station). Zool. Zhurn., V. 50, N9: 1429-1433, 1971, (in Russian).
3. Greze, V. N., Ecosystem of the Black Sea: Results and prospects for investigations. Visn. Acad. Nauk. USSR. N: 4: 24-32, 1985, (in Ukrainian).
4. Greze, V. N., Kovalev, A. V., Main stages of centenary investigations of zooplankton at the IBSS. Zool Record, N. S: 12-17, 1971, (in Russian).
5. Ünlüata, Ü., Aubry, D. G., Belberov, Z., Bologa, A., Emereev, V., & Vinogradov, V., Cooperative marine science program for the Black Sea. EOS, 74: 401-412, 1993.
6. Ostroumov, A. A., Preliminary report about participation in the Black Sea deep-water researches in 1891. Notes of Novrosiysk Naturalists Soc., V.16, N2: 135-148, 1891. (in Russian).
7. Karavaev, V. A., Materials at fauna of pelagic crustacean of the Black Sea. Notes of Kiev Naturalists Soc., XIII, 1-2: 35-62, 1894, (in Russian).
8. Karavaev, V. A., Materials at fauna of pelagic crustacean of the Black Sea. Notes of Kiev Naturalists Soc., V. XIX, 1: 117-174, 1895, (in Russian).
9. Zenkevich, L. A., Biology of the seas in USSR. M.: Pub. Ac. Sci. USSR., 739 p., 1963, (in Russian).

Monitoring of species composition (community structures) will enable investigators to estimate changing trends within the ecosystem. Trends in species successions, and the recording of acclimatisation or disappearance of immigrated or accidentally introduced species could be recorded only with systematic studies. Such investigations relate well to the actual problem of saving the biodiversity. Qualitative and quantitative monitoring of the living and non-living system is necessary in order to determine the healthiness of the Black Sea ecosystem and to detect changes due to natural or anthropogenic impacts, both on a short and long-term time scale.

It is important to continue studying the ctenophore *M. leidyi*, including its dynamics and interrelations with other zooplankton species. It should be monitored regularly to clearly understand if its biomass is now at a stable level.

Steps already made regarding the unification of sampling methods and continued cooperation amongst the riparian countries of the Black Sea should be maintained. This is the prerequisite for the successful research and management of this basin.

Acknowledgements

The present investigation was carried out with the support of the NATO Linkage Grant (ENVIR.LG 951569; NATO-TU.BLACK SEA PROGRAMME), under the Science for Stability Programme of Scientific Affairs Division of NATO, and was also funded by the Scientific and Technical Research Council of Turkey (TUBİTAK). We thank Mrs Alison M. Kideyş for improving the English of the text.

10. Petipa, T. S., Development and tasks of biological investigations in pelagial of the Black Sea. Changeability of the Black Sea ecosystem. Natural and anthropogenic factors. (ed. Vinogradov, M. E.) Moscow, Nauka, 336-345. 1991.
11. Mechnikov, I., Notes about pelagic fauna of the Black Sea. Proc. of 1st Congress of Russian naturalists. Zoological Dep., S. Petersburg: 268-270. 1868, (in Russian).
12. Ul'yanin, V. N., About pelagic fauna of the Black Sea. News of Soc., of Nat. Sciences, Anthropology and Ethnography lovers in Moscow. v. 8, N1: 58-62. 1870, (in Russian).
13. Pereyaslavtseva, S. M., Protoza of the Black Sea. Notes of Novorossiysk naturalists society, V.10, N2: 79-144. 1886, (in Russian).
14. Ostroumov, A. A., Reports about dredging and plankton catches of "Selyanika" expedition. News of Ac. Sci., V.5, N1: 33-92. 1896, (in Russian).
15. Sovinskiy, V. K., Introduction to studying of Pontic-Caspian-Aral basin fauna. Notes of Kiev Naturalists Soc., v.18: 497 p. 1904, (in Russian).
16. Kovalev, A. V. About systematical state and distribution of *Centropages kroyeri* var. *pontica* Karavajev (Crustacea, Copepoda). Biology and distribution of plankton of the southern seas. M: Nauka.: 94-98. 1967, (in Russian).
17. Kovalev, A. V., Shelukhin, E. P., Ivanov, V. N. About taxonomic state of the Black Sea *Calanus* (Copepoda). Zool. Zhurn., V. 54, N2: 195-199. 1975, (in Russian).
18. Fleminger, A., Hulsemann, K., Geographical variation in *Calanus helgolandicus* I. (Copepoda, Calanoida) and evidence of recent speciation of the Black Sea population. Biol. Oceanography. N5: 43-81. 1987.
19. Hulsemann, K., *Calanus euxinus*, new name, a replacement name for *Calanus ponticus* Karavajev, (Copepoda: Calanoida). Proc. Biol. Soc. Wash., 104 (3): 620-621. 1991.
20. Kiseleva, M. I., Pelagic larvae of polychaeta of the Black Sea. Proceedings of Sev. Biol. St. V.9: 59-111. 1957, (in Russian).
21. Petran, A., Sur la faune meroplantonique du littoral Roumain de la mer Noire (donnees quantitatives pour les annees 1972-1979). CIESM. Journées Etud. System. Biogeogr. Médit., Cagliari: 89-90. 1980.
22. Murina, V. V., Meroplankton. Plankton of the Black Sea (eds. Kovlev A. V., Finenko Z. Z.). Kiev, Naukova dumka: 194-204. 1993, (in Russian).
23. Sazhina, L. I., Development of the Black Sea Copepoda. I. Nauplii stages of *Acartia clausi* Giesbr., *Centropages kroyeri* Giesbr., *Oithona minuta* Kritcz. Proc. of Sevast. Biol. Stat., 13: 49-67. 1960, (in Russian).
24. Sazhina, L. I., Nauplii of mass species of pelagic copepods of the World Ocean. (ed. Petipa, T. S.). Kiev, Naukova dumka: 238 p. 1985, (in Russian).
25. Braiko, V. D., Bachesku, M., Vinogradov, K. A., Identification book for the Black Sea and the Azov Sea fauna. V. 1. Free living invertebrates. Protozoa, Porifera, Coelenterata, Nematelminthes, Annelida, Tentaculata (ed. Vodyanitskiy V. A.). Kiev, Naukova dumka: 437 p., 1968, (in Russian).
26. Bachesku, M., Vasilenko, S. V., Greze, I. I., Identification book for the Black Sea and the Azov Sea fauna. V.2. Free living invertebrates. Crustacea (ed. Vodyanitskiy V. A.). Kiev, Naukova dumka: 536 p., 1969, (in Russian).
27. Baranova, Z. I., Bachesku, M., Golikov, A. N., Identification book for the Black Sea and the Azov Sea fauna. V. III. Free living invertebrates. Arthropoda (except Crustacea), Mollusca, Echinodermata, Chaetognatha, Chordata (ed. Vodyanitskiy V. A.). Kiev, Naukova dumka: 337p., 1971, (in Russian).
28. Mutlu, E., Ünsal, M., Bingel, F., A preliminary view on the faunal assemblage of soft-bottom crustaceans along the near-shores of the Turkish Black Sea. Acta Adriat. 33 (1/2): 177-189. 1992.
29. Mutlu, E., Ünsal, M., Bingel, F., Faunal community of soft-bottom mollusks of the Turkish Black Sea coast. Tr. J. of Zoology 17: 189-206. 1993.
30. Zaitsev, Yu. P., Impact of eutrophication on the Black Sea fauna. GFCM., Stud. & Rev. Rome, FAO, 64: 63-86. 1993.
31. Konsulov, A., Qualitative composition of zooplankton in the Black Sea in front of the Bulgarian coast. A. I. D. National Biological Resources Conservation Stragetegy: 11-15. 1993.
32. Kovalev, A. V., Gubanova, A. D., Kideyş, A. E., Melnikov, V. V., Niermann, U., Ostrovskaya, N. A., Scryabin, V. A., Uysal, Z., Zagorodnyaya, Yu. A., Long-term changes in the biomass and composition of fodder zooplankton in coastal regions of the Black Sea during the period 1957-1996. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Reuslts, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 1: pp. 209-220, 1998.
33. Zernov, S. A., To the question on annual changes of the Black Sea zooplankton at Sevastopol. News of Imperial Academy of Sciences, V.20. N4: 119-134. 1904, (in Russian).
34. Zernov, S. A., To the question on studying of the Black Sea life. Notes of Imperial Academy of Sciences. Physics-mathematics department. Series 7, V.32. S.-Petersburg: 299 p. 1913, (in Russian).
35. Nikitin, A. N., Vertical distribution of plankton in the Black Sea. Proc. Sp. zool. lab. and Sevast. Biol. St. AcSci, Ser.2. N9: 93-140. 1926, (in Russian).

36. Nikitin, V. N., Vertical distribution of plankton in the Black Sea. Zooplankton except Copepoda and Cladocera. Proc. of Sevast. Biol. St., 1: 27-117. 1929, (in Russian).
37. Nikitin, V. N., Distribution of biomass of the Black Sea plankton. Rep. Ac. Sci. USSR, 47.N7: 529-532. 1945, (in Russian).
38. Galadzhiev, M. A., Materials on the Black Sea Rotatoria fauna. Proceedings of Sevastopol Biological Station. v.6: 224-235. 1948, (in Russian).
39. Kusmorskaya, A. P., About zooplankton of the Black Sea. Proceedings of AzCHERNIRO, 14: 177-214. 1950, (in Russian).
40. Kusmorskaya, A. P., Seasonal and annual changes of the Black Sea zooplankton. Proceedings of VGBO of Acad. Nauk. USSR, 6: 158-192. 1955, (in Russian).
41. Demir, M., Pontellidae and Parapontellidae (Pelagic Copepoda) from the southern Black, Marmara and north-east Aegean seas. Hydrobiol. Istanbul. B.4, N4: 176-179. 1959.
42. Braiko, V. D., Goromosova, S. A., Pitsyk, G. K., Fedorina, A. I., Dynamics of the Black Sea zooplankton on observations of 1956-1958. Proceedings of AzCHERNIRO, 18: 29-49. 1960, (in Russian).
43. Petipa, T. S., Sazhina, L. I., Delalo, E. P., Vertical distribution of zooplankton of the Black Sea connected with hydrological conditions. Rep. Ac. Sci. USSR, 133, N4: 964-967. 1960, (in Russian).
44. Petipa, T. S., Sazhina, L. I., Delalo, E. P., Distribution of zooplankton in the Black Sea in 1951-1956. Oceanology, 3, N4: 110-129. 1963, (in Russian).
45. Dimov, I. Zooplankton of the Bulgarian Black Sea coast in 1954, 1955 and 1956. Pr. Inst. of Fisheries. N2: 85-147. 1960, (in Bulgarian).
46. Dimov, I. Zooplankton of the western coast of the Black Sea in 1960-1964. -Pr. Inst. of Fisheries and Oceanography. Varna, N7: 45-66. 1966, (in Bulgarian).
47. Goromosova, S. A., Seasonal development and distribution of zooplankton in the north-western part of the Black Sea. Proceedings of AzCHERNIRO, 23: 69-80. 1964, (in Russian).
48. Konsulov, A., Seasonal and annual dynamics of zooplankton in the Black Sea coast in the 1967-1970 period. Institute of Oceanography and Fisheries, Varna, 12: 64-78. 1974.
49. Konsulov, A., On the vertical distribution of zooplankton of the Bulgarian Black Sea coast. Institute of Fisheries, Varna, 15: 67-82. 1977.
50. Mamaeva, N. V., Microzooplankton of the open Black Sea in: Ecosystems of pelagial of the Black Sea. (ed. Vinogradov, M. E.). M.: Nauka: 168-173. 1980, (in Russian).
51. Moiseev, E. V., To the study of zooflagellates of the Black Sea. In: Seasonal changes of the Black Sea zooplankton (ed. Vinogradov, M. E.). Moscow, Nauka: 100-112. 1983, (in Russian).
52. Zagorodnyaya, Yu. A., Unevenness of space distribution of meso- and macroplankton in the summer period in 1980. Waters dynamics and plankton productivity in the Black Sea (eds. Zats V. I., Finenko, Z. Z.). Moscow. Coord. Centre of SEV on problem "The World Ocean": 340-356. 1988, (in Russian).
53. Kopylov, A. J., Ciliates of the oxic zone of the Black Sea. In Structure and production parameters of planktonic communities of the Black Sea (ed. Vinogradov, M. E.). M.: Nauka: 156-172. 1989, (in Russian).
54. Dolgopolskaya, M. A., Zooplankton of the Black Sea in the Karadag region. Proceedings of Karadag Biological Station, 6: 57-111. 1940, (Russian).
55. Zaitsev, Yu. P., Marine neustonology. Kiev: Naukova dumka, 264 p. 1970, (in Russian).
56. Vinogradov, M. E., Flint, M. V., Shushkina, E. A., Vertical distribution of mesoplankton in the open area of the Black Sea. Mar. Biology. Vol. 89: 95-107. 1985.
57. Petran, A., Evolution saisoniere du zooplancton dans les eaux du large de la mer Noire. Rapp. Comm. Int. mer Medit., Monaco, 30 (2): 203-204. 1986.
58. Kovalev, A. V., Structure of zooplankton communities of the Atlantic and the Mediterranean basin. Kiev: Naukova dumka, 141 p. 1991, (in Russian).
59. Polischuk, L. N., Nastenko, E. V., Garkavaya, G. P. Some peculiarities of modern state of pelagic and neustonic zoocenosis of the Black Sea. Ecologiya Morya. v.18: 25-34. 1984, (in Russian).
60. Koval, L. G., Zoo- and necro-zooplankton of the Black Sea. Kiev: Naukova dumka.: 127 p. 1984, (in Russian).
61. Kovalev, A. V., Seasonal changes of zooplankton in the Sevastopol Bay. Hidrobiol. Zhurn. 16, N6: 9-14. 1980, (in Russian).
62. Petran, A., Evolution des biomasses du zooplancton de la mer Noire dans le secteur situe devant les embouchures du Danube, pendant les annees 1977-1980. Rapp. Comm. Int. mer Medit., Monaco, 29 (9): 319-320. 1985.
63. Porumb, F. Evolution du zooplancton des eaux du plateau continental Roumain de la mer Noire au cours de trois decennies. Rapp. et proc. -verb. reun. Commis. Int. Explor. Sci. Mer. Medit., Monaco. v. 33: 266. 1992.
64. Greze, V. N., Baldina, E. P., Bileva, O. K., Dynamics of numbers and production of main zooplankton components in the neritic zone of the Black Sea. Biologiya morya. Kiev. v.24: 12-49. 1971, (in Russian).

65. Fedorina, A. I. Dynamics of the Black Sea zooplankton development and reasons causing it. VNIRO. M.: 49 p. Dep. in CNIITACRH 11.05.78. N149. 1978, (in Russian).
66. Greze, V. N., Boguslavskiy, S. G., Belyakov, Yu. M., Principles of biological productivity of the Black Sea (eds. Greze V. N.). Kiev: Naukova dumka: 391, 1979, (in Russian).
67. Mashtakova, G. P., Long-term dynamics of planktonic community of eastern part of the Black Sea. Oceanological and fishery researches of the Black Sea. M.: Nauka, 50-61. 1985, (in Russian).
68. Bingel, F., Occurrence of jelly fish in the Black Sea-Marmara junctions of the Bosphorus. 2nd workshop on Jellyfish in the Mediterranean Sea, Trieste, Italy, 2-5 September, 1987, UNEP MAP Tech. Rep., 47: 58-70. 1991.
69. Simonov, A. U., Ryabinin, A. J., Gershanovich, D. E (eds) Project "Seas of the USSR". Hydrometeorology and Hydrochemistry of the Seas in the USSR. V. IV. Black Sea II. Hydrochemical condition and oceanographical basis of biological productivity. St. Petersburg, Hydrometeoizdat: 219 p. 1992, (in Russian).
70. Bologa, A. S., Bodeanu, N., Petran, A., Tiganus, V., Major modifications of the Black Sea benthic and planktonic communities in the last decades. Rapp. et proc. verb. des reunions, 33: 15. 1992.
71. Kovalev, A., Niermann, U., Melnikov, V., Belokopitov, V., Uysal, Z., Kideyş, A. E., Unsäl, M., Altukhov, D., Long-term changes in the Black Sea zooplankton: the role of natural and anthropogenic factors. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 1: pp. 221-234, 1998.
72. Konsulov, A., Kamburska, L., Sensitivity to anthropogenic factors of the plankton fauna adjacent to the Bulgarian coast of the Black Sea. In: Özsoy and Mikaelyan (eds.), Sensitivity to Change: Black Sea, Baltic Sea and North Sea. Kluwer Academic Publisher, Dordrecht, 95-104. 1997.
73. Niermann, U., Bingel, F., Gorban, A., Gordina, A. D., Gücü, A. C., Kideyş, A. E., Konsulov, A., Radu, G., Subbotin, A. A., Zaika, V. E., Distribution of anchovy eggs and larvae (*Engraulis encrasicolus* Cuv.) in the Black Sea in 1991 and 1992 in comparison to former surveys. ICES J. of Mar. Sci., 51: 395-406. (1994).
74. Niermann, U., Kideyş, A. E., Kovalev, A. V., Melnikov, V., Belokopytov, V., Long-term fluctuation of the zooplankton of the open Black Sea in comparison to other regions of the world. In: NATO-ARW: Environmental Degradation of the Black Sea: Challenges and Remedies. Ü. Ünlüata & A. Bologa (eds.), Kluwer Sci. Publ. in press.
75. Mironov, G. N., About feeding of some planktonic organisms of the Black Sea. Proc. of Zool. Inst. of Ac. Sci. USSR, v.7, N2: 217-220. 1941, (in Russian).
76. Delalo, E. P., Preliminary data on feeding of *Paracalanus parvus* (Claus) in the Black Sea. Pr. of Sevast. Biol. St., v. 14: 127-135. 1961, (in Russian).
77. Petipa, T. S., Pavlova, E. V., Mironov, G. N., Power balance of mass organisms from different ecological systems of the Black Sea. In: Book of II International Oceanographic Congress. Moscow: 307-398. 1966, (in Russian).
78. Petipa, T. S., Trophic dynamics of copepods in marine planktonic communities. Kiev, Naukova dumka: 242 p. 1981.
79. Shulman, G. E., Yuneva, T. A., Yunev, O. A., Svetlichnyi, L. S., Anninsky, B. E., Romanova Z. A., Arashkevich, E. G., Kideyş, A. E., Bingel, F., Uysal Z., Yılmaz, A. A biochemical approach for the estimation of food provision for heterotrophic organisms of the Black Sea. In: NATO TU Black Sea project: Sympos. on Sci. Results. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 1: pp. 263-280, 1998.
80. Anninskiy, B. E., Romanova, Z. A., Abolmasova, G. I., Gücü, A. C., Kideyş, A. E., The ecological and physiological state of the ctenophore *Mnemiopsis leidyi* (Agassiz) in the Black Sea in autumn 1996. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 1: pp. 249-262, 1998.
81. Beşiktepe, Ş., Kideyş, A. E., Unsäl, M., In situ grazing pressure and diel vertical migration of female *Calanus euxinus* in the Black Sea. Hydrobiologia 1988 (in press).
82. Beşiktepe, Ş., Studies on some ecological aspects of copepods and chaetognaths in the southern Black Sea, with particular reference to *Calanus euxinus*. Ph.D. Thesis, IMS-Middle East Technical University/Ankara, Turkey, 201 pp. 1998.
83. Greze, V. N., Kovalev, A. V., Main stages of centenary investigations of zooplankton at the IBSS. Zool Record. N. S: 12-17. 1971, (in Russian).
84. Petran, A., Elemente privind productia zooplanctonului trophic in anul 1976. Cercetari Marine, IRCM, Constanta, 9: 105-117. 1976, (in Romanian).
85. Zaika, V. E., Comparative productivity of hydrobionts (ed. Greze, V. N.). Kiev, Naukova dumka. 206 p. 1983, (in Russian).
86. Zagorodnyaya, Yu. A., Change of the size structure and productive indices of zooplankton. In: The modern state of the Black Sea ichthyofauna (ed. Konovalov, S. M.). Sevastopol: 145-152. 1995, (in Russian).

87. Ostrovskaya, N. A., Gubanova, A. D., Kideys, A. E., Melnikov, V. V., Niermann, U., Ostrovsky, E. V., Production and biomass of *Acartia clausi* in the Black Sea during summer before and after the *Mnemiopsis* outburst. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 1: pp. 163-170, 1998.
88. Petipa, T. S., Pavlova, E. V., Mironov, G. N., Structure of fodder nets, transferring and using of substance and energy in planktonic communities of the Black Sea. *Biologiya morya*, 19: 3-43. 1970, (in Russian).
89. Belyaev, V. I., A model of the shelf ecosystem for estimation of its potential bio-productivity. System analysis and processes modelling on the Black Sea shelf (ed. Nelepo B. A.). Sevastopol MHI AN USSR: 7-18, 1983, (in Russian).
90. Vinogradov, M. E., Shushkina, E. A., Lebedeva, L. P., Nezhlin, N. P., Structural and functional analysis of planktonic communities of the Black Sea. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 2: pp. 237-256, 1998.
91. Shushkina, E. A., Vinogradov, M. E., Lebedeva, L. P., Oğuz, T., Dyakonov, V. Yu., Nezhlin, N. P. and Anokhina, L. L., Studies of structural parameters of planktonic communities of the open part of the Black Sea relevant to ecosystem modeling. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 1: pp. 311-326, 1998.
92. Oğuz, T., Ducklow, H., Shushkina, E. A., Malanotte-Rizzoli, P., Tuğrul, S., Lebedeva, L. P., Simulation of upper layer biochemical structure in the Black Sea. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 2: pp. 257-300, 1998.
93. Oğuz, T., Ducklow, H. W., Malanotte-Rizzoli, P., Murray, J. W., Vedernikov, V. I., Ünlüata, Ü., A physical-biochemical model of plankton productivity and nitrogen cycling in the Black Sea. Accepted., *Deep Sea Research*, 1997. 35p+17 Figs.
94. Gücü, A. C., Oğuz, T., Modeling trophic interrelationships in the Black Sea. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 2: pp. 359-372, 1998.
95. Zaika, V. E., Moryakova, V. K., Ostrovskaya, N. A., Tsalkina, A. V., Distribution of marine micro-zooplankton. Kiev.: Naukova dumka. 92 p. 1976, (in Russian).
96. Melnikov, V. V., Device for collection of zooplankton samples. A.1486129 SSR MKI A.01 K 61/00- N 425756/31-13/Dec. 08.04.87; pub. 15.06.89. Bull. N22/Discoveries and inventions. 22: 17. 1989, (in Russian).
97. Vinogradov, M. E., Shushkina, E. A., Peculiarities of vertical distribution of the Black Sea zooplankton. *Ecosystems of the Black Sea pelagial*. M.: Nauka, 179-191. 1980, (in Russian).
98. Kovalev, A. V., Bileva, O. K., Moryakova, V. K., Combined method of collecting and registration of marine zooplankton. *Biologiya morya* (Vladivostok), N4: 78-82. 1977, (in Russian).
99. Saydam, C., Tuğrul, S., Baştürk, Ö., Oğuz, T., Identification of the oxic/anoxic interface by isopycnal surface in the Black Sea. *Deep Sea Res.* 40: 1405-1412. 1993.
100. Tuğrul, S., Baştürk, Ö., Saydam, C., Yılmaz, A., Changes in the hydrochemistry of the Black Sea inferred from water density profiles. *Nature*, 359: 137-139. 1992.
101. Niermann, U., Kideys, A., (eds), Assessment of the Phyto- and Zooplankton investigations during the CoMSBlack 1993 and the joint TU-Black Sea-CoMSBlack cruises 1994 and future plans. Report on the meeting of biologists in Erdemli, Turkiye (20.Feb.-3.March). Institute of Marine Science, Middle East Technical University, 33731 Erdemli, İçel, Turkey: 100 pp. 1995.
102. Petipa, T. S., About average weight of main forms of zooplankton of the Black Sea *Proc. Sevast. Biol. St.*, 9: 39-57. 1959.
103. Kovalev, A. V., Structure of zooplankton communities and its changes in the open Atlantic and in the Mediterranean basin. Inag. Diss. of Biol. Sciences Sevastopol., 454 p. 1988, (in Russian).
104. Sazhina, L. I., Kovalev, A. V., About synonyms of crustacean Copepoda of the Black Sea. *Zool. zhurn.*, V.50: 1099-1101. 1971, (in Russian).
105. Zaitsev, Yu. P., Recent changes in the trophic structure of the Black Sea. *Fisheries Oceanography*, 1: 180-189. 1992.
106. Pavlova, E. V., About discovering of the Mediterranean organisms in the Black Sea plankton. *Zool. zhurn.* V. 43, N11: 1710-1713. 1964, (in Russian).
107. Bogdanova, A. K., Shmeleva, A. A., Hydrological conditions of penetration of the Mediterranean planktonic species to the Black Sea. Waters dynamics and questions on hydrochemistry of the Black Sea. Kiev: Naukova dumka, 156-166. 1967, (in Russian).
108. Porumb, F., Presence de quelques especes Meditteraneeenes dans le zooplancton de la mer Noire. *Rev. roum. biol. ser. biol. animal.* Acad RSR. 25.2: 167-170. 1980.
109. Kovalev, A. V., Shmeleva, A. A., Petran, A., Zooplankton of the north-western part from the Bosphorus to the Danube in May, 1982. Waters dynamics and plankton productivity in the Black Sea. M.: Coord. Centre of SEV on problem "The World Ocean", 356-367. 1988 (in Russian).
110. Kovalev, A. V., Shmeleva, A. A., Penetration of Copepoda into the Black Sea through the Bosphorus - an indication of further Mediterraneanization of the Black Sea fauna. Second International Conference on Copepoda: Programme and Abstr. Ottawa: 52. 1984.

111. Kovalev, A., Beşiktepe, Ş., Zagorodnyaya, Yu. A., Kideyş, A. E., The Mediterranization of the Black Sea zooplankton is continuing. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 1: pp. 199-208, 1998.
112. Puzanov, I. I., Mediterranization of the Black Sea fauna and prospects of its strengthening. Zool. zhurn. 46, v.9: 1287. 1967. (in Russian).
113. Gordina, A. D., Nikolskiy, V. N., Niermann, U., Bingel, F., Subbotin, A. A., New data on the morphological differences of anchovy eggs (*Engraulis encrasicolus* L) in the Black Sea. Fisheries Res. 31: 139-145. 1997.
114. Belmonte, I., Mazzocchi, M. I., Prusova, I. Yu., Shadrin, N. V., *Acartia tonsa* a species new for the Black Sea fauna. Hydrobiologia 292/293: 9-15. 1994.
115. Zaitsev, Yu. P., Vorob'eva, L. V., Alexandrov, B. G., New Ctenophora species in the Black Sea. Dep. in VINITI. N5846. V.88-M. VINITI. 1-6. 1988. (in Russian).
116. Vinogradov, M. Ye., Shushkina, E. A., Musayeva, E. I., Sorokin, P. Yu., A newly acclimated species in the Black Sea: The ctenophore *Mnemiopsis* (Ctenophora: Lobata). Oceanology 29(2): 220-224. 1989.
117. Kovalev, A. V., Zaika, V. E., Sergeeva, N. G., Distribution and ecologo-physiological parameters of ctenophora - An invader in the Black Sea and the changes of plankton caused. VI meeting of All-Union Hydrobiol. Soc. Murmansk: Polyarnaya pravda, P.1: 137-138. 1991. (in Russian).
118. Vinogradov, M. E., Shushkina, E. A., Mass development of ctenophore *Mnemiopsis* as display of anthropogenic impact on the marine ecosystem. Practical ecology of the marine regions. The Black Sea (ed. Keondjan V. E., Kudin A. M., Terekhin, Yu. V.), Kiev: Naukova dumka, 94-106. 1990. (in Russian).
119. Zaitsev, Yu. P., Garkavaya, G. P., Nesterova D. A., Modern state of north-western Black Sea ecosystem. Modern state of the Black Sea ecosystem. M.: Nauka, 216-228. 1987. (in Russian).
120. Kovalev, A. V., Melnikov, V. V., Ostrovskaya, N. A., Prusova, I. Yu. Macroplankton. Plankton of the Black Sea (eds. Kovalev, A. V., Finenko, Z. Z.). Kiev: Naukova dumka, 183-193. 1993. (in Russian).
121. Nikitin, V. N., Basic regularities of life distribution in the Black Sea. Proc. of Inst. Oceanology Ac. Sci. USSR. 3: 173-190. 1949. (in Russian).
122. Petran, A., Moldoveanu, M., Post-invasion ecological impact of the Atlantic ctenophore *Mnemiopsis leidyi* Agassiz, 1865 on the zooplankton from the Romanian Black Sea waters. Cercetari Marine, 27-28: 135-157. 1997.
123. Vodryanitskiy, V. A., About problem of biological productivity of reservoirs and of the Black Sea in particular. Proc. of Sevast. Biol. St. of Ac. Sci. USSR. 8: 347-433. 1954. (in Russian).
124. Oğuz, T., Latun, V. S., Latif, M. A., Vladimirov, V. V., Sur, H. I., Markov, A. A., Özsoy, E., Kotovshchikov, B. B., Eremeev, V. V., Ünlüata, Ü., Circulation in the surface and intermediate layers of the Black Sea. Deep Sea Research I, 40(8): 1597-1612. 1993.
125. Sur, H. I., Ilyin, Y. P., Özsoy, E., Ünlüata, Ü., The impacts of continental Shelf. Deep Water Interactions in the Black Sea. J. Mar. Sys., 7, 293-320. 1996.
126. Vinogradov, M. Ye., Flint, M. V., Shushkina, E. A., Vertical distribution of mesoplankton in the open area of the Black Sea. Marine Biology 89: 95-107. 1985.
127. Kovaleva, T. M., Shumakova, G. V., Skryabin, V. A., Ostrovskaya, N. A., Plankton of the redox-zone. Composition and distribution of main components. In: Plankton of the Black Sea (eds. Kovalev, A.V., Finenko, Z. Z.) Kiev: Naukova dumka: 216-245. 1993. (in Russian).
128. Chokhuri, N. I., To the question on diurnal vertical migrations of plankton in the Batumi Bay. Pr. Sci. Fisheries Biol. St. of Georgia. 2: 153-172. 1939. (in Russian).
129. Petipa, T. S., Observations of zooplankton behaviour during solar eclipse. Rep. Ac. Sci. USSR. 104, N2: 323. 1965. (in Russian).
130. Zagorodnyaya, Yu. A., Vertical migrations and diurnal ration of Copepoda Crustacea *Pseudocalanus elongatus* (Boeck) in the Black Sea. Biologiya morya, 33: 11-18. 1975. (in Russian).
131. Konsulov, A., Daily vertical migration of zooplankton in the Bulgarian Black Sea coastal region. Oceanology, 19: 35-48. 1990.
132. Kovalev, A. V., Zagorodnyaya, Yu. A., Diurnal variability of quantitative indices of marine zooplankton in whole layer of inhabitation. DAN USSR. Ser. B. N9: 71-72. 1985. (in Russian).
133. Zagorodnyaya, Yu. A., Kovalev, A. V., Diurnal rhythm of reproduction and ontogenetic changes of marine Copepoda. Gidrobiol. Zhurn. 24, N2: 20-23. 1988. (in Russian).
134. Zagorodnyaya, Yu. A., Kovalev, A. V., Size structure of plankton zoocenosis of the open Black Sea. Ecologiya morya, 31: 1-6. 1989. (in Russian).
135. Nikitin, V. N., Plankton of the Batumi Bay and its annual quantitative changes. Book in honour of Prof. N. M. Knipovich. M.: L.: Pischepromizdat, 63-86. 1939. (in Russian).
136. Lazareva, L. P., To question on seasonal dynamics of biomass of the Black Sea zooplankton in Karadag region. Proc. of Karadag Biol. St., 14: 127-134. 1957. (in Russian).
137. Sazhina, L. I., Seasonal changes of zooplankton in the north-western part of the Black Sea in 1957. Proc. of Sevast. Biol. St., 17: 262-275. 1964. (in Russian).

138. Delalo, E. P., Baldina, E. P., Bileva, O. K., Seasonal changes of zooplankton distribution in the western part of the Black Sea in 1957. Investigations of plankton of the Black and Azov Seas. Kiev, Naukova dumka: 92-101. 1965, (in Russian).
139. Pasternak, A. F., Seasonal dynamics of numbers and biomass of zooplankton. Seasonal changes of the Black Sea zooplankton. Moscow, Nauka: 139-177. 1983, (in Russian).
140. Konsulov, A., Seasonal and annual dynamics of zooplankton in the Black Sea along the Bulgarian coast for the 1974-1984 period. Oceanology, 16: 23-33. 1986.
141. Slavin, L. B. Temperature regime. Hydrometeorological reference book of the Azov Sea. L.: Nauka, 160-267. 1962, (in Russian).
142. Nesterova, D. A., Phytoplanktonic communities of the shelf ecosystem. System analysis and processes modelling on the Black Sea shelf. Sevastopol: Ed. MHI Ac. Sci. UkrSSR.: 102-106. 1983, (in Russian).
143. Gomou, M. T., Kupriyanov, S. S., Estimation of abundance and distribution of medusae *Aurelia aurita* in the eastern part of the Black Sea. M.: Nauka, 191-198. 1980, (in Russian).
144. Zaitsev, Yu. P., Polischuk, L. N., Outburst of abundance of medusae *Aurelia aurita* (L.) in the Black Sea. Ecologiya morya. N17: 35-46. 1984, (in Russian).
145. Shushkina, E. A., Vinogradov, M. E. Changes of planktonic community in the open Black Sea and influence on it by ctenophore *Mnemiopsis* (1978-1989). Changeability of the Black Sea ecosystem. Natural and anthropogenic factors. (ed. Vinogradov M. E.), Moscow, Nauka: 248-261. 1991, (in Russian).
146. Shiganova, T. A., Kideyş, A. E., Gücü, A. C., Niermann, U., Khoroshilov, V. S., Changes in species diversity and abundance of the main components of the Black Sea pelagic community during the last decade. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 1: pp. 171-188, 1998.
147. Konsulov, A., Kamburska, L., Black Sea zooplankton structural dynamic and variability off the Bulgarian Black Sea coast during 1991-1995. In: NATO TU Black Sea project: Ecosystem Modeling as a Management Tool for the Black Sea, Symposium on Sci. Results, L. Ivanov & T. Oğuz (eds.), Kluwer Acad. Publ. Vol. 1: pp. 281-292, 1998.
148. Zaika, V. E., Sergeeva, N. G., Diurnal dynamics of population structure and vertical distribution of ctenophore *Mnemiopsis maccradyi* MAYER in the Black Sea. Zhurnal Obshchbiologii, Kiev, 27(2): 15-19. 1991, (in Russian).
149. Mutlu, E., Bingel, F., Gucu A. C., Melnikov V. V., Niermann, U., Ostrovskaya N. A., Zaika V. E., Distribution of the new invader *Mnemiopsis sp.* and the resident *Aurelia aurita* and *Pleurobrachia pileus* populations in the Black Sea in the years 1991-1993. ICES J. of Mar. Sci., 51: 407-421. 1994.
150. Vinogradov, M. E., Shushkina, E. A., Musaeva E. I., Nikolaeva, G. G., Vertical distribution of the Black Sea mesoplankton in winter 1991. In: Winter state of the open Black Sea ecosystem (ed. Vinogradov, M. E.), Moscow, IO RAN: 103-119, 1992, (in Russian).
151. Kovalev, A. V., Ostrovskaya, N. A., Skryabin, V. A., Zagorodnyaya, Yu. A., State of zooplankton, as a feed base of fishes in the Black Sea. Modern state of the Black Sea ichthyofauna (ed. Kononov, S.M.). Sevastopol: 131-152. 1995, (in Russian).
152. Vinogradov, M. E., Arnautov, G. N., Investigations of modern state of the Black Sea ecosystem Investigations of ecosystem of the Black Sea pelagial. M. Nauka, 1-8. 1986, (in Russian).
153. Tsikhon-Lukanina, Ye. A., Reznitchenko, O. G., Lukasheva, T. A., Quantitative patterns of feeding of the Black Sea ctenophore *Mnemiopsis leidyi*. Oceanology, 31: 196-199. 1991.
154. Tsikhon-Lukanina, Ye. A., Reznitchenko, O. G., Diet of the ctenophore *Mnemiopsis* in the Black Sea as a function of size. Oceanology, 31: 320-323. 1991.
155. Bogdanova, A. K., Vodyanitskiy, V. A., Cycleness of fluctuations of hydrological factors and fisheries in the Azov and Black Seas. Biologiya morya, v. 27: 3-19, 1972, (in Russian).
156. Zhorov, V. A., Boguslavskiy, S. G., Tendencies of some hydrological and hydrochemical processes in the Black Sea. Meteorologiya i gidrologiya. N11: 63-69. 1985, (in Russian).
157. Childers, D. L., Day, J. W., Muller, Jr., R. A., Relating climatological forcing to coastal water levels in Louisiana estuaries and the potential importance of El Niño-southern oscillation events. Climate Research 1: 31-42. 1990.
158. Man, K. H., Lazier, J. R. N., Fronts in coastal waters. In: Dynamics of marine ecosystems (Man, K. H., Lazier, J. R. N. eds), Blackwell Sci Publ.: 213-254. 1991.
159. Mann, K. H., Physical oceanography, food chain and fish stocks: A review, ICES J. of Mar. Sci. 50: 105-119. 1993.
160. Cushing, D. N., The long-term relationship between zooplankton and fish. JCES J. of Mar.Sci. N52: 611-626. 1995.
161. Alheit, J., Hagen, E., Long-term climate forcing of European herring and sardine populations. Fisheries Oceanography 6, 2: 130-139. 1997.
162. Özsoy, E., Ünlüata, Ü., Oceanography of the Black Sea: a review of some recent results. Earth-Science Reviews, 42: 231-272. 1997.

163. Polonsky, A., Voskresenskaya, E., Belokopytov, V., Variability of northwestern Black Sea Hydrography and the river discharges as part of global ocean-atmosphere fluctuations. In: Özsoy & Mikaelean (eds.): Sensitivity to change: Black Sea, Baltic Sea and North Sea. NATO ASI Series, 2. Environment-Vol.27. Kluwer Academic Publishers (pp: 516): 11-24. 1997.
164. Rodionov, S., Global and regional climate interaction: the Caspian Sea experience. Kluwer Academic Publishers, The Netherlands: 241 pp. 1994.
165. Schlesinger, M. E., Ramankutty, N., An oscillation in the global climate system of period 65-70 years. *Nature*, 367: 723-726. 1994.
166. Halpert, M. S., Ropelewski, C. F., Climate assessment. A decadal review, 1981-1990. Camp Springs, MD: US. Department of Commerce, NWS/NMC/CAC. 1991.
167. Trenberth, K. E., Hurrell, J. W., Decadal atmosphere-ocean variations in the North Pacific. *Climate Dynamic*, 9: 303-319, 1994.
168. Kideyş, A. E., Recent dramatic changes in the Black Sea ecosystems: The reason for the sharp decline in Turkish anchovy fisheries. *J. of Marine Systems* 5: 171-181. 1994.
169. Venrick, E. L., McGowan, I. A., Gowan, D. R., Hayward, T. R., Climate and chlorophyll a: long-term changes in the Central North Pacific Ocean. *Science*, 238: 70-72. 1987.