

# 2-Blocks with minimal nonabelian defect groups

Benjamin Sambale  
 Mathematisches Institut  
 Friedrich-Schiller-Universität  
 07743 Jena  
 Germany  
 benjamin.sambale@uni-jena.de

December 9, 2010

## Abstract

We study numerical invariants of 2-blocks with minimal nonabelian defect groups. These groups were classified by Rédei (see [41]). If the defect group is also metacyclic, then the block invariants are known (see [43]). In the remaining cases there are only two (infinite) families of “interesting” defect groups. In all other cases the blocks are nilpotent. We prove Brauer’s  $k(B)$ -conjecture and the Olsson-conjecture for all 2-blocks with minimal nonabelian defect groups. For one of the two families we also show that Alperin’s weight conjecture and Dade’s conjecture is satisfied. This paper is a part of the author’s PhD thesis.

**Keywords:** blocks of finite groups, minimal nonabelian defect groups, Alperin’s conjecture, Dade’s conjecture.

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Fusion systems</b>	<b>2</b>
<b>3</b>	<b>The case <math>r &gt; s = 1</math></b>	<b>3</b>
3.1	The $B$ -subsections . . . . .	3
3.2	The numbers $k(B)$ , $k_i(B)$ and $l(B)$ . . . . .	4
3.3	Generalized decomposition numbers . . . . .	6
3.4	The Cartan matrix . . . . .	9
3.5	Dade’s conjecture . . . . .	11
3.6	Alperin’s weight conjecture . . . . .	13
3.7	The gluing problem . . . . .	14
<b>4</b>	<b>The case <math>r = s &gt; 1</math></b>	<b>14</b>
4.1	The $B$ -subsections . . . . .	14
4.2	The gluing problem . . . . .	17
4.3	Special cases . . . . .	18

## 1 Introduction

Let  $R$  be a discrete complete valuation ring with quotient field  $K$  of characteristic 0. Moreover, let  $(\pi)$  be the maximal ideal of  $R$  and  $F := R/(\pi)$ . We assume that  $F$  is algebraically closed of characteristic 2. We fix a finite group  $G$ , and assume that  $K$  contains all  $|G|$ -th roots of unity. Let  $B$  be a block of  $RG$  with defect group  $D$ . We

denote the number of irreducible ordinary characters of  $B$  by  $k(B)$ . These characters split in  $k_i(B)$  characters of height  $i \in \mathbb{N}_0$ . Similarly, let  $k^i(B)$  be the number of characters of defect  $i \in \mathbb{N}_0$ . Finally, let  $l(B)$  be the number of irreducible Brauer characters of  $B$ . The defect group  $D$  is called *minimal nonabelian* if every proper subgroup of  $D$  is abelian, but not  $D$  itself. Rédei has shown that  $D$  is isomorphic to one of the following groups (see [41]):

- (i)  $\langle x, y \mid x^{2^r} = y^{2^s} = 1, xyx^{-1} = y^{1+2^{s-1}} \rangle$ , where  $r \geq 1$  and  $s \geq 2$ ,
- (ii)  $\langle x, y \mid x^{2^r} = y^{2^s} = [x, y]^2 = [x, x, y] = [y, x, y] = 1 \rangle$ , where  $r \geq s \geq 1$ ,  $[x, y] := xyx^{-1}y^{-1}$  and  $[x, x, y] := [x, [x, y]]$ ,
- (iii)  $Q_8$ .

In the first and last case  $D$  is also metacyclic. In this case  $B$  is well understood (see [43]). Thus, we may assume that  $D$  has the form (ii).

## 2 Fusion systems

To analyse the possible fusion systems on  $D$  we start with a group theoretical lemma.

**Lemma 2.1.** *Let  $z := [x, y]$ . Then the following hold:*

- (i)  $|D| = 2^{r+s+1}$ .
- (ii)  $\Phi(D) = Z(D) = \langle x^2, y^2, z \rangle \cong C_{2^{r-1}} \times C_{2^{s-1}} \times C_2$ .
- (iii)  $D' = \langle z \rangle \cong C_2$ .
- (iv)  $|\text{Irr}(D)| = 5 \cdot 2^{r+s-2}$ .
- (v) *If  $r = s = 1$ , then  $D \cong D_8$ . For  $r \geq 2$  the maximal subgroups of  $D$  are given by*

$$\begin{aligned} \langle x^2, y, z \rangle &\cong C_{2^{r-1}} \times C_{2^s} \times C_2, \\ \langle x, y^2, z \rangle &\cong C_{2^r} \times C_{2^{s-1}} \times C_2, \\ \langle xy, x^2, z \rangle &\cong C_{2^r} \times C_{2^{s-1}} \times C_2. \end{aligned}$$

We omit the (elementary) proof of this lemma. However, notice that  $|P'| = 2$  and  $|P : \Phi(P)| = |P : Z(P)| = p^2$  hold for every minimal nonabelian  $p$ -group  $P$ . Rédei has also shown that for different pairs  $(r, s)$  one gets nonisomorphic groups. This gives precisely  $\lfloor \frac{n-1}{2} \rfloor$  isomorphism classes of these groups of order  $2^n$ . For  $r \neq 1$  (that is  $|D| \geq 16$ ) the structure of the maximal subgroups shows that all these groups are nonmetacyclic.

Now we investigate the automorphism groups.

**Lemma 2.2.** *The automorphism group  $\text{Aut}(D)$  is a 2-group, if and only if  $r \neq s$  or  $r = s = 1$ .*

*Proof.* If  $r \neq s$  or  $r = s = 1$ , then there exists a characteristic maximal subgroup of  $D$  by Lemma 2.1(v). In these cases  $\text{Aut}(D)$  must be a 2-group. Thus, we may assume  $r = s \geq 2$ . Then one can show that the map  $x \mapsto y, y \mapsto x^{-1}y^{-1}$  is an automorphism of order 3.  $\square$

**Lemma 2.3.** *Let  $P \cong C_{2^{n_1}} \times \dots \times C_{2^{n_k}}$  with  $n_1, \dots, n_k, k \in \mathbb{N}$ . Then  $\text{Aut}(P)$  is a 2-group, if and only if the  $n_i$  are pairwise distinct.*

*Proof.* See for example Lemma 2.7 in [34].  $\square$

Now we are able to decide, when a fusion system on  $D$  is nilpotent.

**Theorem 2.4.** *Let  $\mathcal{F}$  be a fusion system on  $D$ . Then  $\mathcal{F}$  is nilpotent or  $s = 1$  or  $r = s$ . If  $r = s \geq 2$ , then  $\mathcal{F}$  is controlled by  $D$ .*

*Proof.* We assume  $s \neq 1$ . Let  $Q < D$  be an  $\mathcal{F}$ -essential subgroup. Since  $Q$  is also  $\mathcal{F}$ -centric, we get  $C_P(Q) = Q$ . This shows that  $Q$  is a maximal subgroup of  $D$ . By Lemma 2.1(v) and Lemma 2.3, one of the following holds:

- (i)  $r = 2$  ( $= s$ ) and  $Q \in \{\langle x^2, y, z \rangle, \langle x, y^2, z \rangle, \langle xy, x^2, z \rangle\}$ ,
- (ii)  $r > s = 2$  and  $Q \in \{\langle x, y^2, z \rangle, \langle xy, x^2, z \rangle\}$ ,
- (iii)  $r = s + 1$  and  $Q = \langle x^2, y, z \rangle$ .

In all cases  $\Omega(Q) \subseteq Z(P)$ . Let us consider the action of  $\text{Aut}_{\mathcal{F}}(Q)$  on  $\Omega(Q)$ . The subgroup  $1 \neq P/Q = N_P(Q)/C_P(Q) \cong \text{Aut}_P(Q) \leq \text{Aut}_{\mathcal{F}}(Q)$  acts trivially on  $\Omega(Q)$ . On the other hand every nontrivial automorphism of odd order acts nontrivially on  $\Omega(Q)$  (see for example 8.4.3 in [19]). Hence, the kernel of this action is a nontrivial normal 2-subgroup of  $\text{Aut}_{\mathcal{F}}(Q)$ . In particular  $O_2(\text{Aut}_{\mathcal{F}}(Q)) \neq 1$ . But then  $\text{Aut}_{\mathcal{F}}(Q)$  cannot contain a strongly 2-embedded subgroup.

This shows that there are no  $\mathcal{F}$ -essential subgroups. Now the claim follows from Lemma 2.2 and Alperin's fusion theorem.  $\square$

Now we consider a kind of converse. If  $r = s = 1$ , then there are nonnilpotent fusion systems on  $D$ . In the case  $r = s \geq 2$  one can construct a nonnilpotent fusion system with a suitable semidirect product (see Lemma 2.2). We show that there is also a nonnilpotent fusion system in the case  $r > s = 1$ .

**Proposition 2.5.** *If  $s = 1$ , then there exists a nonnilpotent fusion system on  $D$ .*

*Proof.* We may assume  $r \geq 2$ . Let  $A_4$  be the alternating group of degree 4, and let  $H := \langle \tilde{x} \rangle \cong C_{2^r}$ . Moreover, let  $\varphi : H \rightarrow \text{Aut}(A_4) \cong S_4$  such that  $\varphi_{\tilde{x}} \in \text{Aut}(A_4)$  has order 4. Write  $\tilde{y} := (12)(34) \in A_4$  and choose  $\varphi$  such that  $\varphi_{\tilde{x}}(\tilde{y}) := (13)(24)$ . Finally, let  $G := A_4 \rtimes_{\varphi} H$ . Since all 4-cycles in  $S_4$  are conjugate,  $G$  is uniquely determined up to isomorphism. Because  $[\tilde{x}, \tilde{y}] = (13)(24)(12)(34) = (14)(23)$ , we get  $\langle \tilde{x}, \tilde{y} \rangle \cong D$ . The fusion system  $\mathcal{F}_G(D)$  is nonnilpotent, since  $A_4$  (and therefore  $G$ ) is not 2-nilpotent.  $\square$

### 3 The case $r > s = 1$

Now we concentrate on the case  $r > s = 1$ , i.e.

$$D := \langle x, y \mid x^{2^r} = y^2 = [x, y]^2 = [x, x, y] = [y, x, y] = 1 \rangle$$

with  $r \geq 2$ . As before  $z := [x, y]$ . We also assume that  $B$  is a nonnilpotent block. By Lemma 2.2,  $\text{Aut}(D)$  is a 2-group, and the inertial index  $t(B)$  of  $B$  equals 1.

#### 3.1 The $B$ -subsections

Olsson has already obtained the conjugacy classes of so called  $B$ -subsections (see [34]). However, his results contain errors. For example he missed the necessary relations  $[x, x, y]$  and  $[y, x, y]$  in the definition of  $D$ .

In the next lemma we denote by  $\text{Bl}(RH)$  the set of blocks of a finite group  $H$ . If  $H \leq G$  and  $b \in \text{Bl}(RH)$ , then  $b^G$  is the Brauer correspondent of  $b$  (if exists). Moreover, we use the notion of subpairs and subsections (see [36]).

**Lemma 3.1.** *Let  $b \in \text{Bl}(RDC_G(D))$  be a Brauer correspondent of  $B$ . For  $Q \leq D$  let  $b_Q \in \text{Bl}(RQC_G(Q))$  such that  $(Q, b_Q) \leq (D, b)$ . Set  $\mathcal{T} := Z(D) \cup \{x^i y^j : i, j \in \mathbb{Z}, i \text{ odd}\}$ . Then*

$$\bigcup_{a \in \mathcal{T}} \left\{ (a, b_{C_D(a)}^{C_G(a)}) \right\}$$

*is a system of representatives for the conjugacy classes of  $B$ -subsections. Moreover,  $|\mathcal{T}| = 2^{r+1}$ .*

*Proof.* If  $r = 2$ , then the claim follows from Proposition 2.14 in [34]. For  $r \geq 3$  the same argument works. However, Olsson refers wrongly to Proposition 2.11 (the origin of this mistake already lies in Lemma 2.8).  $\square$

From now on we write  $b_a := b_{C_D(a)}^{C_G(a)}$  for  $a \in \mathcal{T}$ .

**Lemma 3.2.** *Let  $P \cong C_{2^s} \times C_2^2$  with  $s \in \mathbb{N}$ , and let  $\alpha$  be an automorphism of  $P$  of order 3. Then  $C_P(\alpha) := \{b \in P : \alpha(b) = b\} \cong C_{2^s}$ .*

*Proof.* We write  $P = \langle a \rangle \times \langle b \rangle \times \langle c \rangle$  with  $|\langle a \rangle| = 2^s$ . It is well known that the kernel of the restriction map  $\text{Aut}(P) \rightarrow \text{Aut}(P/\Phi(P))$  is a 2-group. Since  $|\text{Aut}(P/\Phi(P))| = |\text{GL}(3, 2)| = 168 = 2^3 \cdot 3 \cdot 7$ , it follows that  $|\text{Aut}(P)|$  is divisible by 3 only once. In particular every automorphism of  $P$  of order 3 is conjugate to  $\alpha$  or  $\alpha^{-1}$ . Thus, we may assume  $\alpha(a) = a$ ,  $\alpha(b) = c$  and  $\alpha(c) = bc$ . Then  $C_P(\alpha) = \langle a \rangle \cong C_{2^s}$ .  $\square$

### 3.2 The numbers $k(B)$ , $k_i(B)$ and $l(B)$

The next step is to determine the numbers  $l(b_a)$ . The case  $r = 2$  needs special attention, because in this case  $D$  contains an elementary abelian maximal subgroup of order 8. We denote the inertial group of a block  $b \in \text{Bl}(RH)$  with  $H \trianglelefteq G$  by  $\text{T}_G(b)$ .

**Lemma 3.3.** *There is an element  $c \in \text{Z}(D)$  of order  $2^{r-1}$  such that  $l(b_a) = 1$  for all  $a \in \mathcal{T} \setminus \langle c \rangle$ .*

*Proof.*

**Case 1:**  $a \in \text{Z}(D)$ .

Then  $b_a = b_D^{C_G(a)}$  is a block with defect group  $D$  and Brauer correspondent  $b_D \in \text{Bl}(RD C_{C_G(a)}(D))$ . Let  $M := \langle x^2, y, z \rangle \cong C_{2^{r-1}} \times C_2^2$ . Since  $B$  is nonnilpotent, there exists an element  $\alpha \in \text{T}_{\text{N}_G(M)}(b_M)$  such that  $\alpha C_G(M) \in \text{T}_{\text{N}_G(M)}(b_M)/C_G(M)$  has order  $q \in \{3, 7\}$ . We will exclude the case  $q = 7$ . In this case  $r = 2$  and  $\text{T}_{\text{N}_G(M)}(b_M)/C_G(M)$  is isomorphic to a subgroup of  $\text{Aut}(M) \cong \text{GL}(3, 2)$ . Since

$$(M, {}^d b_M) = {}^d(M, b_M) \leq {}^d(D, b_D) = (D, b_D)$$

for all  $d \in D$ , we have  $D \subseteq \text{T}_{\text{N}_G(M)}(b_M)$ . This implies  $\text{T}_{\text{N}_G(M)}(b_M)/C_G(M) \cong \text{GL}(3, 2)$ , because  $\text{GL}(3, 2)$  is simple. By Satz 1 in [2], this contradicts the fact that  $\text{T}_{\text{N}_G(M)}(b_M)/C_G(M)$  contains a strongly 2-embedded subgroup (of course this can be shown “by hand” without invoking [2]). Thus, we have shown  $q = 3$ . Now

$$\text{T}_{\text{N}_G(M)}(b_M)/C_G(M) \cong S_3$$

follows easily. By Lemma 3.2 there is an element  $c := x^{2i} y^j z^k \in C_M(\alpha)$  ( $i, j, k \in \mathbb{Z}$ ) of order  $2^{r-1}$ . Let us assume that  $j$  is odd. Since  $x\alpha x \equiv x\alpha x^{-1} \equiv \alpha^{-1} \pmod{C_G(M)}$  we get

$$\begin{aligned} \alpha(x^{2i} y^j z^{k+1}) \alpha^{-1} &= \alpha x (x^{2i} y^j z^k) x^{-1} \alpha^{-1} = x \alpha^{-1} (x^{2i} y^j z^k) \alpha x^{-1} \\ &= x (x^{2i} y^j z^k) x^{-1} = x^{2i} y^j z^{k+1}. \end{aligned}$$

But this contradicts Lemma 3.2. Hence, we have proved that  $j$  is even. In particular  $c \in \text{Z}(D)$ . For  $a \notin \langle c \rangle$  we have  $\alpha \notin C_G(a)$  and  $l(b_a) = 1$ . While in the case  $a \in \langle c \rangle$  we get  $\alpha \in C_G(a)$ , and  $b_a$  is nonnilpotent. Thus, in this case  $l(b_a)$  remains unknown.

**Case 2:**  $a \notin \text{Z}(D)$ .

Let  $C_D(a) = \langle \text{Z}(D), a \rangle =: M$ . Since  $(M, b_M)$  is a Brauer subpair,  $b_M$  has defect group  $M$ . It follows from  $(M, b_M) \trianglelefteq (D, b_D)$  that also  $b_a$  has defect group  $M$  and Brauer correspondent  $b_M$ . In case  $M \cong C_{2^r} \times C_2$  we get  $l(b_a) = 1$ . Now let us assume  $M \cong C_{2^{r-1}} \times C_2^2$ . As in the first case, we choose  $\alpha \in \text{T}_{\text{N}_G(M)}(b_M)$  such that  $\alpha C_G(M) \in \text{T}_{\text{N}_G(M)}(b_M)/C_G(M)$  has order 3. Since  $a \notin \text{Z}(D)$ , we derive  $\alpha \notin C_G(a)$  and  $l(b_a) = 1$ .  $\square$

We denote by  $\text{IBr}(b_u) := \{\varphi_u\}$  for  $u \in \mathcal{T} \setminus \langle c \rangle$  the irreducible Brauer character of  $b_u$ . Then the generalized decomposition numbers  $d_{\chi\varphi_u}^u$  for  $\chi \in \text{Irr}(B)$  form a column  $d(u)$ . Let  $2^k$  be the order of  $u$ , and let  $\zeta := \zeta_{2^k}$  be a primitive  $2^k$ -th root of unity. Then the entries of  $d(u)$  lie in the ring of integers  $\mathbb{Z}[\zeta]$ . Hence, there exist integers  $a_i^u(\chi) \in \mathbb{Z}$  such that

$$d_{\chi\varphi_u}^u = \sum_{i=0}^{2^k-1} a_i^u(\chi) \zeta^i.$$

We expand this by

$$a_{i+2^{k-1}}^u := -a_i^u$$

for all  $i \in \mathbb{Z}$ .

Let  $|G| = 2^a m$  where  $2 \nmid m$ . We may assume  $\mathbb{Q}(\zeta_{|G|}) \subseteq K$ . Then  $\mathbb{Q}(\zeta_{|G|})|\mathbb{Q}(\zeta_m)$  is a Galois extension, and we denote the corresponding Galois group by

$$\mathcal{G} := \text{Gal}(\mathbb{Q}(\zeta_{|G|})|\mathbb{Q}(\zeta_m)).$$

Restriction gives an isomorphism

$$\mathcal{G} \cong \text{Gal}(\mathbb{Q}(\zeta_{2^a})|\mathbb{Q}).$$

In particular  $|\mathcal{G}| = 2^{a-1}$ . For every  $\gamma \in \mathcal{G}$  there is a number  $\tilde{\gamma} \in \mathbb{N}$  such that  $\gcd(\tilde{\gamma}, |G|) = 1$ ,  $\tilde{\gamma} \equiv 1 \pmod{m}$ , and  $\gamma(\zeta_{|G|}) = \zeta_{|G|}^{\tilde{\gamma}}$  hold. Then  $\mathcal{G}$  acts on the set of subsections by

$$\gamma(u, b) := (u^{\tilde{\gamma}}, b).$$

For every  $\gamma \in \mathcal{G}$  we get

$$d(u^{\tilde{\gamma}}) = \sum_{s \in \mathcal{S}} a_s^u \zeta_{2^k}^{s\tilde{\gamma}}$$

for every system  $\mathcal{S}$  of representatives of the cosets of  $2^{k-1}\mathbb{Z}$  in  $\mathbb{Z}$ . It follows that

$$a_s^u = 2^{1-a} \sum_{\gamma \in \mathcal{G}} d(u^{\tilde{\gamma}}) \zeta_{2^k}^{-\tilde{\gamma}s} \quad (1)$$

for  $s \in \mathcal{S}$ .

Now let  $u \in \mathcal{T} \setminus Z(D)$  and  $M := C_D(u)$ . Then  $b_u$  and  $b_M^{\text{TN}_G(M)(b_M) \cap \text{N}_G(\langle u \rangle)}$  have  $M$  as defect group, because  $D \not\subseteq \text{N}_G(\langle u \rangle)$ . By (6B) in [6] it follows that the  $2^{r-1}$  distinct  $B$ -subsections of the form  $\gamma(u, b_u)$  with  $\gamma \in \mathcal{G}$  are pairwise nonconjugate. The same holds for  $u \in Z(D) \setminus \{1\}$ . Using this and equation (1) we can adapt Lemma 3.9 in [33]:

**Lemma 3.4.** *Let  $c \in Z(D)$  as in Lemma 3.3, and let  $u, v \in \mathcal{T} \setminus \langle c \rangle$  with  $|\langle u \rangle| = 2^k$  and  $|\langle v \rangle| = 2^l$ . Moreover, let  $i \in \{0, 1, \dots, 2^{k-1} - 1\}$  and  $j \in \{0, 1, \dots, 2^{l-1} - 1\}$ . If there exist  $\gamma \in \mathcal{G}$  and  $g \in G$  such that  $g(u, b_u) = \gamma(v, b_v)$ , then*

$$(a_i^u, a_j^v) = \begin{cases} 2^{d(B)-k+1} & \text{if } u \in Z(D) \text{ and } j\tilde{\gamma} - i \equiv 0 \pmod{2^k} \\ -2^{d(B)-k+1} & \text{if } u \in Z(D) \text{ and } j\tilde{\gamma} - i \equiv 2^{k-1} \pmod{2^k} \\ 2^{d(B)-k} & \text{if } u \notin Z(D) \text{ and } j\tilde{\gamma} - i \equiv 0 \pmod{2^k} \\ -2^{d(B)-k} & \text{if } u \notin Z(D) \text{ and } j\tilde{\gamma} - i \equiv 2^{k-1} \pmod{2^k} \\ 0 & \text{otherwise} \end{cases}.$$

Otherwise  $(a_i^u, a_j^v) = 0$ . In particular  $(a_i^u, a_j^v) = 0$  if  $k \neq l$ .

Using the theory of contributions we can also carry over Lemma (6.E) in [20]:

**Lemma 3.5.** *Let  $u \in Z(D)$  with  $l(b_u) = 1$ . If  $u$  has order  $2^k$ , then for every  $\chi \in \text{Irr}(B)$  holds:*

$$(i) \quad 2^{h(\chi)} \mid a_i^u(\chi) \text{ for } i = 0, \dots, 2^{k-1} - 1,$$

$$(ii) \quad \sum_{i=0}^{2^{k-1}-1} a_i^u(\chi) \equiv 2^{h(\chi)} \pmod{2^{h(\chi)+1}}.$$

By Lemma 1.1 in [39] we have

$$k(B) \leq \sum_{i=0}^{\infty} 2^{2i} k_i(B) \leq |D|. \quad (2)$$

In particular Brauer's  $k(B)$ -conjecture holds. Olsson's conjecture

$$k_0(B) \leq |D : D'| = 2^{r+1} \quad (3)$$

follows by Theorem 3.1 in [39]. Now we are able to calculate the numbers  $k(B)$ ,  $k_i(B)$  and  $l(B)$ .

**Theorem 3.6.** *We have*

$$k(B) = 5 \cdot 2^{r-1} = |\text{Irr}(D)|, \quad k_0(B) = 2^{r+1} = |D : D'|, \quad k_1(B) = 2^{r-1}, \quad l(B) = 2.$$

*Proof.* We argue by induction on  $r$ . Let  $r = 2$ , and let  $c \in Z(D)$  as in Lemma 3.3. By way of contradiction we assume  $c = z$ . If  $\alpha$  and  $M$  are defined as in the proof of Lemma 3.3, then  $\alpha$  acts nontrivially on  $M/\langle z \rangle \cong C_2^2$ . On the other hand  $x$  acts trivially on  $M/\langle z \rangle$ . This contradicts  $x\alpha x^{-1}\alpha \in C_G(M)$ .

This shows  $c \in \{x^2, x^2z\}$  and  $D/\langle c \rangle \cong D_8$ . Thus, we can apply Theorem 2 in [8]. For this let

$$M_1 := \begin{cases} \langle x, z \rangle & \text{if } c = x^2 \\ \langle xy, z \rangle & \text{if } c = x^2z \end{cases}.$$

Then  $M \neq M_1 \cong C_4 \times C_2$  and  $\overline{M} := M/\langle c \rangle \cong C_2^2 \cong M_1/\langle c \rangle =: \overline{M}_1$ . Let  $\beta$  be the block of  $R\overline{C_G(c)} := R[C_G(c)/\langle c \rangle]$  which is dominated by  $b_c$ . By Theorem 1.5 in [33] we have

$$3 \mid |\text{T}_{\overline{C_G(c)}(\overline{M})}(\beta_{\overline{M}})/\overline{C_{\overline{C_G(c)}}(\overline{M})}|$$

and

$$3 \nmid |\text{T}_{\overline{C_G(c)}(\overline{M}_1)}(\beta_{\overline{M}_1})/\overline{C_{\overline{C_G(c)}}(\overline{M}_1)}|,$$

where  $(\overline{M}, \beta_{\overline{M}})$  and  $(\overline{M}_1, \beta_{\overline{M}_1})$  are  $\beta$ -subpairs. This shows that case (ab) in Theorem 2 in [8] occurs. Hence,  $l(b_c) = l(\beta) = 2$ . Now Lemma 3.3 yields

$$k(B) \geq 1 + k(B) - l(B) = 9.$$

It is well known that  $k_0(B)$  is divisible by 4. Thus, the equations (2) and (3) imply  $k_0(B) = 8$ . Moreover,

$$d_{\chi\varphi_z}^z = a_0^z(\chi) = \pm 1$$

holds for every  $\chi \in \text{Irr}(B)$  with  $h(\chi) = 0$ . This shows  $4k_1(B) \leq |D| - k_0(B) = 8$ . It follows that  $k_1(B) = l(B) = 2$ .

Now we consider the case  $r \geq 3$ . Since  $z$  is not a square in  $D$ , we have  $z \notin \langle c \rangle$ . Let  $a \in \langle c \rangle$  such that  $|\langle a \rangle| = 2^k$ . If  $k = r - 1$ , then  $l(b_a) = 2$  as before. Now let  $k < r - 1$ . Then  $D/\langle a \rangle$  has the same isomorphism type as  $D$ , but one has to replace  $r$  by  $r - k$ . By induction we get  $l(b_a) = 2$  for  $k \geq 1$ . This shows

$$k(B) \geq 1 + k(B) - l(B) = 2^{r+1} + 2^{r-1} - 1.$$

Equation (2) yields

$$\begin{aligned} 2^{r+2} - 4 &= 2^{r+1} + 4(2^{r-1} - 1) \leq k_0(B) + 4(k(B) - k_0(B)) \\ &\leq \sum_{i=0}^{\infty} 2^{2i} k_i(B) \leq |D| = 2^{r+2}. \end{aligned}$$

Now the conclusion follows easily. □

As a consequence, Brauer's height zero conjecture and the Alperin-McKay-conjecture hold for  $B$ .

### 3.3 Generalized decomposition numbers

Now we will determine some of the generalized decomposition numbers. Again let  $c \in Z(D)$  as in Lemma 3.3, and let  $u \in Z(D) \setminus \langle c \rangle$  with  $|\langle u \rangle| = 2^k$ . Then  $(a_i^u, a_i^u) = 2^{r+3-k}$  and  $2 \mid a_i^u(\chi)$  for  $h(\chi) = 1$  and  $i = 0, \dots, 2^{k-1} - 1$ . This gives

$$|\{\chi \in \text{Irr}(B) : a_i^u(\chi) \neq 0\}| \leq 2^{r+3-k} - 3|\{\chi \in \text{Irr}(B) : h(\chi) = 1, a_i^u(\chi) \neq 0\}|.$$

Moreover, for every character  $\chi \in \text{Irr}(B)$  there exists  $i \in \{0, \dots, 2^{k-1} - 1\}$  such that  $a_i^u(\chi) \neq 0$ . Hence,

$$\begin{aligned} k(B) &\leq \sum_{i=0}^{2^{k-1}-1} \sum_{\substack{\chi \in \text{Irr}(B), \\ a_i^u(\chi) \neq 0}} 1 \leq \sum_{i=0}^{2^{k-1}-1} \left( 2^{r+3-k} - 3 \sum_{\substack{\chi \in \text{Irr}(B), \\ h(\chi)=1, \\ a_i^u(\chi) \neq 0}} 1 \right) = |D| - 3 \sum_{i=0}^{2^{k-1}-1} \sum_{\substack{\chi \in \text{Irr}(B), \\ h(\chi)=1, \\ a_i^u(\chi) \neq 0}} 1 \\ &\leq |D| - 3k_1(B) = k(B). \end{aligned}$$

This shows that for every  $\chi \in \text{Irr}(B)$  there exists  $i(\chi) \in \{0, \dots, 2^{k-1} - 1\}$  such that

$$d_{\chi\varphi_u}^u = \begin{cases} \pm \zeta_{2^k}^{i(\chi)} & \text{if } h(\chi) = 0 \\ \pm 2\zeta_{2^k}^{i(\chi)} & \text{if } h(\chi) = 1 \end{cases}.$$

In particular

$$d_{\chi\varphi_u}^u = a_0^u(\chi) = \begin{cases} \pm 1 & \text{if } h(\chi) = 0 \\ \pm 2 & \text{if } h(\chi) = 1 \end{cases}$$

for  $k = 1$ .

By Lemma 3.4 we have  $(a_i^u, a_i^u) = 4$  for  $u \in \mathcal{T} \setminus Z(D)$  and  $i = 0, \dots, 2^{r-1} - 1$ . If  $a_i^u$  has only one nonvanishing entry, then  $a_i^u$  would not be orthogonal to  $a_{\tilde{j}}$ . Hence,  $a_i^u$  has up to ordering the form

$$(\pm 1, \pm 1, \pm 1, \pm 1, 0, \dots, 0)^T,$$

where the signs are independent of each other. The proof of Theorem 3.1 in [39] gives

$$|d_{\chi\varphi_u}^u| = 1$$

for  $u \in \mathcal{T} \setminus Z(D)$  and  $\chi \in \text{Irr}(B)$  with  $h(\chi) = 0$ . In particular  $d_{\chi\varphi_u}^u = 0$  for characters  $\chi \in \text{Irr}(B)$  of height 1. By suitable ordering we get

$$a_i^u(\chi_j) = \begin{cases} \pm 1 & \text{if } j - 4i \in \{1, \dots, 4\} \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad d_{\chi_j\varphi_u}^u = \begin{cases} \pm \zeta_{2^r}^{\lfloor \frac{j-1}{4} \rfloor} & \text{if } 1 \leq j \leq k_0(B) \\ 0 & \text{if } k_0(B) < j \leq k(B) \end{cases}$$

for  $i = 0, \dots, 2^{r-1} - 1$ , where  $\chi_1, \dots, \chi_{k_0(B)}$  are the characters of height 0.

Now let  $\text{IBr}(b_c) := \{\varphi_1, \varphi_2\}$ . We determine the numbers  $d_{\chi\varphi_1}^c, d_{\chi\varphi_2}^c \in \mathbb{Z}[\zeta_{2^{r-1}}]$ . By (4C) in [6] we have  $d_{\chi\varphi_1}^c \neq 0$  or  $d_{\chi\varphi_2}^c \neq 0$  for all  $\chi \in \text{Irr}(B)$ . As in the proof of Theorem 3.6,  $b_c$  dominates a block  $\overline{b_c} \in \text{Bl}(R[C_G(c)/\langle c \rangle])$  with defect group  $D_8$ . The table at the end of [14] shows that the Cartan matrix of  $\overline{b_c}$  has the form

$$\begin{pmatrix} 8 & 4 \\ 4 & 3 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 4 & 2 \\ 2 & 3 \end{pmatrix}.$$

We label these possibilities as the ‘‘first’’ and the ‘‘second’’ case. The Cartan matrix of  $b_c$  is

$$2^{r-1} \begin{pmatrix} 8 & 4 \\ 4 & 3 \end{pmatrix} \quad \text{or} \quad 2^{r-1} \begin{pmatrix} 4 & 2 \\ 2 & 3 \end{pmatrix}$$

respectively. The inverses of these matrices are

$$2^{-r-2} \begin{pmatrix} 3 & -4 \\ -4 & 8 \end{pmatrix} \quad \text{and} \quad 2^{-r-2} \begin{pmatrix} 3 & -2 \\ -2 & 4 \end{pmatrix}.$$

Let  $m_{\chi\psi}^{(c,b_c)}$  be the contribution of  $\chi, \psi \in \text{Irr}(B)$  with respect to the subsection  $(c, b_c)$  (see [6]). Then we have

$$\begin{aligned} |D|m_{\chi\psi}^{(c,b_c)} &= 3d_{\chi\varphi_1}^c \overline{d_{\psi\varphi_1}^c} - 4(d_{\chi\varphi_1}^c \overline{d_{\psi\varphi_2}^c} + d_{\chi\varphi_2}^c \overline{d_{\psi\varphi_1}^c}) + 8d_{\chi\varphi_2}^c \overline{d_{\psi\varphi_2}^c} \\ &\quad \text{or} \\ |D|m_{\chi\psi}^{(c,b_c)} &= 3d_{\chi\varphi_1}^c \overline{d_{\psi\varphi_1}^c} - 2(d_{\chi\varphi_1}^c \overline{d_{\psi\varphi_2}^c} + d_{\chi\varphi_2}^c \overline{d_{\psi\varphi_1}^c}) + 4d_{\chi\varphi_2}^c \overline{d_{\psi\varphi_2}^c} \end{aligned} \quad (4)$$

respectively. For a character  $\chi \in \text{Irr}(B)$  with height 0 we get

$$0 = h(\chi) = \nu(|D|m_{\chi\chi}^{(c,b_c)}) = \nu(3d_{\chi\varphi_1}^c \overline{d_{\chi\varphi_1}^c}) = \nu(d_{\chi\varphi_1}^c)$$

by (5H) in [6]. In particular  $d_{\chi\varphi_1}^c \neq 0$ . We define  $c_i^j \in \mathbb{Z}^{k(B)}$  by

$$d_{\chi\varphi_j}^c = \sum_{i=0}^{2^{r-2}-1} c_i^j(\chi) \zeta_{2^{r-1}}^i$$

for  $j = 1, 2$ . Then

$$(c_i^1, c_j^1) = \begin{cases} \delta_{ij} 16 & \text{first case} \\ \delta_{ij} 8 & \text{second case} \end{cases}, \quad (c_i^1, c_j^2) = \begin{cases} \delta_{ij} 8 & \text{first case} \\ \delta_{ij} 4 & \text{second case} \end{cases}, \quad (c_i^2, c_j^2) = \delta_{ij} 6$$

as in Lemma 3.4. (Since the  $2^{r-2}$   $B$ -subsections of the form  $\gamma(c, b_c)$  for  $\gamma \in \mathcal{G}$  are pairwise nonconjugate, one can argue like in Lemma 3.4.) Hence, in the second case

$$d_{\chi_i\varphi_1}^c = \begin{cases} \pm \zeta_{2^{r-1}}^{\lfloor \frac{i-1}{8} \rfloor} & \text{if } 1 \leq i \leq k_0(B) \\ 0 & \text{if } k_0(B) < i \leq k(B) \end{cases} \quad (\text{second case})$$

holds for a suitable arrangement. Again  $\chi_1, \dots, \chi_{k_0(B)}$  are the characters of height 0. In the first case

$$1 = h(\psi) = \nu(|D|m_{\chi\psi}^{(c,b_c)}) = \nu(3d_{\chi\varphi_1}^c \overline{d_{\psi\varphi_1}^c}) = \nu(d_{\psi\varphi_1}^c)$$

by (5G) in [6] for  $h(\psi) = 1$  and  $h(\chi) = 0$ . As in Lemma 3.5 we also have  $2 \mid c_i^1(\psi)$  for  $h(\psi) = 1$  and  $i = 0, \dots, 2^{r-2} - 1$ . Analogously as in the case  $u \in Z(D) \setminus \langle c \rangle$  we conclude

$$d_{\chi\varphi_1}^c = \begin{cases} \pm \zeta_{2^{r-1}}^{i(\chi)} & \text{if } h(\chi) = 0 \\ \pm 2 \zeta_{2^{r-1}}^{i(\chi)} & \text{if } h(\chi) = 1 \end{cases} \quad (\text{first case}) \quad (5)$$

for suitable indices  $i(\chi) \in \{0, \dots, 2^{r-2} - 1\}$ . Since  $(c_i^2, c_j^2) = \delta_{ij} 6$ , in both cases  $c_i^2$  has the form

$$(\pm 1, \pm 1, \pm 1, \pm 1, \pm 1, \pm 1, \pm 1, 0, \dots, 0)^T \text{ or } (\pm 2, \pm 1, \pm 1, 0, \dots, 0)^T.$$

We show that the latter possibility does not occur. In the second case for every character  $\chi \in \text{Irr}(B)$  with height 1 there exists  $i \in \{0, \dots, 2^{r-2} - 1\}$  such that  $c_i^2(\chi) \neq 0$ . In this case we get

$$d_{\chi_i\varphi_2}^c = \begin{cases} \pm \zeta_{2^{r-1}}^{\lfloor \frac{i-1}{4} \rfloor} & \text{if } 1 \leq i \leq 2^r \\ 0 & \text{if } 2^r < i \leq k_0(B) \\ \pm \zeta_{2^{r-1}}^{\lfloor \frac{i-k_0(B)-1}{2} \rfloor} & \text{if } k_0(B) < i \leq k(B) \end{cases} \quad (\text{second case}),$$

where  $\chi_1, \dots, \chi_{k_0(B)}$  are again the characters of height 0. Now let us consider the first case. Since  $(c_i^1, c_j^2) = \delta_{ij} 8$ , the value  $\pm 2$  must occur in every column  $c_i^1$  for  $i = 0, \dots, 2^{r-2} - 1$  at least twice. Obviously exactly two entries have to be  $\pm 2$ . Thus, one can improve equation (5) to

$$d_{\chi_i\varphi_1}^c = \begin{cases} \pm \zeta_{2^{r-1}}^{\lfloor \frac{i-1}{8} \rfloor} & \text{if } 1 \leq i \leq k_0(B) \\ \pm 2 \zeta_{2^{r-1}}^{\lfloor \frac{i-k_0(B)-1}{2} \rfloor} & \text{if } k_0(B) < i \leq k(B) \end{cases} \quad (\text{first case}).$$

It follows

$$d_{\chi_i\varphi_2}^c = \begin{cases} \pm \zeta_{2^{r-1}}^{\lfloor \frac{i-1}{4} \rfloor} & \text{if } 1 \leq i \leq 2^r \\ 0 & \text{if } 2^r < i \leq k_0(B) \\ \pm \zeta_{2^{r-1}}^{\lfloor \frac{i-k_0(B)-1}{2} \rfloor} & \text{if } k_0(B) < i \leq k(B) \end{cases} \quad (\text{first case}).$$

Hence, the numbers  $d_{\chi\varphi_2}^c$  are independent of the case. Of course, one gets similar results for  $d_{\chi\varphi_i}^u$  with  $\langle u \rangle = \langle c \rangle$ .



### 3.4 The Cartan matrix

Now we investigate the Cartan matrix of  $B$ .

**Lemma 3.7.** *The elementary divisors of the Cartan matrix of  $B$  are  $2^{r-1}$  and  $|D|$ .*

*Proof.* Let  $C$  be the Cartan matrix of  $B$ . Since  $l(B) = 2$ , it suffices to show that  $2^{r-1}$  occurs as elementary divisor of  $C$  at least once. In order to prove this, we use the notion of lower defect groups (see [35]). Let  $(u, b)$  be a  $B$ -subsection with  $|\langle u \rangle| = 2^{r-1}$  and  $l(b) = 2$ . Let  $b_1 := b^{\text{N}_G(\langle u \rangle)}$ . Then  $b_1$  has also defect group  $D$ , and  $l(b_1) = 2$  holds. Moreover,  $u^{2^{r-2}} \in Z(\text{N}_G(\langle u \rangle))$ . Let  $\bar{b}_1 \in \text{Bl}(R[\text{N}_G(u)/\langle u^{2^{r-2}} \rangle])$  be the block which is covered by  $b_1$ . Then  $\bar{b}_1$  has defect group  $D/\langle u^{2^{r-2}} \rangle$ . We argue by induction on  $r$ . Thus, let  $r = 2$ . Then  $b = b_1$  and  $D/\langle u^{2^{r-2}} \rangle = D/\langle u \rangle \cong D_8$ . By Proposition (5G) in [8] the Cartan matrix of  $\bar{b}$  has the elementary divisors 1 and 8. Hence,  $2 = 2^{r-1}$  and  $16 = |D|$  are the elementary divisors of the Cartan matrix of  $b$ . Hence, the claim follows from Theorem 7.2 in [35].

Now assume that the claim already holds for  $r - 1 \geq 2$ . By induction the elementary divisors of the Cartan matrix of  $\bar{b}_1$  are  $2^{r-2}$  and  $|D|/2$ . The claim follows easily as before.  $\square$

Now we are in a position to calculate the Cartan matrix  $C$  up to equivalence of quadratic forms. Here we call two matrices  $M_1, M_2 \in \mathbb{Z}^{l \times l}$  equivalent if there exists a matrix  $S \in \text{GL}(l, \mathbb{Z})$  such that  $A = SBS^T$ , where  $S^T$  denotes the transpose of  $S$ .

By Lemma 3.7 all entries of  $C$  are divisible by  $2^{r-1}$ . Thus, we can consider  $\tilde{C} := 2^{1-r}C \in \mathbb{Z}^{2 \times 2}$ . Then  $\det \tilde{C} = 8$  and the elementary divisors of  $\tilde{C}$  are 1 and 8. If we write

$$\tilde{C} = \begin{pmatrix} c_1 & c_2 \\ c_2 & c_3 \end{pmatrix},$$

then  $\tilde{C}$  corresponds to the positive definite binary quadratic form  $q(x_1, x_2) := c_1x_1^2 + 2c_2x_1x_2 + c_3x_2^2$ . Obviously  $\gcd(c_1, c_2, c_3) = 1$ . If one reduces the entries of  $\tilde{C}$  modulo 2, then one gets a matrix of rank 1 (this is just the multiplicity of the elementary divisor 1). This shows that  $c_1$  or  $c_3$  must be odd. Hence,  $\gcd(c_1, 2c_2, c_3) = 1$ , i. e.  $q$  is primitive (see [10] for example). Moreover,  $\Delta := -4 \det \tilde{C} = -32$  is the discriminant of  $q$ . Now it is easy to see that  $q$  (and  $\tilde{C}$ ) is equivalent to exactly one of the following matrices (see page 20 in [10]):

$$\begin{pmatrix} 1 & 0 \\ 0 & 8 \end{pmatrix} \text{ or } \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}.$$

The Cartan matrices for the block  $\bar{b}_c$  with defect group  $D_8$  (used before) satisfy

$$\begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 8 & 4 \\ 4 & 3 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}^T = \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 4 & 2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix}^T = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}.$$

Hence, only the second matrix occurs up to equivalence. We show that this holds also for the block  $B$ .

**Theorem 3.8.** *The Cartan matrix of  $B$  is equivalent to*

$$2^{r-1} \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}.$$

*Proof.* We argue by induction on  $r$ . The smallest case was already considered by  $b_c$  (this would correspond to  $r = 1$ ). Thus, we may assume  $r \geq 2$  (as usual). First, we determine the generalized decomposition numbers  $d_{\chi\varphi}^u$  for  $u \in \langle c \rangle \setminus \{1\}$  with  $|\langle u \rangle| = 2^k < 2^{r-1}$ . As in the proof of Theorem 3.6, the group  $D/\langle u \rangle$  has the same isomorphism type as  $D$ , but one has to replace  $r$  by  $r - k$ . Hence, by induction we may assume that  $b_u$  has a Cartan matrix which is equivalent to the matrix given in the statement of the theorem. Let  $C_u$  be the Cartan matrix of  $b_u$ , and let  $S_u \in \text{GL}(2, \mathbb{Z})$  such that

$$C_u = 2^{r-1} S_u^T \begin{pmatrix} 4 & 2 \\ 2 & 3 \end{pmatrix} S_u,$$

i. e. with the notations of the previous section, we assume that the “second case” occurs. (This is allowed, since we can only compute the generalized decomposition numbers up to multiplication with  $S_u$  anyway.) As before we write  $\text{IBr}(b_u) = \{\varphi_1, \varphi_2\}$ ,  $D_u := (d_{\chi\varphi_i}^u)$  and  $(\tilde{d}_{\chi\varphi_i}^u) := D_u S_u^{-1}$ . The consideration in the previous section carries over, and one gets

$$\tilde{d}_{\chi\varphi_1}^u = \begin{cases} \pm \zeta_{2^k}^{\lfloor \frac{i-1}{2^{r+2-k}} \rfloor} & \text{if } 1 \leq i \leq k_0(B) \\ 0 & \text{if } k_0(B) < i \leq k(B) \end{cases}$$

and

$$\tilde{d}_{\chi\varphi_2}^u = \begin{cases} \pm \zeta_{2^k}^{\lfloor \frac{i-1}{2^{r-k+1}} \rfloor} & \text{if } 1 \leq i \leq 2^r \\ 0 & \text{if } 2^r < i \leq k_0(B) \\ \pm \zeta_{2^k}^{\lfloor \frac{i-k_0(B)-1}{2^{r-k}} \rfloor} & \text{if } k_0(B) < i \leq k(B) \end{cases},$$

where  $\chi_1, \dots, \chi_{k_0(B)}$  are the characters of height 0. But notice that the ordering of those characters for  $\varphi_1$  and  $\varphi_2$  is different.

Now assume that there is a matrix  $S \in \text{GL}(2, \mathbb{Z})$  such that

$$C = 2^{r-1} S^T \begin{pmatrix} 1 & 0 \\ 0 & 8 \end{pmatrix} S.$$

If  $Q$  denotes the decomposition matrix of  $B$ , we set  $(\tilde{d}_{\chi\varphi_i}^u) := QS^{-1}$  for  $\text{IBr}(B) = \{\varphi_1, \varphi_2\}$ . Then we have

$$|D|m_{\chi\psi}^{(1,B)} = 8\tilde{d}_{\chi\varphi_1} \tilde{d}_{\psi\varphi_1} + \tilde{d}_{\chi\varphi_2} \tilde{d}_{\psi\varphi_2} \text{ for } \chi, \psi \in \text{Irr}(B).$$

In particular  $|D|m_{\chi\chi}^{(1,B)} \equiv 1 \pmod{4}$  for a character  $\chi \in \text{Irr}(B)$  of height 0. For  $u \in \mathcal{T} \setminus Z(D)$  we have  $|D|m_{\chi\chi}^{(u,b_u)} = 2$ , and for  $u \in Z(D) \setminus \langle c \rangle$  we have  $|D|m_{\chi\chi}^{(u,b_u)} = 1$ . Let  $u \in \langle c \rangle \setminus \{1\}$ . Equation (4) and the considerations above imply  $|D|m_{\chi\chi}^{(u,b_u)} \equiv 3 \pmod{4}$ . Now (5B) in [6] reveals the contradiction

$$|D| = \sum_{u \in \mathcal{T}} |D|m_{\chi\chi}^{(u,b_u)} \equiv |D|m_{\chi\chi}^{(1,B)} + 2^{r+1} + 2^{r-1} + 3 \cdot (2^{r-1} - 1) \equiv 2 \pmod{4}. \quad \square$$

With the proof of the last theorem we can also obtain the ordinary decomposition numbers (up to multiplication with an invertible matrix):

$$d_{\chi\varphi_1} = \begin{cases} \pm 1 & \text{if } h(\chi) = 0 \\ 0 & \text{if } h(\chi) = 1 \end{cases}, \quad d_{\chi\varphi_2} = \begin{cases} \pm 1 & \text{if } 0 \leq i \leq 2^r \\ 0 & \text{if } 2^r < i \leq k_0(B) \\ \pm 1 & \text{if } k_0(B) < i \leq k(B) \end{cases}.$$

Again  $\chi_1, \dots, \chi_{k_0(B)}$  are the characters of height 0.

Since we know how  $\mathcal{G}$  acts on the  $B$ -subsections, we can investigate the action of  $\mathcal{G}$  on  $\text{Irr}(B)$ .

**Theorem 3.9.** *The irreducible characters of height 0 of  $B$  split in  $2(r+1)$  families of 2-conjugate characters. These families have sizes  $1, 1, 1, 1, 2, 2, 4, 4, \dots, 2^{r-1}, 2^{r-1}$  respectively. The characters of height 1 split in  $r$  families with sizes  $1, 1, 2, 4, \dots, 2^{r-2}$  respectively. In particular there are exactly six 2-rational characters in  $\text{Irr}(B)$ .*

*Proof.* We start by determining the number of orbits of the action of  $\mathcal{G}$  on the columns of the generalized decomposition matrix. The columns  $\{d_{\chi\varphi_u}^u : \chi \in \text{Irr}(B)\}$  with  $u \in \mathcal{T} \setminus Z(D)$  split in two orbits of length  $2^{r-1}$ . For  $i = 1, 2$  the columns  $\{d_{\chi\varphi_i}^u : \chi \in \text{Irr}(B)\}$  with  $u \in \langle c \rangle$  split in  $r$  orbits of lengths  $1, 1, 2, 4, \dots, 2^{r-2}$  respectively. Finally, the columns  $\{d_{\chi\varphi_u}^u : \chi \in \text{Irr}(B)\}$  with  $u \in Z(D) \setminus \langle c \rangle$  consist of  $r$  orbits of lengths  $1, 1, 2, 4, \dots, 2^{r-2}$  respectively. This gives  $3r+2$  orbits altogether. By Theorem 11 in [3] there also exist exactly  $3r+2$  families of 2-conjugate characters. (Since  $\mathcal{G}$  is noncyclic, one cannot conclude a priori that also the lengths of the orbits of these two actions coincide.)

By considering the column  $\{d_{\chi\varphi_x}^x : \chi \in \text{Irr}(B)\}$ , we see that the irreducible characters of height 0 split in at most  $2(r+1)$  orbits of lengths  $1, 1, 1, 1, 2, 2, 4, 4, \dots, 2^{r-1}, 2^{r-1}$  respectively. Similarly the column  $\{d_{\chi\varphi_2}^c : \chi \in \text{Irr}(B)\}$  shows that there are at most  $r$  orbits of lengths  $1, 1, 2, 4, \dots, 2^{r-2}$  of characters of height 1. Since  $2(r+1) + r = 3r+2$ , these orbits do not merge further, and the claim is proved.  $\square$

Let  $M = \langle x^2, y, z \rangle$  as in Lemma 3.3. Then  $D \subseteq \mathbb{T}_{N_G(M)}(b_M)$ . Since  $e(B) = 1$ , Alperin's fusion theorem implies that  $\mathbb{T}_{N_G(M)}(b_M)$  controls the fusion of  $B$ -subpairs. By Lemma 3.3 we also have  $\mathbb{T}_{N_G(M)}(b_M) \subseteq C_G(c)$  for a  $c \in Z(D)$ . This shows that  $B$  is a so called "centrally controlled block" (see [22]). In [22] it was shown that then the centers of the blocks  $B$  and  $b_c$  (regarded as blocks of  $FG$ ) are isomorphic.

### 3.5 Dade's conjecture

In this section we will verify Dade's (ordinary) conjecture for the block  $B$  (see [12]). First, we need a lemma.

**Lemma 3.10.** *Let  $\tilde{B}$  be a block of  $RG$  with defect group  $\tilde{D} \cong C_{2^s} \times C_2^2$  ( $s \in \mathbb{N}_0$ ) and inertial index 3. Then  $k(\tilde{B}) = k_0(\tilde{B}) = |\tilde{D}| = 2^{s+2}$  and  $l(\tilde{B}) = 3$  hold.*

*Proof.* Let  $\alpha$  be an automorphism of  $\tilde{D}$  of order 3 which is induced by the inertial group. By Lemma 3.2 we have  $C_{\tilde{D}}(\alpha) \cong C_{2^s}$ . We choose a system of representatives  $x_1, \dots, x_k$  for the orbits of  $\tilde{D} \setminus C_{\tilde{D}}(\alpha)$  under  $\alpha$ . Then  $k = 2^s$ . If  $b_i \in \text{Bl}(RC_G(x_i))$  for  $i = 1, \dots, k$  and  $b_u \in \text{Bl}(RC_G(u))$  for  $u \in C_{\tilde{D}}(\alpha)$  are Brauer correspondents of  $\tilde{B}$ , then

$$\bigcup_{i=1}^k \{(x_i, b_i)\} \cup \bigcup_{u \in C_{\tilde{D}}(\alpha)} \{(u, b_u)\}$$

is a system of representatives for the conjugacy classes of  $\tilde{B}$ -subsections. Since  $\alpha \notin C_G(x_i)$ , we have  $l(b_i) = 1$  for  $i = 1, \dots, k$ . In particular  $k(\tilde{B}) \leq 2^{s+2}$  holds. Now we show the opposite inequality by induction on  $s$ .

For  $s = 0$  the claim is well known. Let  $s \geq 1$ . By induction  $l(b_u) = 3$  for  $u \in C_{\tilde{D}}(\alpha) \setminus \{1\}$ . This shows  $k(\tilde{B}) - l(\tilde{B}) = k + (2^s - 1)3 = 2^{s+2} - 3$  and  $l(\tilde{B}) \leq 3$ . An inspection of the numbers  $d_{x\varphi}^{x_1}$  implies  $k(\tilde{B}) = k_0(\tilde{B}) = 2^{s+2} = |\tilde{D}|$  and  $l(\tilde{B}) = 3$ . (This would also follow from Theorem 1 in [46].)  $\square$

Now assume  $O_2(G) = 1$  (this is a hypothesis of Dade's conjecture). In order to prove Dade's conjecture it suffices to consider chains

$$\sigma : P_1 < P_2 < \dots < P_n$$

of nontrivial elementary abelian 2-subgroups of  $G$  (see [12]). (Note that also the empty chain is allowed.) In particular  $P_i \trianglelefteq P_n$  and  $P_n \trianglelefteq N_G(\sigma)$  for  $i = 1, \dots, n$ . Hence, for a block  $b \in \text{Bl}(RN_G(\sigma))$  with  $b^G = B$  and defect group  $Q$  we have  $P_n \leq Q$ . Moreover, there exists a  $g \in G$  such that  ${}^gQ \leq D$ . Thus, by conjugation with  $g$  we may assume  $P_n \leq Q \leq D$  (see also Lemma 6.9 in [12]). This shows  $n \leq 3$ .

In the case  $|P_n| = 8$  we have  $P_n = \langle x^{2^{r-1}}, y, z \rangle =: E$ , because this is the only elementary abelian subgroup of order 8 in  $D$ . Let  $b \in \text{Bl}(RN_G(\sigma))$  with  $b^G = B$ . We choose a defect group  $Q$  of  $\tilde{B} := b^{N_G(E)}$ . Since  $\Omega(Q) = P_n$ , we get  $N_G(Q) \leq N_G(E)$ . Then Brauer's first main theorem implies  $Q = D$ . Hence,  $\tilde{B}$  is the unique Brauer correspondent of  $B$  in  $RN_G(E)$ . For  $M := \langle x^2, y, z \rangle \leq D$  we also have  $N_G(M) \leq N_G(\Omega(M)) = N_G(E)$ . Hence,  $\tilde{B}$  is nonnilpotent. Now consider the chain

$$\tilde{\sigma} : \begin{cases} \emptyset & \text{if } n = 1 \\ P_1 & \text{if } n = 2 \\ P_1 < P_2 & \text{if } n = 3 \end{cases}$$

for the group  $\tilde{G} := N_G(E)$ . Then  $N_G(\sigma) = N_{\tilde{G}}(\tilde{\sigma})$  and

$$\sum_{\substack{b \in \text{Bl}(RN_G(\sigma)), \\ b^G = B}} k^i(b) = \sum_{\substack{b \in \text{Bl}(RN_{\tilde{G}}(\tilde{\sigma})), \\ b^{\tilde{G}} = \tilde{B}}} k^i(b).$$

The chains  $\sigma$  and  $\tilde{\sigma}$  account for all possible chains of  $G$ . Moreover, the lengths of  $\sigma$  and  $\tilde{\sigma}$  have opposite parity. Thus, it seems plausible that the contributions of  $\sigma$  and  $\tilde{\sigma}$  in the alternating sum cancel out each other (this would imply Dade's conjecture). The question which remains is: Can we replace  $(\tilde{G}, \tilde{B}, \tilde{\sigma})$  by  $(G, B, \tilde{\sigma})$ ? We make this more precise in the following lemma.

**Lemma 3.11.** *Let  $\mathcal{Q}$  be a system of representatives for the  $G$ -conjugacy classes of pairs  $(\sigma, b)$ , where  $\sigma$  is a chain (of  $G$ ) of length  $n$  with  $P_n < E$  and  $b \in \text{Bl}(RN_G(\sigma))$  is a Brauer correspondent of  $B$ . Similarly, let  $\tilde{\mathcal{Q}}$  be a system of representatives for the  $\tilde{G}$ -conjugacy classes of pairs  $(\tilde{\sigma}, \tilde{b})$ , where  $\tilde{\sigma}$  is a chain (of  $\tilde{G}$ ) of length  $n$  with  $P_n < E$  and  $\tilde{b} \in \text{Bl}(RN_{\tilde{G}}(\tilde{\sigma}))$  is a Brauer correspondent of  $\tilde{B}$ . Then there exists a bijection between  $\mathcal{Q}$  and  $\tilde{\mathcal{Q}}$  which preserves the numbers  $k^i(b)$ .*

*Proof.* Let  $b_D \in \text{Bl}(RN_G(D))$  be a Brauer correspondent of  $B$ . We consider chains of  $B$ -subpairs

$$\sigma : (P_1, b_1) < (P_2, b_2) < \dots < (P_n, b_n) < (D, b_D),$$

where the  $P_i$  are nontrivial elementary abelian 2-subgroups such that  $P_n < E$ . Then  $\sigma$  is uniquely determined by these subgroups  $P_1, \dots, P_n$  (see Theorem 1.7 in [36]). Moreover, the empty chain is also allowed. Let  $\mathcal{U}$  be a system of representatives for  $G$ -conjugacy classes of such chains. For every chain  $\sigma \in \mathcal{U}$  we define

$$\tilde{\sigma} : (P_1, \tilde{b}_1) < (P_2, \tilde{b}_2) < \dots < (P_n, \tilde{b}_n) < (D, b_D)$$

with  $\tilde{b}_i \in \text{Bl}(RC_{\tilde{G}}(P_i))$  for  $i = 1, \dots, n$ . Finally we set  $\tilde{\mathcal{U}} := \{\tilde{\sigma} : \sigma \in \mathcal{U}\}$ . By Alperin's fusion theorem  $\tilde{\mathcal{U}}$  is a system of representatives for the  $\tilde{G}$ -conjugacy classes of corresponding chains for the group  $\tilde{B}$ . Hence, it suffices to show the existence of bijections  $f$  (resp.  $\tilde{f}$ ) between  $\mathcal{U}$  (resp.  $\tilde{\mathcal{U}}$ ) and  $\mathcal{Q}$  (resp.  $\tilde{\mathcal{Q}}$ ) such that the following property is satisfied: If  $f(\sigma) = (\tau, b)$  and  $\tilde{f}(\tilde{\sigma}) = (\tilde{\tau}, \tilde{b})$ , then  $k^i(b) = k^i(\tilde{b})$  for all  $i \in \mathbb{N}_0$ .

Let  $\sigma \in \mathcal{U}$ . Then we define the chain  $\tau$  by only considering the subgroups of  $\sigma$ , i. e.  $\tau : P_1 < \dots < P_n$ . This gives  $C_G(P_n) \subseteq N_G(\tau)$ , and we can define

$$f : \mathcal{U} \rightarrow \mathcal{Q}, \sigma \mapsto (\tau, b_n^{N_G(\tau)}).$$

Now let  $(\sigma, b) \in \mathcal{Q}$  arbitrary. We write  $\sigma : P_1 < \dots < P_n$ . By Theorem 5.5.15 in [29] there exists a Brauer correspondent  $\beta_n \in \text{Bl}(RC_G(P_n))$  of  $b$ . Since  $(P_n, \beta_n)$  is a  $B$ -subpair, we may assume  $(P_n, \beta_n) < (D, b_D)$  after a suitable conjugation. Then there are uniquely determined blocks  $\beta_i \in \text{Bl}(RC_G(P_i))$  for  $i = 1, \dots, n-1$  such that

$$(P_1, \beta_1) < (P_2, \beta_2) < \dots < (P_n, \beta_n) < (D, b_D).$$

This shows that  $f$  is surjective.

Now let  $\sigma_1, \sigma_2 \in \mathcal{U}$  be given. We write

$$\sigma_i : (P_1^i, \beta_1^i) < \dots < (P_n^i, \beta_n^i)$$

for  $i = 1, 2$ . Let us assume that  $f(\sigma_1) = (\tau_1, b_1)$  and  $f(\sigma_2) = (\tau_2, b_2)$  are conjugate in  $G$ , i. e. there is a  $g \in G$  such that

$$(\tau_2, ({}^g\beta_n^1)^{N_G(\tau_2)}) = {}^g(\tau_1, b_1) = (\tau_2, b_2) = (\tau_2, (\beta_n^2)^{N_G(\tau_2)}).$$

Since  ${}^g\beta_n^1 \in \text{Bl}(RC_G(P_n^2))$  and  $\beta_n^2$  are covered by  $b_2$ , there is  $h \in N_G(\tau_2)$  with  ${}^{hg}\beta_n^1 = \beta_n^2$ . Then

$${}^{hg}(P_n^1, \beta_n^1) = (P_n^2, \beta_n^2).$$

Since the blocks  $\beta_j^i$  for  $i = 1, 2$  and  $j = 1, \dots, n-1$  are uniquely determined by  $P_j^i$ , we also have  ${}^{gh}\sigma_1 = \sigma_2 = \sigma_1$ . This proves the injectivity of  $f$ . Analogously, we define the map  $\tilde{f}$ .

It remains to show that  $f$  and  $\tilde{f}$  satisfy the property given above. For this let  $\sigma \in \mathcal{U}$  with  $\sigma : (P_1, b_1) < \dots < (P_n, b_n)$ ,  $\tilde{\sigma} : (P_1, \tilde{b}_1) < \dots < (P_n, \tilde{b}_n)$ ,  $f(\sigma) = (\tau, b_n^{N_G(\tau)})$  and  $\tilde{f}(\tilde{\sigma}) = (\tilde{\tau}, \tilde{b}_n^{N_{\tilde{G}}(\tilde{\tau})})$ . We have to prove  $k^i(b_n^{N_G(\tau)}) = k^i(\tilde{b}_n^{N_{\tilde{G}}(\tilde{\tau})})$  for  $i \in \mathbb{N}_0$ .

Let  $Q$  be a defect group of  $b_n^{N_G(\tau)}$ . Then  $QC_G(Q) \subseteq N_G(\tau)$ , and there is a Brauer correspondent  $\beta_n \in \text{Bl}(RC_G(Q))$  of  $b_n^{N_G(\tau)}$ . In particular  $(Q, \beta_n)$  is a  $B$ -Brauer subpair. As in Lemma 3.1 we may assume  $Q \in \{D, M, \langle x, z \rangle, \langle xy, z \rangle\}$ . The same considerations also work for the defect group  $\tilde{Q}$  of  $\tilde{b}_n^{N_{\tilde{G}}(\tilde{\tau})}$ . Since  $b_n^{D C_G(P_n)} = \tilde{b}_n^{D C_G(P_n)}$ , we get:

$$Q = D \iff D \subseteq N_G(\tau) \iff D \subseteq N_{\tilde{G}}(\tilde{\tau}) \iff \tilde{Q} = D.$$

Let us consider the case  $Q = D (= \tilde{Q})$ . Let  $b_M \in \text{Bl}(RC_G(M))$  such that  $(M, b_M) \leq (D, b_D)$  and  $\alpha \in \text{T}_{N_G(M)}(b_M) \setminus DC_G(M) \subseteq N_G(M) \subseteq \tilde{G}$ . Then:

$$b_n^{N_G(\tau)} \text{ is nilpotent} \iff \alpha \notin N_G(\tau) \iff \alpha \notin N_{\tilde{G}}(\tau) \iff \tilde{b}_n^{N_{\tilde{G}}(\tau)} \text{ is nilpotent.}$$

Thus, the claim holds in this case. Now let  $Q < D$  (and  $\tilde{Q} < D$ ). Then we have  $QC_G(Q) = C_G(Q) \subseteq C_G(P_n)$ . Since  $\beta_n^{C_G(P_n)}$  is also a Brauer correspondent of  $b_n^{N_G(\tau)}$ , the blocks  $\beta_n^{C_G(P_n)}$  and  $b_n$  are conjugate. In particular  $b_n$  (and  $\tilde{b}_n$ ) has defect group  $Q$ . Hence, we obtain  $Q = \tilde{Q}$ . If  $Q \in \{\langle x, z \rangle, \langle xy, z \rangle\}$ , then  $b_n^{N_G(\tau)}$  and  $\tilde{b}_n^{N_{\tilde{G}}(\tau)}$  are nilpotent, and the claim holds. Thus, we may assume  $Q = M$ . Then as before:

$$b_n^{N_G(\tau)} \text{ is nilpotent} \iff \alpha \notin N_G(\tau) \iff \alpha \notin N_{\tilde{G}}(\tau) \iff \tilde{b}_n^{N_{\tilde{G}}(\tau)} \text{ is nilpotent.}$$

We may assume that the nonnilpotent case occurs. Then  $t(b_n^{N_G(\tau)}) = t(\tilde{b}_n^{N_{\tilde{G}}(\tau)}) = 3$ , and the claim follows from Lemma 3.10.  $\square$

As explained in the beginning of the section, the Dade conjecture follows.

**Theorem 3.12.** *The Dade conjecture holds for  $B$ .*

### 3.6 Alperin's weight conjecture

In this section we prove Alperin's weight conjecture for  $B$ . Let  $(P, \beta)$  be a weight for  $B$ , i.e.  $P$  is a 2-subgroup of  $G$  and  $\beta$  is a block of  $R[N_G(P)/P]$  with defect 0. Moreover,  $\beta$  is dominated by a Brauer correspondent  $b \in \text{Bl}(RN_G(P))$  of  $B$ . As usual, one can assume  $P \leq D$ . If  $\text{Aut}(P)$  is a 2-group, then  $N_G(P)/C_G(P)$  is also a 2-group. Then  $P$  is a defect group of  $b$ , since  $\beta$  has defect 0. Moreover,  $\beta$  is uniquely determined by  $b$ . By Brauer's first main theorem we have  $P = D$ . Thus, in this case there is exactly one weight for  $B$  up to conjugation.

Now let us assume that  $\text{Aut}(P)$  is not a 2-group (in particular  $P < D$ ). As usual,  $\beta$  covers a block  $\beta_1 \in \text{Bl}(R[C_G(P)/P])$ . By the Fong-Reynolds theorem (see [29] for example) also  $\beta_1$  has defect 0. Hence,  $\beta_1$  is dominated by exactly one block  $b_1 \in \text{Bl}(RC_G(P))$  with defect group  $P$ . Since  $\beta\beta_1 \neq 0$ , we also have  $bb_1 \neq 0$ , i.e.  $b$  covers  $b_1$ . Thus, the situation is as follows:

$$\begin{array}{ccc} \beta \in \text{Bl}(R[N_G(P)/P]) & \longleftrightarrow & b \in \text{Bl}(RN_G(P)) \\ \updownarrow & & \updownarrow \\ \beta_1 \in \text{Bl}(R[C_G(P)/P]) & \longleftrightarrow & b_1 \in \text{Bl}(RC_G(P)) \end{array}$$

By Theorem 5.5.15 in [29] we have  $b_1^{N_G(P)} = b$  and  $b_1^G = B$ . This shows that  $(P, b_1)$  is a  $B$ -Brauer subpair. Then  $P = M (= \langle x^2, y, z \rangle)$  follows. By Brauer's first main theorem  $b$  is uniquely determined (independent of  $\beta$ ). Now we prove that also  $\beta$  is uniquely determined by  $b$ .

In order to do so it suffices to show that  $\beta$  is the only block with defect 0 which covers  $\beta_1$ . By the Fong-Reynolds theorem it suffices to show that  $\beta_1$  is covered by only one block of  $R\overline{\text{T}_{N_G(M)/M}}(\beta_1) = R[\overline{\text{T}_{N_G(M)}(b_1)}/M]$  with defect 0. For convenience we write  $\overline{C_G(M)} := C_G(M)/M$ ,  $\overline{N_G(M)} := N_G(M)/M$  and  $\overline{\text{T}} := \overline{\text{T}_{N_G(M)}(b_1)}/M$ . Let  $\chi \in \text{Irr}(\beta_1)$ . The irreducible constituents of  $\text{Ind}_{\overline{C_G(M)}}^{\overline{\text{T}}}(\chi)$  belong to blocks which covers  $\beta_1$  (where  $\text{Ind}$  denote induction). Conversely, every block of  $R\overline{\text{T}}$  which covers  $\beta_1$  arises in this way (see Lemma 5.5.7 in [29]). Let

$$\text{Ind}_{\overline{C_G(M)}}^{\overline{\text{T}}}(\chi) = \sum_{i=1}^t e_i \psi_i$$

with  $\psi_i \in \text{Irr}(\overline{\text{T}})$  and  $e_i \in \mathbb{N}$  for  $i = 1, \dots, t$ . Then

$$\sum_{i=1}^t e_i^2 = |\overline{\text{T}} : \overline{C_G(M)}| = |\overline{\text{T}_{N_G(M)}(b_1)} : C_G(M)| = 6$$

(see page 84 in [17]). Thus, there is some  $i \in \{1, \dots, t\}$  with  $e_i = 1$ , i. e.  $\chi$  is extendible to  $\overline{\mathbb{T}}$ . We may assume  $e_1 = 1$ . By Corollary 6.17 in [17] it follows that  $t = |\text{Irr}(\overline{\mathbb{T}/C_G(M)})| = |\text{Irr}(S_3)| = 3$  and

$$\{\psi_1, \psi_2, \psi_3\} = \{\psi_1\tau : \tau \in \text{Irr}(\overline{\mathbb{T}/C_G(M)})\},$$

where the characters in  $\text{Irr}(\overline{\mathbb{T}/C_G(M)})$  were identified with their inflations in  $\text{Irr}(\overline{\mathbb{T}})$ . Thus, we may assume  $e_2 = 1$  and  $e_3 = 2$ . Then it is easy to see that  $\psi_1$  and  $\psi_2$  belong to blocks with defect at least 1. Hence, only the block with contains  $\psi_3$  is allowed. This shows uniqueness.

Finally we show that there is in fact a weight of the form  $(M, \beta)$ . For this we choose  $b, b_1, \beta_1, \chi$  and  $\psi_i$  as above. Then  $\chi$  vanishes on all nontrivial 2-elements. Moreover,  $\psi_1$  is an extension of  $\chi$ . Let  $\tau \in \text{Irr}(\overline{\mathbb{T}/C_G(M)})$  be the character of degree 2. Then  $\tau$  vanishes on all nontrivial 2-elements of  $\overline{\mathbb{T}/C_G(M)}$ . Hence,  $\psi_3 = \psi_1\tau$  vanishes on all nontrivial 2-elements of  $\overline{\mathbb{T}}$ . This shows that  $\psi_3$  belongs in fact to a block  $\tilde{\beta} \in \text{Bl}(R\overline{\mathbb{T}})$  with defect 0. Then  $(M, \tilde{\beta}^{\overline{N_G(M)}})$  is the desired weight for  $B$ .

Hence, we have shown that there are exactly two weights for  $B$  up to conjugation. Since  $l(B) = 2$ , Alperin's weight conjecture is satisfied.

**Theorem 3.13.** *Alperin's weight conjecture holds for  $B$ .*

### 3.7 The gluing problem

Finally we show that the gluing problem (see Conjecture 4.2 in [26]) for the block  $B$  has a unique solution. We will not recall the very technical statement of the gluing problem. Instead we refer to [37] for most of the notations. Observe that the field  $F$  is denoted by  $k$  in [37].

**Theorem 3.14.** *The gluing problem for  $B$  has a unique solution.*

*Proof.* As in [37] we denote the fusion system induced by  $B$  with  $\mathcal{F}$ . Then the  $\mathcal{F}$ -centric subgroups of  $D$  are given by  $M_1 := \langle x^2, y, z \rangle$ ,  $M_2 := \langle x, z \rangle$ ,  $M_3 := \langle xy, z \rangle$  and  $D$ . We have seen so far that  $\text{Aut}_{\mathcal{F}}(M_1) \cong \text{Out}_{\mathcal{F}}(M_1) \cong S_3$ ,  $\text{Aut}_{\mathcal{F}}(M_i) \cong D/M_i \cong C_2$  for  $i = 2, 3$  and  $\text{Aut}_{\mathcal{F}}(D) \cong D/Z(D) \cong C_2^2$  (see proof of Lemma 3.3). Using this, we get  $H^i(\text{Aut}_{\mathcal{F}}(\sigma), F^\times) = 0$  for  $i = 1, 2$  and every chain  $\sigma$  of  $\mathcal{F}$ -centric subgroups (see proof of Corollary 2.2 in [37]). Hence,  $H^0([S(\mathcal{F}^c)], \mathcal{A}_{\mathcal{F}}^2) = H^1([S(\mathcal{F}^c)], \mathcal{A}_{\mathcal{F}}^1) = 0$ . Now the claim follows from Theorem 1.1 in [37].  $\square$

## 4 The case $r = s > 1$

In the section we assume that  $B$  is a nonnilpotent block of  $RG$  with defect group

$$D := \langle x, y \mid x^{2^r} = y^{2^r} = [x, y]^2 = [x, x, y] = [y, x, y] = 1 \rangle$$

for  $r \geq 2$ . As before we define  $z := [x, y]$ . Since  $|D/\Phi(D)| = 4$ , 2 and 3 are the only prime divisors of  $|\text{Aut}(D)|$ . In particular  $t(B) \in \{1, 3\}$ . If  $t(B) = 1$ , then  $B$  would be nilpotent by Theorem 2.4. Thus, we have  $t(B) = 3$ .

### 4.1 The $B$ -subsections

We investigate the automorphism group of  $D$ .

**Lemma 4.1.** *Let  $\alpha \in \text{Aut}(D)$  be an automorphism of order 3. Then  $z$  is the only nontrivial fixed-point of  $Z(D)$  under  $\alpha$ .*

*Proof.* Since  $D' = \langle z \rangle$ ,  $z$  remains fixed under all automorphisms of  $D$ . Moreover,  $\alpha(x) \in yZ(D) \cup xyZ(D)$ , because  $\alpha$  acts nontrivially on  $D/Z(D)$ . In both cases we have  $\alpha(x^2) \neq x^2$ . This shows that  $\alpha|_{Z(D)} \in \text{Aut}(Z(D))$  is also an automorphism of order 3. Obviously  $\alpha$  induces an automorphism of order 3 on  $Z(D)/\langle z \rangle \cong C_{2^{r-1}}^2$ . But this automorphism is fixed-point-free (see Lemma 1 in [27]). The claim follows.  $\square$

Using this, we can find a system of representatives for the conjugacy classes of  $B$ -subsections.

**Lemma 4.2.** *Let  $b \in \text{Bl}(RDC_G(D))$  be a Brauer correspondent of  $B$ , and for  $Q \leq D$  let  $b_Q$  be the unique block of  $RQC_G(Q)$  with  $(Q, b_Q) \leq (D, b)$ . We choose a system  $\mathcal{S} \subseteq Z(D)$  of representatives for the orbits of  $Z(D)$  under the action of  $\text{TN}_{G(D)}(b)$ . We set  $\mathcal{T} := \mathcal{S} \cup \{y^i x^{2j} : i, j \in \mathbb{Z}, i \text{ odd}\}$ . Then*

$$\bigcup_{a \in \mathcal{T}} \{(a, b_{C_D(a)}^{C_G(a)})\}$$

is a system of representatives for the conjugacy classes of  $B$ -subsections. Moreover,

$$|\mathcal{T}| = \frac{5 \cdot 2^{2(r-1)} + 4}{3}.$$

*Proof.* Proposition 2.12.(ii) in [34] states the desired system wrongly. More precisely the claim  $I_D = Z(D)$  in the proof is false. Indeed Lemma 4.1 shows  $I_D = \mathcal{S}$ . Now the claim follows easily.  $\square$

From now on we write  $b_a := b_{C_D(a)}^{C_G(a)}$  for  $a \in \mathcal{T}$ . We are able to determine the difference  $k(B) - l(B)$ .

**Proposition 4.3.** *We have*

$$k(B) - l(B) = \frac{5 \cdot 2^{2(r-1)} + 7}{3}.$$

*Proof.* Consider  $l(b_a)$  for  $1 \neq a \in \mathcal{T}$ .

**Case 1:**  $a \in Z(D)$ .

Then  $b_a$  is a block with defect group  $D$ . Moreover,  $b_a$  and  $B$  have a common Brauer correspondent in  $\text{Bl}(RDC_{C_G(a)}(D)) = \text{Bl}(RDC_G(D))$ . In case  $a \neq z$  we have  $t(b_a) = 1$  by Lemma 4.1. Hence,  $b_a$  is nilpotent and  $l(b_a) = 1$ . Now let  $a = z$ . Then there exists a block  $\overline{b_z}$  of  $C_G(z)/\langle z \rangle$  with defect group  $D/\langle z \rangle \cong C_{2^r}^2$  and  $l(\overline{b_z}) = l(b_z)$ . By Theorem 1.5(iv) in [33],  $t(\overline{b_z}) = t(b_z) = 3$  holds. Thus, Theorem 2 in [43] implies  $l(b_z) = l(\overline{b_z}) = 3$ .

**Case 2:**  $a \notin Z(D)$ .

Then  $b_{C_P(a)} = b_M$  is a block with defect group  $M := \langle x^2, y, z \rangle$ . Since  $b_M^{DC_G(M)} = b_D^{DC_G(M)}$ , also  $b_M^{C_G(a)} = b_a$  has defect group  $M$ . For every automorphism  $\alpha \in \text{Aut}(D)$  of order 3 we have  $\alpha(M) \neq M$ . Since  $D$  controls the fusion of  $B$ -subpairs, we get  $t(b_a) = l(b_a) = 1$ .

Now the conclusion follows from  $k(B) = \sum_{a \in \mathcal{T}} l(b_a)$ .  $\square$

The next result concerns the Cartan matrix of  $B$ .

**Lemma 4.4.** *The elementary divisors of the Cartan matrix of  $B$  are contained in  $\{1, 2, |D|\}$ . The elementary divisor 2 occurs twice and  $|D|$  occurs once (as usual). In particular  $l(B) \geq 3$ .*

*Proof.* Let  $C$  be the Cartan matrix of  $B$ . As in Lemma 3.7 we use the notion of lower defect groups. For this let  $P < D$  such that  $|P| \geq 4$ , and let  $b \in \text{Bl}(RN_G(P))$  be a Brauer correspondent of  $B$  with defect group  $Q \leq D$ . Brauer's first main theorem implies  $P < Q$ . By Proposition 1.3 in [33] there exists a block  $\beta \in \text{Bl}(RC_G(P))$  with  $\beta^{N_G(P)} = b$  such that at most  $l(\beta)$  lower defect groups of  $b$  contain a conjugate of  $P$ . Let  $S \leq Q$  be a defect group of  $\beta$ . First, we consider the case  $S = D$ . Then  $P \subseteq Z(D)$ . By Lemma 4.1 we have  $l(\beta) = 1$ , since  $|P| \geq 4$ . It follows that  $m_b^1(P) = m_b(P) = 0$ , because  $P$  is contained in the (lower) defect group  $Q$  of  $b$ .

Now assume  $S < D$ . In particular  $S$  is abelian. If  $S$  is even metacyclic, then  $l(\beta) = 1$  and  $m_b^1(P) = 0$ , since  $P \subseteq Z(C_G(P))$ . Thus, let us assume that  $S$  is nonmetacyclic. By (3C) in [5],  $x^2 \in Z(D)$  is conjugate to an element of  $Z(S)$ . This shows  $S \cong C_{2^k} \times C_{2^l} \times C_2$  with  $k \in \{r, r-1\}$  and  $1 \leq l \leq r$ . If  $1, k, l$  are pairwise distinct, then  $l(\beta) = 1$  and  $m_b^1(P) = 0$  follow from Lemma 2.3. Let  $k = l$ . Then every automorphism of  $S$  of order 3 has only one nontrivial fixed-point. Since  $|P| \geq 4$ , it follows again that  $l(\beta) = 1$  and  $m_b^1(P) = 0$ .

Now let  $S \cong C_{2^k} \times C_2^2$  with  $2 \leq k \in \{r-1, r\}$ . Assume first that  $P$  is noncyclic. Then  $S/P$  is metacyclic. If  $S/P$  is not a product of two isomorphic cyclic groups, then  $l(\beta) = 1$  and  $m_b^1(P) = 0$ . Hence, we may assume

$S/P \cong C_2^2$ . It is easy to see that there exists a subgroup  $P_1 \leq P$  with  $S/P_1 \cong C_4 \times C_2$ . We get  $l(\beta) = 1$  and  $m_b^1(P) = 0$  also in this case.

Finally, let  $P = \langle u \rangle$  be cyclic. Then  $(u, \beta)$  is a  $B$ -subsection. Since  $|P| \geq 4$ ,  $u$  is not conjugate to  $z$ . As in the proof of Proposition 4.3 we have  $l(\beta) = 1$  and  $m_b^1(P) = 0$ . This shows  $m_B^1(P) = 0$ . Since  $P$  was arbitrary, the multiplicity of  $|P|$  as an elementary divisor of  $C$  is 0.

It remains to consider the case  $|P| = 2$ . We write  $P = \langle u \rangle \leq D$ . As before let  $b \in \text{Bl}(RN_G(P))$  be a Brauer correspondent of  $B$ . Then  $(u, b)$  is a  $B$ -subsection. If  $(u, b)$  is not conjugate to  $(z, b_z)$ , then  $l(b) = 1$  and  $m_b^1(P) = 0$  as in the proof of Proposition 4.3. Since we can replace  $P$  by a conjugate, we may assume  $P = \langle z \rangle$  and  $(u, b) = (z, b_z)$ . Then  $l(b) = 3$  and  $D$  is a defect group of  $b$ . Now let  $\bar{b} \in \text{Bl}(R[N_G(P)/P])$  be the block which is dominated by  $b$ . By Corollary 1 in [16] the elementary divisors of the Cartan matrix of  $\bar{b}$  are 1, 1,  $|D|/2$ . Hence, the elementary divisors of the Cartan matrix of  $b$  are 2, 2,  $|D|$ . This shows

$$2 = \sum_{\substack{Q \in \mathcal{P}(N_G(P)), \\ |Q|=2}} m_b^1(Q),$$

where  $\mathcal{P}(N_G(P))$  is a system of representatives for the conjugacy classes of  $p$ -subgroups of  $N_G(P)$ . The same arguments applied to  $b$  instead of  $B$  imply  $m_b^1(Q) = 0$  for  $P \neq Q \leq N_G(P)$  with  $|Q| = 2$ . Hence,  $2 = m_b^1(P) = m_B^1(P)$ , and 2 occurs as elementary divisors of  $C$  twice.  $\square$

As in Section 3 we write  $\text{IBr}(b_u) = \{\varphi_u\}$  for  $u \in \mathcal{T} \setminus \langle z \rangle$ . In a similar manner we define the integers  $a_i^u$ . If  $u \in \mathcal{T} \setminus \langle z \rangle$  with  $|\langle u \rangle| = 2^k > 2$ , then the  $2^{k-1}$  distinct subsections of the form  ${}^\gamma(u, b_u)$  for  $\gamma \in \mathcal{G}$  are pairwise nonconjugate (same argument as in the case  $r > s = 2$ ). Hence, Lemma 3.4 carries over in a corresponding form. Apart from that we can also carry over Lemma (6.B) in [20]:

**Lemma 4.5.** *Let  $\chi \in \text{Irr}(B)$  and  $u \in \mathcal{T} \setminus Z(D)$ . Then  $\chi$  has height 0 if and only if the sum*

$$\sum_{i=0}^{2^{r-1}-1} a_i^u(\chi)$$

*is odd.*

*Proof.* If  $\chi$  has height 0, the sum is odd by Proposition 1 in [9]. The other implication follows easily from (5G) in [6].  $\square$

The next lemma is the analogon to Lemma 3.5.

**Lemma 4.6.** *Let  $u \in Z(D) \setminus \langle z \rangle$  of order  $2^k$ . Then for all  $\chi \in \text{Irr}(B)$  we have:*

$$(i) \quad 2^{h(\chi)} \mid a_i^u(\chi) \text{ for } i = 0, \dots, 2^{k-1} - 1,$$

$$(ii) \quad \sum_{i=0}^{2^{k-1}-1} a_i^u(\chi) \equiv 2^{h(\chi)} \pmod{2^{h(\chi)+1}}.$$

As in the case  $r > s = 1$ , Lemma 1.1 in [39] implies

$$k(B) \leq \sum_{i=0}^{\infty} 2^{2i} k_i(B) \leq |D|. \quad (6)$$

In particular Brauer's  $k(B)$ -conjecture holds. Moreover, Theorem 3.1 in [39] gives  $k_0(B) \leq |D|/2 = |D : D'|$ , i. e. Olsson's conjecture is satisfied. Using this, we can improve the inequality (6) to

$$|D| \geq k_0(B) + 4(k(B) - k_0(B)) = 4k(B) - 3k_0(B) \geq 4k(B) - \frac{3|D|}{2}$$

and

$$\frac{5 \cdot 2^{2(r-1)} + 16}{3} \leq k(B) - l(B) + l(B) = k(B) \leq \frac{5|D|}{8} = 5 \cdot 2^{2(r-1)}.$$



We will improve this further. Let  $\overline{b_z}$  be the block of  $\text{Bl}(RC_G(z)/\langle z \rangle)$  which is dominated by  $b_z$ . Then  $\overline{b_z}$  has defect group  $D/\langle z \rangle \cong C_{2^r}^2$ . Using the existence of a perfect isometry (see [44, 45, 38]), one can show that the Cartan matrix of  $\overline{b_z}$  is equivalent to

$$\overline{C} := \frac{1}{3} \begin{pmatrix} 2^{2r} + 2 & 2^{2r} - 1 & 2^{2r} - 1 \\ 2^{2r} - 1 & 2^{2r} + 2 & 2^{2r} - 1 \\ 2^{2r} - 1 & 2^{2r} - 1 & 2^{2r} + 2 \end{pmatrix}.$$

Hence, the Cartan matrix of  $b_z$  is equivalent to  $2\overline{C}$ . Now inequality (\*\*) in [24] yields

$$k(B) \leq 2 \frac{2^{2r} + 8}{3} = \frac{|D| + 16}{3}.$$

(Notice that the proof of Theorem A in [24] also works for  $b_z$  instead of  $B$ , since the generalized decomposition numbers corresponding to  $(z, b_z)$  are integral. See also Lemma 3 in [42].)

In addition we have

$$k_i(B) = 0 \text{ for } i \geq 4$$

by Corollary (6D) in [7]. This means that the heights of the characters in  $\text{Irr}(B)$  are bounded independently of  $r$ . We remark also that Alperin's weight conjecture is equivalent to

$$l(B) = l(b)$$

for the Brauer correspondent  $b \in \text{Bl}(RN_G(D))$  of  $B$  (see Consequence 5 in [1]). Since  $z \in Z(N_G(D))$ ,  $l(B) = l(b) = 3$  and  $k(B) = (5 \cdot 2^{2(r-1)} + 16)/3$  would follow in this case (see proof of Proposition 4.3).

## 4.2 The gluing problem

As in section 3.7 we use the notations of [37].

**Theorem 4.7.** *The gluing problem for  $B$  has a unique solution.*

*Proof.* Let  $\mathcal{F}$  be the fusion system induced by  $B$ . Then the  $\mathcal{F}$ -centric subgroups of  $D$  are given by  $M := \langle x^2, y, z \rangle$  and  $D$  (up to conjugation in  $\mathcal{F}$ ). We have  $\text{Aut}_{\mathcal{F}}(M) \cong D/M \cong C_2$  and  $\text{Aut}_{\mathcal{F}}(D) \cong A_4$ . This shows  $H^2(\text{Aut}_{\mathcal{F}}(\sigma), F^\times) = 0$  for every chain  $\sigma$  of  $\mathcal{F}$ -centric subgroups. Consequently,  $H^0([S(\mathcal{F}^c)], \mathcal{A}_{\mathcal{F}}^2) = 0$ . On the other hand, we have  $H^1(\text{Aut}_{\mathcal{F}}(D), F^\times) \cong H^1(C_3, F^\times) \cong C_3$  and  $H^1(\text{Aut}_{\mathcal{F}}(\sigma), F^\times) = 0$  for all chains  $\sigma \neq D$ . Hence, the situation is as in Case 3 of the proof of Theorem 1.2 in [37]. However, the proof in [37] is pretty short. For the convenience of the reader, we give a more complete argument.

Since  $[S(\mathcal{F}^c)]$  is partially ordered by taking subchains, one can view  $[S(\mathcal{F}^c)]$  as a category, where the morphisms are given by the pairs of ordered chains. In particular  $[S(\mathcal{F}^c)]$  has exactly five morphisms. With the notations of [47] the functor  $\mathcal{A}_{\mathcal{F}}^1$  is a *representation* of  $[S(\mathcal{F}^c)]$  over  $\mathbb{Z}$ . Hence, we can view  $\mathcal{A}_{\mathcal{F}}^1$  as a module  $\mathcal{M}$  over the incidence algebra of  $[S(\mathcal{F}^c)]$ . More precisely, we have

$$\mathcal{M} := \bigoplus_{a \in \text{Ob}[S(\mathcal{F}^c)]} \mathcal{A}_{\mathcal{F}}^1(a) = \mathcal{A}_{\mathcal{F}}^1(D) \cong C_3.$$

Now we can determine  $H^1([S(\mathcal{F}^c)], \mathcal{A}_{\mathcal{F}}^1)$  using Lemma 6.2(2) in [47]. For this let  $d : \text{Hom}[S(\mathcal{F}^c)] \rightarrow \mathcal{M}$  a derivation. Then we have  $d(\alpha) = 0$  for all  $\alpha \in \text{Hom}[S(\mathcal{F}^c)]$  with  $\alpha \neq (D, D) =: \alpha_1$ . However,

$$d(\alpha_1) = d(\alpha_1 \alpha_1) = (\mathcal{A}_{\mathcal{F}}^1(\alpha_1))(d(\alpha_1)) + d(\alpha_1) = 2d(\alpha_1) = 0.$$

Hence,  $H^1([S(\mathcal{F}^c)], \mathcal{A}_{\mathcal{F}}^1) = 0$ . □

### 4.3 Special cases

Since the general methods do not suffice to compute the invariants of  $B$ , we restrict ourself to certain special situations.

**Proposition 4.8.** *If  $O_2(G) \neq 1$ , then*

$$k(B) = \frac{5 \cdot 2^{2(r-1)} + 16}{3}, \quad k_0(B) \geq \frac{2^{2r} + 8}{3}, \quad l(B) = 3.$$

*Proof.* Let  $1 \neq Q := O_2(G)$ . Then  $Q \subseteq D$ . In the case  $Q = D'$  we have  $C_G(z) = N_G(Q) = G$  and  $B = b_z$ . Then the assertions on  $k(B)$  and  $l(B)$  are clear. Moreover,  $b_z$  dominates a block  $\overline{b_z} \in \text{Bl}(RC_G(z)/\langle z \rangle)$  with defect group  $C_{2^r}^2$ . By Theorem 2 in [43] we have

$$k_0(B) \geq k_0(\overline{b_z}) = k(\overline{b_z}) = \frac{2^{2r} + 8}{3}.$$

Hence, we may assume  $Q \neq D'$ . With the same argument we may also assume  $Q < D$ . In particular  $Q$  is abelian. We consider a  $B$ -subpair  $(Q, b_Q)$ . Then  $D$  or  $M$  is a defect group of  $b_Q$  (see proof of Lemma 4.2). If  $D$  is a defect group of  $b_Q$ , then  $D \subseteq C_G(Q)$  and  $Q \subseteq Z(D)$ . By Lemma 4.1 it follows that  $b_Q$  is nilpotent.

Now let us assume that  $M$  is a defect group of  $b_Q$ . Since  $D$  controls the fusions of  $B$ -subpairs, we have  $t(b_Q) = 1$  (see Case 2 in the proof of Proposition 4.3). Hence, again  $b_Q$  is nilpotent. Thus, in both cases  $B$  is an extension of a nilpotent block of  $\text{Bl}(RC_G(Q))$ . In this situation the Külshammer-Puig theorem applies. In particular we can replace  $B$  by a block with normal defect group (see [23]). Hence,  $B = b_z$ , and the claim follows as before.  $\square$

Since  $N_G(D) \subseteq C_G(z)$ ,  $B$  is a ‘‘centrally controlled block’’ (see [22]). In [22] it was shown that then an epimorphism  $Z(B) \rightarrow Z(b_z)$  exists, where one has to regard  $B$  (resp.  $b_z$ ) as blocks of  $FG$  (resp.  $FC_G(z)$ ). Moreover, we conjecture that the blocks  $B$  and  $b_z$  are Morita-equivalent. For the similar defect group  $Q_8$  this holds in fact (see [18]). In this context the work [11] is also interesting. There it was shown that there is a perfect isometry between any two blocks with the same quaternion group as defect group and the same fusion of subpairs. Thus, it would be also possible that there is a perfect isometry between  $B$  and  $b_z$ .

**Proposition 4.9.** *In order to determine  $k(B)$  (and thus also  $l(B)$ ), we may assume that  $O_2(G)$  is trivial and  $O_2'(G) = Z(G) = F(G)$  is cyclic. Moreover, we can assume that  $G$  is an extension of a solvable group by a quasisimple group. In particular  $G$  has only one nonabelian composition factor.*

*Proof.* By Proposition 4.8 we may assume  $O_2(G) = 1$ . Now we consider  $O(G) := O_{2'}(G)$ . Using Clifford theory we may assume that  $O(G)$  is central and cyclic (see e.g. Theorem X.1.2 in [15]). Since  $O_2(G) = 1$ , we get  $O(G) = Z(G)$ . Let  $E(G)$  be the normal subgroup of  $G$  generated by the components. As usual,  $B$  covers a block  $b$  of  $E(G)$ . By Fong-Reynolds we can assume that  $b$  is stable in  $G$ . Then  $d := D \cap E(G)$  is a defect group of  $b$ . By the Külshammer-Puig result we may assume that  $b$  is nonnilpotent. In particular  $d$  has rank at least 2. Let  $C_1, \dots, C_n$  be the components of  $G$ . Then  $E(G)$  is the central product of  $C_1, \dots, C_n$ . Since  $[C_i, C_j] = 1$  for  $i \neq j$ ,  $b$  covers exactly one block  $\beta_i$  of  $RC_i$  for  $i = 1, \dots, n$ . Then  $b$  is dominated by the block  $\beta_1 \otimes \dots \otimes \beta_n$  of  $R[C_1 \times \dots \times C_n]$ . Since  $Z(C_1)$  is abelian and subnormal in  $G$ , it must have odd order. Hence, we may identify  $b$  with  $\beta_1 \otimes \dots \otimes \beta_n$  (see Proposition 1.5 in [13]). In particular  $d = \delta_1 \times \dots \times \delta_n$ , where  $\delta_i := d \cap C_i$  is a defect group of  $\beta_i$  for  $i = 1, \dots, n$ . Assume that  $\delta_1$  is cyclic. Then  $\beta_1$  is nilpotent and isomorphic to  $(R\delta_1)^{m \times m}$  for some  $m \in \mathbb{N}$  by Puig. Let  $\{C_1, \dots, C_k\}$  be the orbit of  $C_1$  under the conjugation action of  $G$  ( $k \leq n$ ). Then  $\beta_1 \otimes \dots \otimes \beta_k \cong (R\delta_1)^{m_1 \times m_1}$  (for some  $m_1 \in \mathbb{N}$ ) is a block of  $R[C_1 \dots C_k]$  with  $l(\beta_1 \otimes \dots \otimes \beta_k) = 1$ . Lemma 2.1(v) implies  $k \leq 2$  or  $k = 3$  and  $|\delta_1| = 2$ . In the first case Theorem 2 in [43] shows that  $\beta_1 \otimes \dots \otimes \beta_k$  is nilpotent. This also holds in the second case by [25]. Since  $C_1 \dots C_k \trianglelefteq G$ ,  $B$  is an extension of a nilpotent block. This shows that we can assume that the groups  $\delta_i$  are noncyclic for  $i = 1, \dots, n$ . By Lemma 2.1(v),  $d$  has rank at most 3. Hence,  $n = 1$  and  $E(G) = C_1$ .

That means in order to determine the invariants of the block  $B$  we may assume that  $G$  contains only one component. Let  $F(G)$  (resp.  $F^*(G)$ ) be the Fitting subgroup (resp. generalized Fitting subgroup) of  $G$ . Since  $F(G) = Z(G)$ , we have  $C_G(E(G)) = C_G(F^*(G)) \leq F(G)$ . Hence,  $C_G(E(G))$  is nilpotent. On the other hand, the quotient  $G/C_G(E(G))$  is isomorphic to a subgroup of the automorphism group of the quasisimple group  $E(G)$ .

Consider the canonical map  $f : \text{Aut}(E(G)) \rightarrow \text{Aut}(E(G)/Z(E(G)))$ . Let  $\alpha \in \ker f$ . Then  $\alpha(g)g^{-1} \in Z(E(G))$  for all  $g \in E(G)$ . Hence, we get a map  $\beta : E(G) \rightarrow Z(E(G))$ ,  $g \mapsto \alpha(g)g^{-1}$ . Moreover, it is easy to see that  $\beta$  is a homomorphism. Since  $E(G)$  is perfect, we get  $\beta = 1$  and thus  $\alpha = 1$ . This shows  $\text{Aut}(E(G)) \leq \text{Aut}(E(G)/Z(E(G)))$ . By Schreier's conjecture (which can be proven using the classification)  $\text{Aut}(E(G)/Z(E(G)))$  is an extension of the solvable group  $\text{Out}(E(G)/Z(E(G)))$  by the simple group  $\text{Inn}(E(G)/Z(E(G))) \cong E(G)/Z(E(G))$ . Taking these facts together, we see that  $G$  has only one nonabelian composition factor. In particular  $G$  is an extension of a solvable group by a quasisimple group.  $\square$

Now we consider blocks of maximal defect, i. e.  $D$  is a Sylow 2-subgroup of  $G$ . These include principal blocks.

**Proposition 4.10.** *If  $B$  has maximal defect, then  $G$  is solvable. In particular Alperin's weight conjecture is satisfied, and we have*

$$\begin{aligned} k(B) &= \frac{5 \cdot 2^{2(r-1)} + 16}{3}, \\ k_0(B) &= \frac{2^{2r} + 8}{3}, \\ k_1(B) &= \frac{2^{2(r-1)} + 8}{3}, \\ l(B) &= 3. \end{aligned}$$

*Proof.* By Feit-Thompson we may assume  $O_{2'}(G) = 1$  in order to show that  $G$  is solvable. We apply the  $Z^*$ -theorem. For this let  $g \in G$  such that  ${}^g z \in D$ . Since all involutions of  $D$  are central (in  $D$ ), we get  ${}^g z \in Z(D)$ . By Burnside's fusion theorem there exists  $h \in N_G(D)$  such that  ${}^h z = {}^g z$ . (For principal blocks this would also follow from the fact that  $D$  controls fusion.) Since  $D' = \langle z \rangle$ , we have  ${}^g z = z$ . Now the  $Z^*$ -theorem implies  $z \in Z(G)$ . Then  $D/\langle z \rangle \cong C_{2^r}^2$  is a Sylow 2-subgroup of  $G/\langle z \rangle$ . By Theorem 1 in [4],  $G/\langle z \rangle$  is solvable. Hence, also  $G$  is solvable. Since Alperin's weight conjecture holds for solvable groups, we obtain the numbers  $k(B)$  and  $l(B)$ .

It is also known that the Alperin-McKay-conjecture holds for solvable groups (see [32]). Thus, in order to determine  $k_0(B)$  we may assume  $D \trianglelefteq G$ . Then we can apply the results of [21]. For this let  $L := D \rtimes C_3$ . Then  $B \cong (RL)^{n \times n}$  for some  $n \in \mathbb{N}$ . Hence,  $k_0(B)$  is just the number of irreducible characters of  $L$  with odd degree. By Clifford, every irreducible character of  $L$  is an extension or an induction of a character of  $D$ . Thus, it suffices to count the characters of  $L$  which arise from linear characters of  $D$ . These linear characters of  $D$  are just the inflations of  $\text{Irr}(D/D')$ . They split into the trivial character and orbits of length 3 under the action of  $L$  by Brauer's permutation lemma. The three inflations of  $\text{Irr}(L/D)$  are the extensions of the trivial character of  $D$ . The other linear characters of  $D$  remain irreducible after induction. Characters in the same orbit amount to the same character of  $L$ . This shows

$$k_0(B) = 3 + \frac{|D/D'| - 1}{3} = \frac{2^{2r} + 8}{3}.$$

By Theorem 1.4 in [28] we have  $k_i(B) = 0$  for  $i \geq 2$ . We conclude

$$k_1(B) = k(B) - k_0(B) = \frac{5 \cdot 2^{2(r-1)} + 16}{3} - \frac{2^{2r} + 8}{3} = \frac{2^{2(r-1)} + 8}{3}. \quad \square$$

The last result implies that Brauer's height zero conjecture is also satisfied for blocks of maximal defect. Moreover, the Dade-conjecture holds for solvable groups (see [40]).

Finally we consider the case  $r = 2$  (i. e.  $|D| = 32$ ) for arbitrary groups  $G$ .

**Proposition 4.11.** *If  $r = 2$ , we have*

$$k(B) = 12, \quad k_0(B) = 8, \quad k_1(B) = 4, \quad l(B) = 3.$$

*There are two pairs of 2-conjugate characters of height 0. The remaining characters are 2-rational. Moreover, the Cartan matrix of  $B$  is equivalent to*

$$\begin{pmatrix} 4 & 2 & 2 \\ 2 & 4 & 2 \\ 2 & 2 & 12 \end{pmatrix}.$$

*Proof.* The proof is somewhat lengthy and consists entirely of technical calculations. For this reason we will only outline the argumenation. Since  $k_0(B)$  is divisible by 4, inequality (6) implies  $k_0(B) \geq 8$ . Since there are exactly two pairs of 2-conjugate  $B$ -subsections, Brauer's permutation lemma implies that we also have two pairs of 2-conjugate characters. Hence, the column  $a_1^y$  contains at most four nonvanishing entries. Since  $(a_1^y, a_1^y) = 8$ , there are just two nonvanishing entries, both are  $\pm 2$ . Now Lemma 4.5 implies  $k_0(B) = 8$ . This shows  $(k(B), k_1(B), l(B)) \in \{(12, 4, 3), (14, 6, 5)\}$ .

By way of contradiction, we assume  $k(B) = 14$ . Then one can determine the numbers  $d_{\chi\varphi}^u$  for  $u \neq 1$  with the help of the contributions. However, there are many possibilities. The ordinary decomposition matrix  $Q$  can be computed as the orthogonal space of the other columns of the generalized decomposition matrix. Finally we obtain the Cartan matrix of  $B$  as  $C = Q^T Q$ . In all cases it turns out that  $C$  has the wrong determinant (see Lemma 4.4). This shows  $k(B) = 12$ ,  $k_1(B) = 4$  and  $l(B) = 3$ .

Again we can determine the numbers  $d_{\chi\varphi}^u$  for  $u \neq 1$ . This yields the heights of the 2-conjugate characters. We also obtain some informations about the Cartan invariants in this way. We regard the Cartan matrix  $C$  as a quadratic form. Using the tables [31, 30] we conclude that  $C$  has the form given in the statement of the proposition.  $\square$

## Acknowledgment

The author thanks his advisor Burkhard Külshammer for his encouragement. Proposition 4.9 was his idea.

## References

- [1] J. L. Alperin, *Weights for finite groups*, in The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986), 369–379, Amer. Math. Soc., Providence, RI, 1987.
- [2] H. Bender, *Transitive Gruppen gerader Ordnung, in denen jede Involution genau einen Punkt festläßt*, J. Algebra **17** (1971), 527–554.
- [3] R. Brauer, *On the connection between the ordinary and the modular characters of groups of finite order*, Ann. of Math. (2) **42** (1941), 926–935.
- [4] R. Brauer, *Some applications of the theory of blocks of characters of finite groups. II*, J. Algebra **1** (1964), 307–334.
- [5] R. Brauer, *On blocks and sections in finite groups. I*, Amer. J. Math. **89** (1967), 1115–1136.
- [6] R. Brauer, *On blocks and sections in finite groups. II*, Amer. J. Math. **90** (1968), 895–925.
- [7] R. Brauer, *Some applications of the theory of blocks of characters of finite groups. IV*, J. Algebra **17** (1971), 489–521.
- [8] R. Brauer, *On 2-blocks with dihedral defect groups*, in Symposia Mathematica, Vol. XIII (Convegno di Gruppi e loro Rappresentazioni, INDAM, Rome, 1972), 367–393, Academic Press, London, 1974.
- [9] M. Broué, *On characters of height zero*, in The Santa Cruz Conference on Finite Groups (Univ. California, Santa Cruz, Calif., 1979), 393–396, Amer. Math. Soc., Providence, R.I., 1980.
- [10] D. A. Buell, *Binary quadratic forms*, Springer-Verlag, New York, 1989.
- [11] M. Cabanes and C. Picaronny, *Types of blocks with dihedral or quaternion defect groups*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **39** (1992), 141–161.
- [12] E. C. Dade, *Counting characters in blocks. I*, Invent. Math. **109** (1992), 187–210.
- [13] O. Dövel, *On Donovan's conjecture*, J. Algebra **272** (2004), 1–26.
- [14] K. Erdmann, *Blocks of tame representation type and related algebras*, Lecture Notes in Mathematics, Springer-Verlag, Berlin, 1990.

- [15] W. Feit, *The representation theory of finite groups*, North-Holland Mathematical Library, North-Holland Publishing Co., Amsterdam, 1982.
- [16] M. Fujii, *On determinants of Cartan matrices of  $p$ -blocks*, Proc. Japan Acad. Ser. A Math. Sci. **56** (1980), 401–403.
- [17] I. M. Isaacs, *Character theory of finite groups*, AMS Chelsea Publishing, Providence, RI, 2006.
- [18] R. Kessar and M. Linckelmann, *On perfect isometries for tame blocks*, Bull. London Math. Soc. **34** (2002), 46–54.
- [19] H. Kurzweil and B. Stellmacher, *The theory of finite groups*, Universitext, Springer-Verlag, New York, 2004.
- [20] B. Külshammer, *On 2-blocks with wreathed defect groups*, J. Algebra **64** (1980), 529–555.
- [21] B. Külshammer, *Crossed products and blocks with normal defect groups*, Comm. Algebra **13** (1985), 147–168.
- [22] B. Külshammer and T. Okuyama, *On centrally controlled blocks of finite groups*, unpublished.
- [23] B. Külshammer and L. Puig, *Extensions of nilpotent blocks*, Invent. Math. **102** (1990), 17–71.
- [24] B. Külshammer and T. Wada, *Some inequalities between invariants of blocks*, Arch. Math. (Basel) **79** (2002), 81–86.
- [25] P. Landrock, *On the number of irreducible characters in a 2-block*, J. Algebra **68** (1981), 426–442.
- [26] M. Linckelmann, *Fusion category algebras*, J. Algebra **277** (2004), 222–235.
- [27] V. D. Mazurov, *Finite groups with metacyclic Sylow 2-subgroups*, Sibirsk. Mat. Ž. **8** (1967), 966–982.
- [28] A. Moretó and G. Navarro, *Heights of characters in blocks of  $p$ -solvable groups*, Bull. London Math. Soc. **37** (2005), 373–380.
- [29] H. Nagao and Y. Tsushima, *Representations of finite groups*, Academic Press Inc., Boston, MA, 1989.
- [30] G. Nebe and N. Sloane, *The Brandt-Intrau-Schiemann table of even ternary quadratic forms*, [http://www2.research.att.com/~njas/lattices/Brandt\\_2.html](http://www2.research.att.com/~njas/lattices/Brandt_2.html).
- [31] G. Nebe and N. Sloane, *The Brandt-Intrau-Schiemann table of odd ternary quadratic forms*, [http://www2.research.att.com/~njas/lattices/Brandt\\_1.html](http://www2.research.att.com/~njas/lattices/Brandt_1.html).
- [32] T. Okuyama and M. Wajima, *Irreducible characters of  $p$ -solvable groups*, Proc. Japan Acad. Ser. A Math. Sci. **55** (1979), 309–312.
- [33] J. B. Olsson, *On 2-blocks with quaternion and quasidihedral defect groups*, J. Algebra **36** (1975), 212–241.
- [34] J. B. Olsson, *On the subsections for certain 2-blocks*, J. Algebra **46** (1977), 497–510.
- [35] J. B. Olsson, *Lower defect groups*, Comm. Algebra **8** (1980), 261–288.
- [36] J. B. Olsson, *On subpairs and modular representation theory*, J. Algebra **76** (1982), 261–279.
- [37] S. Park, *The gluing problem for some block fusion systems*, J. Algebra **323** (2010), 1690–1697.
- [38] L. Puig and Y. Usami, *Perfect isometries for blocks with abelian defect groups and cyclic inertial quotients of order 4*, J. Algebra **172** (1995), 205–213.
- [39] G. R. Robinson, *On Brauer’s  $k(B)$  problem*, J. Algebra **147** (1992), 450–455.
- [40] G. R. Robinson, *Dade’s projective conjecture for  $p$ -solvable groups*, J. Algebra **229** (2000), 234–248.
- [41] L. Rédei, *Das „schiefe Produkt“ in der Gruppentheorie*, Comment. Math. Helv. **20** (1947), 225–264.
- [42] B. Sambale, *Cartan matrices and Brauer’s  $k(B)$ -conjecture*, Journal of Algebra (to appear), <http://www.sciencedirect.com/science/article/B6WH2-51H1NP7-1/2/591ade320b5d95adf75b60f4fad8ada>.
- [43] B. Sambale, *Fusion systems on metacyclic 2-groups*, <http://arxiv.org/abs/0908.0783>.

- [44] Y. Usami, *On  $p$ -blocks with abelian defect groups and inertial index 2 or 3. I*, J. Algebra **119** (1988), 123–146.
- [45] Y. Usami, *On  $p$ -blocks with abelian defect groups and inertial index 2 or 3. II*, J. Algebra **122** (1989), 98–105.
- [46] A. Watanabe, *Notes on  $p$ -blocks of characters of finite groups*, J. Algebra **136** (1991), 109–116.
- [47] P. Webb, *An introduction to the representations and cohomology of categories*, in Group representation theory, 149–173, EPFL Press, Lausanne, 2007.