

TOEPLITZ OPERATORS ON BERGMAN SPACES OF POLYANALYTIC FUNCTIONS

ŽELJKO ČUČKOVIĆ AND TRIEU LE

ABSTRACT. We study algebraic properties of Toeplitz operators on Bergman spaces of polyanalytic functions on the unit disk. We obtain results on finite-rank commutators and semi-commutators of Toeplitz operators with harmonic symbols. We also raise and discuss some open questions.

1. INTRODUCTION

The last twenty years have brought a lot of progress in the study of spaces of analytic functions and their operators. Particularly interesting to us are Bergman spaces and Toeplitz operators acting on them. One can notice the development of the field in several directions. One promising direction is to study these operators on Bergman spaces of different domains in \mathbb{C}^n . The other generalization concerns harmonic Bergman spaces. We choose to go to a different generalization, namely we would like to study algebraic properties of Toeplitz operators on Bergman spaces of polyanalytic functions on the unit disk \mathbb{D} in the complex plane. For $n \geq 1$, a function h is called n -analytic (or polyanalytic of order n) on \mathbb{D} if it satisfies the generalized Cauchy-Riemann equation $\frac{\partial^n h}{\partial \bar{z}^n} = 0$. It is clear that h is n -analytic if and only if there are analytic functions h_0, \dots, h_{n-1} on \mathbb{D} such that

$$h(z) = h_0(z) + h_1(z)\bar{z} + \dots + h_{n-1}(z)\bar{z}^{n-1},$$

for all $z \in \mathbb{D}$.

Throughout the paper, we use $L^p(\mathbb{D})$ ($1 \leq p \leq \infty$) to denote the space $L^p(\mathbb{D}, dA)$, where dA is the normalized Lebesgue area measure on \mathbb{D} . The n -analytic Bergman space $A_n^2 = A_n^2(\mathbb{D})$ consists of all n -analytic functions which also belong to the Hilbert space $L^2(\mathbb{D})$ (with inner product denoted by $\langle \cdot, \cdot \rangle$). In other words, a function h belongs to A_n^2 if and only if h is n -analytic and

$$\|h\| = \langle h, h \rangle^{1/2} = \left(\int_{\mathbb{D}} |h(z)|^2 dA(z) \right)^{1/2} < \infty.$$

The space A_1^2 is the familiar Bergman space of the unit disk. It is well known that A_1^2 is a closed subspace of $L^2(\mathbb{D})$. It turns out that the same thing holds for all $n \geq 1$.

2010 *Mathematics Subject Classification.* Primary 47B35; Secondary 46E22.

One of the first studies of A_n^2 appeared in Košelev's paper [10]. We also refer the interested reader to Ramazanov's papers [14, 15] and Balk's book [6] and the references therein for more details about the structure of these spaces.

For any function $f \in L^1(\mathbb{D}, dA)$, the Toeplitz operator T_f is an integral operator defined by the formula

$$(T_f h)(z) = \int_{\mathbb{D}} f(w)h(w)\bar{K}_z(w)dA(w),$$

for $h \in A_n^2$ for which the integral on the right hand side is well defined for all $z \in \mathbb{D}$. Here $K_z(w)$ is the reproducing kernel function of A_n^2 . When f is a bounded function, $T_f h$ is defined for all $h \in A_n^2$ and it is also given by $T_f h = P(fh)$, where P is the orthogonal projection from $L^2(\mathbb{D})$ onto A_n^2 .

The first study of Toeplitz operators on A_n^2 seems to be Wolf's paper [18]. In the paper, Wolf considered Toeplitz operators T_f where the function f is continuous on the closed unit disk. He showed that such an operator is unitarily equivalent to a compact perturbation of the direct sum of n copies of the Hardy-Toeplitz operator whose symbol is the restriction of f on the unit circle. As a consequence, a criterion for Fredholmness and a Fredholm index formula for T_f were obtained.

The goal of the current paper is to use the reproducing kernels and Berezin transform to understand the algebraic and quantitative properties of Toeplitz operators on A_n^2 . We show that certain analytic Bergman space results remain true in the new setting. In particular, we generalize the result of the first author [8] on finite rank perturbations of products of Toeplitz operators with harmonic symbols. Such a generalization is non trivial since the new Berezin transform is no longer one-to-one when $n \geq 2$, unlike the analytic Bergman space case when $n = 1$. We hope to be able to deepen our understanding of operators on these function spaces in our future work.

2. PRELIMINARIES

In this section we discuss in more detail the spaces A_n^2 and their reproducing kernel functions.

We begin by showing that the evaluation functional at each point in the unit disk is bounded on A_n^2 . Recall that the pseudo-hyperbolic distance between two points z and w in the disk is defined by $|\varphi_z(w)|$, where $\varphi_z(w) = \frac{z-w}{1-w\bar{z}}$ is the Moebius transformation of the disk that interchanges 0 and z . For $z \in \mathbb{D}$ and $0 < r < 1$, let $E(z, r) = \{w : |\varphi_z(w)| < r\}$ be the pseudo-hyperbolic disk of radius r centered at z . It is well known (see [19, Proposition 4.4]) that $E(z, r)$ is the Euclidean disk of radius R centered at c , where

$$c = \frac{1-r^2}{1-r^2|z|^2}z, \quad \text{and} \quad R = \frac{1-|z|^2}{1-r^2|z|^2}r.$$

Since $\tilde{R} = R - |c - z| = \frac{(1 - |z|^2)r}{1 + r|z|} > 0$, we see that $E(z, r)$ contains the Euclidean disk $\{w : |w - z| < \tilde{R}\}$.

In [13, Lemma 5], Pavlović showed that there is a constant $C > 0$ such that for any $h \in A_n^2$, $z \in \mathbb{D}$ and $0 < \delta < 1 - |z|$,

$$|h(z)|^2 \leq \frac{C}{\delta^2} \int_{|w-z|<\delta} |h(w)|^2 dA(w). \quad (2.1)$$

Pavlović in fact studied spaces of polyharmonic functions but since any n -analytic functions is also n -harmonic, his lemma may be used in the current setting. Applying (2.1) with $\delta = \tilde{R}$ and using the inclusion $\{w : |w - z| < \tilde{R}\} \subset E(z, r)$, we obtain

$$\begin{aligned} |h(z)|^2 &\leq \frac{C(1+r|z|)^2}{r^2(1-|z|^2)^2} \int_{|w-z|<\tilde{R}} |h(w)|^2 dA(w) \\ &\leq \frac{C(1+r)^2}{r^2(1-|z|^2)^2} \int_{E(z,r)} |h(w)|^2 dA(w) \\ &= \frac{C_r}{(1-|z|^2)^2} \int_{E(z,r)} |h(w)|^2 dA(w), \end{aligned} \quad (2.2)$$

where C_r is independent of h and z . This implies that A_n^2 is a closed subspace of $L^2(\mathbb{D})$ and that the functional $h \mapsto h(z)$ is bounded on A_n^2 for any $z \in \mathbb{D}$. Therefore there is a function K_z in A_n^2 such that $h(z) = \langle h, K_z \rangle$ for $z \in \mathbb{D}$. The function $K(w, z) = K_z(w)$ for $(z, w) \in \mathbb{D} \times \mathbb{D}$ is called, as usual, the kernel function for A_n^2 . It can be showed (see [10]) that

$$\begin{aligned} K_z(w) &= \frac{n}{(1-w\bar{z})^{2n}} \sum_{j=0}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} |1-w\bar{z}|^{2(n-1-j)} |w-z|^{2j} \\ &= \frac{n|1-w\bar{z}|^{2(n-1)}}{(1-w\bar{z})^{2n}} \sum_{j=0}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} |\varphi_z(w)|^{2j}. \end{aligned}$$

Using properties of reproducing kernel functions we obtain, for $z \in \mathbb{D}$, $\|K_z\|^2 = \langle K_z, K_z \rangle = K_z(z) = n^2/(1-|z|^2)^2$. The normalized reproducing kernel at z is defined by

$$\begin{aligned} k_z(w) &= \frac{K_z(w)}{\|K_z\|} = \frac{1-|z|^2}{n} K_z(w) \\ &= \frac{(1-|z|^2)|1-w\bar{z}|^{2(n-1)}}{(1-w\bar{z})^{2n}} \sum_{j=0}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} |\varphi_z(w)|^{2j}. \end{aligned} \quad (2.3)$$

Using the formula $1 - |\varphi_z(w)|^2 = \frac{(1-|z|^2)(1-|w|^2)}{|1-w\bar{z}|^2}$ we obtain

$$|k_z(w)| = \frac{1-|z|^2}{|1-w\bar{z}|^2} \left| \sum_{j=0}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} |\varphi_z(w)|^{2j} \right|$$

$$\begin{aligned}
&= \frac{1 - |\varphi_z(w)|^2}{1 - |w|^2} \left| \sum_{j=0}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} |\varphi_z(w)|^{2j} \right| \\
&= \frac{1 - |\varphi_z(w)|^2}{1 - |w|^2} \left| n + \sum_{j=1}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} |\varphi_z(w)|^{2j} \right|.
\end{aligned}$$

The last formula shows that $(1 - |w|^2)|k_z(w)| \rightarrow n$ as $|\varphi_z(w)| \rightarrow 0$, that is, for any $\epsilon > 0$, there is a $\delta > 0$ such that $|(1 - |w|^2)|k_z(w)| - n| < \epsilon$ for all $w, z \in \mathbb{D}$ satisfying $|\varphi_z(w)| < \delta$. This implies that there is a number $0 < r < 1$ such that $(1 - |w|^2)|k_z(w)| > n/2 \geq 1/2$ whenever $|\varphi_z(w)| < r$. Using this we obtain the following estimate.

Lemma 2.1. *There is a constant $c > 0$ such that for any function $h \in A_n^2$ and any $w \in \mathbb{D}$,*

$$|h(w)|^2 \leq c \int_{\mathbb{D}} |h(z)|^2 |k_z(w)|^2 dA(z).$$

Proof. Let r be the number mentioned in the paragraph preceding the lemma. Since $|k_z(w)| > \frac{1}{2(1-|w|^2)}$ for all $z \in E(w, r)$, using (2.2), we have

$$\int_{\mathbb{D}} |h(z)|^2 |k_z(w)|^2 dA(z) \geq \frac{1}{4(1-|w|^2)^2} \int_{E(w, r)} |h(z)|^2 dA(z) \geq \frac{|h(w)|^2}{4C_r}.$$

Set $c = 4C_r$. The conclusion of the lemma follows. \square

The last lemma of this section, which is well known in the case of analytic Bergman space, is a consequence of the fact that $\|K_z\| \rightarrow \infty$ as $|z| \uparrow 1$ and n -analytic polynomials are dense in A_n^2 .

Lemma 2.2. *We have $k_z \rightarrow 0$ weakly as $|z| \uparrow 1$. As a result, for any $f \in A_n^2$,*

$$\lim_{|z| \uparrow 1} (1 - |z|^2) f(z) = n \cdot \lim_{|z| \uparrow 1} \langle f, k_z \rangle = 0.$$

3. WEIGHTED BEREZIN TRANSFORM

For any real number $\alpha \geq 0$, the weighted Berezin transform B_α is defined by

$$B_\alpha u(z) = (\alpha + 1) \int_{\mathbb{D}} u(\varphi_z(w)) (1 - |w|^2)^\alpha dA(w)$$

for u in $L^1(\mathbb{D})$ and z in \mathbb{D} . Note that B_0 is just the standard unweighted Berezin transform.

It follows from the mean value theorem for harmonic functions that if $u \in L^1(\mathbb{D})$ is harmonic, then $B_\alpha u = u$. Fixed points of B_0 (even in the setting of the unit ball in higher dimensions) were studied by Ahern, Flores and Rudin in [4]. In the current setting of the unit disk, their results say that all fixed points of B_0 are integrable harmonic functions. This is not

true for all B_α and the interested reader is referred to [19, Chapter 6] and the references therein.

By a change of variable, we see that

$$B_\alpha u(z) = (\alpha + 1) \int_{\mathbb{D}} u(w) \frac{(1 - |z|^2)^{2+\alpha} (1 - |w|^2)^\alpha}{|1 - z\bar{w}|^{2\alpha+4}} dA(w).$$

This shows that $B_\alpha u$ is a real analytic function on \mathbb{D} , hence it is infinitely differentiable.

There is a relation between $B_{\alpha+1}$ and B_α that involves the invariant Laplacian $\tilde{\Delta}$. Recall that $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} = 4 \frac{\partial^2}{\partial z \partial \bar{z}}$ is the ordinary Laplacian. The invariant Laplacian is defined by $(\tilde{\Delta}u)(z) = (1 - |z|^2)^2 (\Delta u)(z)$ for any twice differentiable function u on \mathbb{D} .

Lemma 3.1 ([4, Proposition 2.4] and also [19, Lemma 6.23]). *If $u \in L^1(\mathbb{D})$, then*

$$\tilde{\Delta}(B_\alpha u) = 4(\alpha + 1)(\alpha + 2)(B_\alpha u - B_{\alpha+1}u).$$

This shows that for such u , $B_{\alpha+1}u = \left(1 - \frac{\tilde{\Delta}}{4(\alpha + 1)(\alpha + 2)}\right) B_\alpha u$.

Lemma 3.1 implies that for any integer $k \geq 1$, we have $B_k u = q_k(\tilde{\Delta})(B_0 u)$ for $u \in L^1(\mathbb{D})$, where $q_k(\lambda) = \prod_{j=1}^k \left(1 - \frac{\lambda}{4j(j+1)}\right)$.

It was showed by Ahern and Čučković in [2] that the Berezin transform B_0 and the invariant Laplacian $\tilde{\Delta}$ commute in the following sense: if u is in $C^2(\mathbb{D})$ such that both u and $\tilde{\Delta}u$ belong to $L^1(\mathbb{D})$, then $\tilde{\Delta}B_0 u = B_0 \tilde{\Delta}u$. By applying this identity multiple times, we see that if u is $2s$ times continuously differentiable such that u and $(\tilde{\Delta})^j u$ belong to $L^1(\mathbb{D})$ for all $1 \leq j \leq s$, then

$$q(\tilde{\Delta})B_0 u = B_0(q(\tilde{\Delta})u) \quad (3.1)$$

for any polynomial q of degree s .

In the study of finite sums of finite products of Toeplitz operators with bounded harmonic symbols, one often encounters the problem of classifying integrable functions u for which $B_0(u)$ satisfies certain conditions. The following theorem due to N.V. Rao [16] shows that for locally bounded u , $B_0 u$ cannot be of the form $f_1 \bar{g}_1 + \cdots + f_m \bar{g}_m$ with analytic functions f_1, \dots, f_m and g_1, \dots, g_m unless u is harmonic. The proof uses recent results on finite rank Toeplitz operators (with distributional symbols) on the analytic Bergman space. In fact Rao considered $u \in L^1(\mathbb{D})$ without any boundedness assumption but the conclusion is different and we do not need it here for our goal.

Theorem 3.2. *Suppose u is an integrable function that is locally bounded on \mathbb{D} such that $B_0(u) = f_1 \bar{g}_1 + \cdots + f_m \bar{g}_m$ for some positive integer m , where f_1, \dots, f_m and g_1, \dots, g_m are analytic on \mathbb{D} . Then u is harmonic, or equivalently, $\tilde{\Delta}u = 0$ on \mathbb{D} , and we have $u = f_1 \bar{g}_1 + \cdots + f_m \bar{g}_m$.*

Remark 3.3. For analytic functions f_1, \dots, f_m and g_1, \dots, g_m , Theorem 3.3 in [7] shows that the function $f_1\bar{g}_1 + \dots + f_m\bar{g}_m$ is harmonic if and only if $(f_1 - f_1(0))(\bar{g}_1 - \bar{g}_1(0)) + \dots + (f_m - f_m(0))(\bar{g}_m - \bar{g}_m(0)) = 0$ on \mathbb{D} . In particular, when $m = 1$, we have $(f_1 - f_1(0))(g_1 - g_1(0)) = 0$, which implies either f_1 or g_1 is a constant function.

4. TOEPLITZ OPERATORS ON A_n^2

Recall that for any function $f \in L^1(\mathbb{D}, dA)$, the Toeplitz operator T_f is defined by the formula

$$(T_f h)(z) = \int_{\mathbb{D}} f(w)h(w)\bar{K}_z(w)dA(w), \quad (4.1)$$

for $h \in A_n^2$ for which the integral on the right hand side is well defined for all $z \in \mathbb{D}$. Since the function $\bar{K}_z(w)$ is n -analytic in the variable z , $T_f h$ is n -analytic on \mathbb{D} . Note that the domain of T_f contains the space \mathcal{W} of all linear combinations of kernel functions K_z , which is dense in A_n^2 . We say that T_f is bounded on A_n^2 if there is a constant $C > 0$ such that for any $h \in \mathcal{W}$, $T_f h$ belongs to A_n^2 and $\|T_f h\| \leq C\|h\|$. In this case T_f extends uniquely to a bounded operator on A_n^2 . Also for $h \in A_n^2$ and $g = \sum_{j=1}^m \alpha_j K_{z_j} \in \mathcal{W}$, we have

$$\begin{aligned} \langle T_f h, g \rangle &= \sum_{j=1}^m \bar{\alpha}_j \langle T_f h, K_{z_j} \rangle = \sum_{j=1}^m \bar{\alpha}_j (T_f h)(z_j) \\ &= \sum_{j=1}^m \bar{\alpha}_j \int_{\mathbb{D}} f(w)h(w)\bar{K}_{z_j}(w)dA(w) \quad (\text{by (4.1)}) \\ &= \int_{\mathbb{D}} f(w)h(w)\bar{g}(w)dA(w). \end{aligned} \quad (4.2)$$

Let P denote the orthogonal projection of $L^2(\mathbb{D})$ onto A_n^2 . If f belongs to $L^2(\mathbb{D})$, then for any bounded h in A_n^2 , the function fh belongs to $L^2(\mathbb{D})$. This implies that

$$P(fh)(z) = \langle P(fh), K_z \rangle = \langle fh, K_z \rangle = \int_{\mathbb{D}} f(w)h(w)\bar{K}_z(w)dA(w) = (T_f h)(z).$$

We then recover the usual definition of T_f , which is given by $T_f h = P(fh)$.

If f is a bounded function, then T_f is a bounded operator on A_n^2 and we have $\|T_f\| \leq \|f\|_{\infty}$ and $T_f^* = T_{\bar{f}}$.

If f is analytic (not necessarily bounded), then T_f is just the operator of multiplication by f . Also for any z, w in \mathbb{D} , we have

$$\langle T_{\bar{f}} K_z, K_w \rangle = \langle K_z, f K_w \rangle = \bar{f}(z)\bar{K}_w(z) = \langle \bar{f}(z)K_z, K_w \rangle.$$

Using the density in A_n^2 of the linear span of $\{K_w : w \in \mathbb{D}\}$, we conclude that $T_{\bar{f}} K_z = \bar{f}(z)K_z$ for $z \in \mathbb{D}$.

Suppose u is a harmonic function that belongs to $L^2(\mathbb{D})$. Then there are analytic functions f, g in $L^2(\mathbb{D})$ such that $u = f + \bar{g}$ and hence,

$$\langle T_u K_z, K_w \rangle = \langle \bar{g} K_z, K_w \rangle + \langle K_z, \bar{f} K_w \rangle = (f(w) + \bar{g}(z)) \langle K_z, K_w \rangle$$

for all z, w in \mathbb{D} . This shows, in particular, that $\langle T_u K_z, K_z \rangle = u(z) \|K_z\|^2$ for any $z \in \mathbb{D}$. Hence, if T_u is a bounded operator, then $\|u\|_\infty \leq \|T_u\|$. On the other hand, as we mentioned above, $\|T_u\| \leq \|u\|_\infty$ whenever u is bounded. We then conclude that $\|T_u\| = \|u\|_\infty$.

Using the kernel function $K(z, w)$, we define the Berezin transform of a bounded operator T on A_n^2 as $B(T)(z) = \|K_z\|^{-2} \langle T K_z, K_z \rangle = \langle T k_z, k_z \rangle$ for $z \in \mathbb{D}$. (Recall that $k_z = \|K_z\|^{-1} K_z$ is the normalized reproducing kernel at z .) It follows from the formula for K_z that B maps the space $\mathcal{B}(A_n^2)$ of bounded linear operators on A_n^2 to the space of real analytic functions on \mathbb{D} . Also $\|B(T)\| \leq \|T\|$ for all $T \in \mathcal{B}(A_n^2)$.

Suppose $f \in L^1(\mathbb{D})$ such that T_f is a bounded operator on $A_n^2(\mathbb{D})$. Then for $z \in \mathbb{D}$, using the formula for k_z as in (2.3), we have

$$\begin{aligned} B(T_f)(z) &= \langle T_f k_z, k_z \rangle = \int_{\mathbb{D}} f(w) |k_z(w)|^2 dA(w) \quad (\text{by (4.2)}) \\ &= \int_{\mathbb{D}} f(w) \frac{(1 - |z|^2)^2}{|1 - w\bar{z}|^4} \left\{ \sum_{j=0}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} |\varphi_z(w)|^{2j} \right\}^2 dA(w). \end{aligned}$$

Note that the last integral is defined for all z in \mathbb{D} and all f in $L^1(\mathbb{D})$ regardless of the boundedness of T_f . We will denote it by $Bf(z)$ and call the function Bf the Berezin transform of f .

4.1. Compactness of Toeplitz operators with non-negative symbols. Compactness and Schatten class membership of Toeplitz operators (on analytic Bergman spaces) with non-negative symbols can be characterized using the Berezin transform of the symbol. See [19, Chapter 7] for more details. We will show that this characterization (for simplicity, we will only work on the compactness characterization) carries over to n -analytic Bergman spaces with a similar approach.

Lemma 4.1. *There is a constant $c > 0$ so that for any non-negative function $f \in L^1(\mathbb{D})$ such that $\tilde{f} = Bf$ belongs to $L^1(\mathbb{D})$ and $T_{\tilde{f}}$ is bounded on A_n^2 , we have $T_f \leq cT_{\tilde{f}}$.*

Proof. Let h be a linear combination of kernel functions. We have

$$\begin{aligned} \langle T_{\tilde{f}} h, h \rangle &= \int_{\mathbb{D}} \tilde{f}(z) |h(z)|^2 dA(z) \quad (\text{by (4.2)}) \\ &= \int_{\mathbb{D}} \int_{\mathbb{D}} f(w) |k_z(w)|^2 |h(z)|^2 dA(w) dA(z) \\ &= \int_{\mathbb{D}} \left(\int_{\mathbb{D}} |k_z(w)|^2 |h(z)|^2 dA(z) \right) f(w) dA(w) \end{aligned}$$

$$\begin{aligned}
&\geq \frac{1}{c} \int_{\mathbb{D}} |h(w)|^2 f(w) dA(w) \quad (\text{by Lemma 2.1}) \\
&= \frac{1}{c} \langle T_f h, h \rangle.
\end{aligned}$$

Since the above inequalities hold for all h in a dense subset of A_n^2 and $T_{\tilde{f}}$ is bounded, we conclude that $T_f \leq cT_{\tilde{f}}$. \square

Suppose f is a bounded function whose support is a compact subset of the unit disk. Since T_f is an integral operator with kernel $f(w)\bar{K}_z(w)$, which is bounded in $(z, w) \in \mathbb{D} \times \mathbb{D}$, we see that T_f is a Hilbert-Schmidt operator, hence compact. Using an approximation argument, we obtain the next lemma, which is well known in the case of analytic Bergman space A_1^2 .

Lemma 4.2. *Suppose f is a bounded function such that $\lim_{|z| \uparrow 1} f(z) = 0$. Then the operator T_f is compact on A_n^2 .*

We are now able to characterize compact Toeplitz operators on A_n^2 with non-negative symbols. These results are analogous to the corresponding results on the analytic Bergman space A_1^2 .

Theorem 4.3. *Let $f \in L^1(\mathbb{D})$ be a non-negative function. Then T_f is compact on A_n^2 if and only if $Bf(z) \rightarrow 0$ as $|z| \uparrow 1$.*

Proof. Suppose T_f is compact. Since $k_z \rightarrow 0$ weakly as $|z| \uparrow 1$ by Lemma 2.2, we have $Bf(z) = B(T_f)(z) = \langle T_f k_z, k_z \rangle \rightarrow 0$ as $|z| \uparrow 1$.

Conversely, suppose $\tilde{f}(z) = Bf(z) \rightarrow 0$ as $|z| \uparrow 1$. This, in particular, implies that \tilde{f} is bounded. Lemma 4.2 then shows that $T_{\tilde{f}}$ is compact. Using Lemma 4.1, we conclude that T_f is also compact. \square

Remark 4.4. It is possible to introduce the notion of Carleson measure on A_n^2 and use it to obtain Theorem 4.3 as it was done for the analytic Bergman space A_1^2 in [19, Section 7.2]. We leave the details to the interested reader.

Remark 4.5. On the analytic Bergman space $A_1^2(\mathbb{D})$, Axler and Zheng showed (see the equivalence of (a) and (c) in [19, Theorem 7.23]) that for a bounded function f , the Toeplitz operator T_f is compact if and only if $B_0(f)(z) \rightarrow 0$ as $|z| \uparrow 1$. We do not know if this result can be extended to any A_n^2 with $n \geq 2$. (Axler-Zheng's proof, when applied to A_n^2 with $n \geq 2$, faces many difficulties which cannot be resolved.) Note that in Theorem 4.3, we do not require the boundedness of f but we do require the positivity, which is crucial for the proof to work.

4.2. Toeplitz operators with harmonic symbols. Suppose u and v are two bounded harmonic functions such that $u = f + \bar{g}$ and $v = h + \bar{k}$, where f, g, h, k are analytic. For any z, w in \mathbb{D} , we have

$$\begin{aligned}
\langle T_u T_v K_z, K_w \rangle &= \langle T_v K_z, T_{\bar{u}} K_w \rangle \\
&= \langle hK_z + \bar{k}(z)K_z, \bar{f}(w)K_w + gK_w \rangle \\
&= (f(w)h(w) + f(w)\bar{k}(z) + \bar{g}(z)\bar{k}(z)) \langle K_z, K_w \rangle + \langle \bar{g}hK_z, K_w \rangle.
\end{aligned} \tag{4.3}$$

In particular, by letting $w = z$ and dividing by $\|K_z\|^2$ we obtain

$$B(T_u T_v) = fh + f\bar{k} + \bar{g}\bar{k} + B(T_{\bar{g}h}) = uv - \bar{g}h + B(T_{\bar{g}h}). \quad (4.4)$$

For two non-zero functions x and y in A_n^2 , the rank-one operator $x \otimes y$ is defined by $(x \otimes y)(f) = \langle f, y \rangle x$ for $f \in A_n^2$. For z, w in \mathbb{D} , we have

$$\langle (x \otimes y)K_z, K_w \rangle = \langle K_z, y \rangle \langle x, K_w \rangle = x(w)\bar{y}(z).$$

This implies

$$B(x \otimes y)(z) = \frac{(1 - |z|^2)^2}{n^2} x(z)\bar{y}(z). \quad (4.5)$$

The Berezin transform is injective on the space of bounded linear operators on A_1^2 : for any $\alpha \geq 0$, if $B_\alpha(T)$ is the zero function on \mathbb{D} , then $T = 0$ (see [19, Proposition 6.2]). It turns out that this is not the case on A_n^2 when $n \geq 2$. In fact, if $f(z) = z$, then f and \bar{f} belong to A_n^2 and by (4.5), $B(f \otimes 1) = B(1 \otimes \bar{f})$ but it is clear that $f \otimes 1 \neq 1 \otimes \bar{f}$. We mention in passing that this example also serves as an example to show that the Berezin transform is not injective on bounded linear operators acting on the harmonic Bergman space of the unit disk. Examples involving Toeplitz operators on A_n^2 can also be produced as follows.

Example 4.6 (Non-injectivity of Berezin transform). Choose analytic monomials u_1, u_2, u_3 and conjugate analytic monomials v_1, v_2, v_3 such that

$$\frac{(1 - \bar{z}w)^2}{n^2} = u_1(w)v_1(z) + u_2(w)v_2(z) + u_3(w)v_3(z).$$

Using (4.4) and (4.5) together with the linearity of the Berezin transform, we obtain

$$B\left(\sum_{j=1}^3 T_{u_j} T_{v_j}\right)(z) = \sum_{j=1}^3 u_j(z)v_j(z) = \frac{(1 - |z|^2)^2}{n^2} = B(1 \otimes 1)(z)$$

for any $z \in \mathbb{D}$. On the other hand, from (4.3),

$$\begin{aligned} \left\langle \left(\sum_{j=1}^3 T_{u_j} T_{v_j}\right)K_z, K_w \right\rangle &= \left\langle \sum_{j=1}^3 u_j(w)v_j(z) \right\rangle \langle K_z, K_w \rangle = \frac{(1 - \bar{z}w)^2}{n^2} K_z(w), \\ \langle (1 \otimes 1)K_z, K_w \rangle &= \langle K_z, 1 \rangle \langle 1, K_w \rangle = 1, \end{aligned}$$

for all z and w in \mathbb{D} . Since $K_z(w)$ is not the same as $\frac{n^2}{(1 - \bar{z}w)^2}$ for $n \geq 2$, we conclude that $T_{u_1} T_{v_1} + T_{u_2} T_{v_2} + T_{u_3} T_{v_3} \neq 1 \otimes 1$.

Even though the Berezin transform is not injective on the space of bounded operators on A_n^2 , the implication $Bf = 0 \implies f = 0$ is in fact valid if appropriate conditions are imposed on f (in particular, it is true for all bounded functions f). We will see this by analyzing the relation between B and B_0 ,

the unweighted Berezin transform. Recall that for $f \in L^1$ and $z \in \mathbb{D}$, the formula for $Bf(z)$ has the form

$$\begin{aligned} Bf(z) &= \int_{\mathbb{D}} f(w) \frac{(1 - |z|^2)^2}{|1 - w\bar{z}|^4} \left\{ \sum_{j=0}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} |\varphi_z(w)|^{2j} \right\}^2 dA(w) \\ &= \int_{\mathbb{D}} f(\varphi_z(\zeta)) \left\{ \sum_{j=0}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} |\zeta|^{2j} \right\}^2 dA(\zeta), \end{aligned}$$

by the change of variable $w = \varphi_z(\zeta)$. Let μ be the polynomial of degree $2n - 2$ defined by

$$\mu(t) = \left\{ \sum_{j=0}^{n-1} (-1)^j \binom{n}{j+1} \binom{n+j}{n} t^j \right\}^2.$$

Rewriting $\mu(t) = b_0 + b_1(1-t) + \dots + b_{2n-2}(1-t)^{2n-2}$ and using the remark after Lemma 3.1 we see that Bf can be written as

$$\begin{aligned} Bf &= b_0 B_0(f) + \frac{b_1}{2} B_1(f) + \dots + \frac{b_{2n-2}}{2n-1} B_{2n-2}(f) \\ &= b_0 B_0(f) + \frac{b_1}{2} q_1(\tilde{\Delta}) B_0(f) + \dots + \frac{b_{2n-2}}{2n-1} q_{2n-2}(\tilde{\Delta}) B_0(f) \\ &= Q(\tilde{\Delta}) B_0(f), \end{aligned} \tag{4.6}$$

where $Q(t) = b_0 + \frac{b_1}{2} q_1(t) + \dots + \frac{b_{2n-2}}{2n-1} q_{2n-2}(t)$ and $q_k(t) = \prod_{j=1}^k \left(1 - \frac{t}{4j(j+1)}\right)$ for $1 \leq k \leq 2n - 2$. The formula $Bf = Q(\tilde{\Delta}) B_0 f$ implies that in order to understand B , we need to understand $Q(\tilde{\Delta})$ and B_0 .

We begin with a result regarding the eigenspaces of the invariant Laplacian. For any complex number λ , let X_λ denote the linear space of all twice differentiable functions u on \mathbb{D} such that $\tilde{\Delta}u = \lambda u$. Theorem 4.2.7 in [17] shows that $X_\lambda \cap L^\infty(\mathbb{D}) \neq \emptyset$ if and only if λ belongs to the set

$$\Omega_\infty = \{\lambda \in \mathbb{C} : 4\operatorname{Re} \lambda + (\operatorname{Im} \lambda)^2 \leq 0\}. \tag{4.7}$$

This shows that if $u \in C^2(\mathbb{D}) \cap L^\infty(\mathbb{D})$, $\lambda \notin \Omega_\infty$ and $\tilde{\Delta}u = \lambda u$, then $u = 0$. As a consequence, we obtain

Lemma 4.7. *Let q be a polynomial of degree $s \geq 0$ whose roots lie outside Ω_∞ . Suppose u is a function which is $2s+2$ times continuously differentiable on \mathbb{D} such that $q(\tilde{\Delta})\tilde{\Delta}u = 0$ and $(\tilde{\Delta})^j u$ is bounded for all $1 \leq j \leq s$. Then $\tilde{\Delta}u = 0$, that is, u is harmonic.*

We now need to study the locations of the roots of the polynomial $Q(t)$. Because of the complexity of Q , we have not fully understood the locations of its roots for all $n \geq 2$. With the help of Maple, we are able to show that for $2 \leq n \leq 25$, all roots of Q lie outside Ω_∞ . As a result, Lemma 4.7 can be applied to Q . We illustrate here the case $n = 2$ in which the computation

can be carried out by hand. With $n = 2$, we have $\mu(t) = (2 - 3t)^2 = 1 - 6(1 - t) + 9(1 - t)^2$ and hence

$$Q(t) = 1 - 3q_1(t) + 3q_2(t) = 1 - \frac{t}{8} + \frac{t^2}{64}.$$

Since the roots of Q are $t = 4 \pm 4\sqrt{-3}$, they lie outside Ω_∞ . Note that in this case all the roots have positive real parts, however this is not true for other values of n .

Motivated by the conditions in Lemma 4.7, we define \mathcal{S} to be the space of all smooth functions g on \mathbb{D} such that $z \mapsto (1 - |z|^2)^{k+l} \frac{\partial^{k+l} g}{\partial z^k \partial \bar{z}^l}(z)$ is bounded on \mathbb{D} for all $0 \leq k + l$. Using Leibniz's rule, it can be verified that \mathcal{S} is an algebra, that is, $g_1 g_2$ belongs to \mathcal{S} whenever g_1 and g_2 belong to \mathcal{S} . Leibniz's rule also implies that \mathcal{S} is invariant under the action of $\tilde{\Delta}$, that is, $\tilde{\Delta}g$ belongs to \mathcal{S} whenever g belongs to \mathcal{S} .

Recall that the Bloch space \mathcal{B} consists of all analytic functions f on \mathbb{D} such that the function $z \mapsto (1 - |z|^2)f'(z)$ is bounded on \mathbb{D} . Theorem 5.4 in [19] shows that if f belongs to \mathcal{B} then $(1 - |z|^2)^m f^{(m)}(z)$ is bounded on \mathbb{D} for all $m \geq 1$. It follows, by induction, that functions of the form $(1 - |z|^2)^m f^{(m)}(z)$ ($m \geq 1$), where $f \in \mathcal{B}$, belong to \mathcal{S} .

If u is a bounded harmonic function on \mathbb{D} then as well known, there are functions f and g in \mathcal{B} such that $u = f + \bar{g}$. It then follows that u belongs to \mathcal{S} and hence, \mathcal{S} contains all finite sums of finite products of bounded harmonic functions.

Remark 4.8. Any function u in \mathcal{S} satisfies the hypothesis of Lemma 4.7. Therefore if q is a polynomial whose roots lie outside Ω_∞ and $q(\tilde{\Delta})\tilde{\Delta}u = 0$, then $\tilde{\Delta}u = 0$.

We are now ready to prove our main theorem. As we discussed above, we need to restrict ourselves to A_n^2 with $1 \leq n \leq 25$. We believe that this restriction is artificial but we are not able to remove it at this time.

Theorem 4.9. *Let u_1, \dots, u_d and v_1, \dots, v_d be bounded harmonic functions on \mathbb{D} . For each j we write $u_j = f_j + \bar{g}_j$ and $v_j = h_j + \bar{k}_j$, where f_j, g_j, h_j, k_j belong to the Bloch space. Let F be a function in \mathcal{S} (for example, F may be taken to be a finite sum of finite products of bounded harmonic functions on \mathbb{D}). If*

$$\sum_{j=1}^d T_{u_j} T_{v_j} - T_F = \sum_{l=1}^s x_l \otimes y_l, \quad (4.8)$$

where x_l, y_l belong to A_n^2 , then the following statements hold

- (i) $\sum_{j=1}^d h_j \bar{g}_j - F$ is harmonic on \mathbb{D} .
- (ii) $\sum_{j=1}^d u_j(z)v_j(z) - F(z) = \frac{(1-|z|^2)^2}{n^2} \sum_{l=1}^s x_l(z)\bar{y}_l(z)$ for $z \in \mathbb{D}$.

Proof. For any $w, z \in \mathbb{D}$, we have

$$\begin{aligned} \left\langle \left(\sum_{l=1}^s x_l \otimes y_l \right) k_z, k_w \right\rangle &= \sum_{l=1}^s \langle x_l, k_w \rangle \langle k_z, y_l \rangle \\ &= \sum_{l=1}^s x_l(w) \bar{y}_l(z) (\|K_z\| \|K_w\|)^{-1}. \end{aligned}$$

If (4.8) holds, then by applying Berezin transform on both sides and using (4.4) and the above identities, we obtain

$$\begin{aligned} B\left(\sum_{j=1}^d h_j \bar{g}_j\right)(z) + \sum_{j=1}^d f_j(z) h_j(z) + f_j(z) \bar{k}_j(z) + \bar{g}_j(z) \bar{k}_j(z) - B(F) \\ = \|K_z\|^{-2} \sum_{l=1}^s x_l(z) \bar{y}_l(z) = \frac{(1 - |z|^2)^2}{n^2} \sum_{l=1}^s x_l(z) \bar{y}_l(z) \quad (4.9) \end{aligned}$$

for all $z \in \mathbb{D}$.

Put $u = \sum_{j=1}^d h_j \bar{g}_j - F$. Since $\tilde{\Delta}u$ belongs to \mathcal{S} , we may apply (4.6) and then (3.1) to obtain $B(u) = Q(\tilde{\Delta})B_0u = B_0(Q(\tilde{\Delta})u)$. Since x_l and y_l are n -analytic, the function $n^{-2}(1 - |z|^2)^2 x_l(z) \bar{y}_l(z)$ may be written as a finite sum of products of the form $f\bar{g}$, where f and g are analytic on \mathbb{D} . Now, (4.9) implies that $B_0(Q(\tilde{\Delta})u)$ is also a finite sum of such products. Using Theorem 3.2, we conclude that $\tilde{\Delta}Q(\tilde{\Delta})u = 0$, or equivalently, $Q(\tilde{\Delta})(\tilde{\Delta}u) = 0$.

Since all roots of Q lie outside Ω_∞ and $\tilde{\Delta}u$ belongs to \mathcal{S} , Remark 4.8 shows that $\tilde{\Delta}u = 0$, which proves the first statement in the conclusion of the theorem. The second statement follows from (4.9) and the fact that harmonic functions are fixed by the Berezin transform. \square

Theorem 4.9 enjoys many interesting consequences which are analogous to the case of unweighted Bergman space A_1^2 . The following results, for A_1^2 , were obtained by Guo, Sun and Zheng in [9, Section 4]. Using Theorem 4.9, we see that they also hold for A_n^2 .

Corollary 4.10. *If u and v are bounded harmonic functions on \mathbb{D} such that $T_u T_v$ is of finite rank, then either u or v is the zero function.*

Proof. Write $u = f + \bar{g}$ and $v = h + \bar{k}$, where f, g, h, k belong to the Bloch space. Suppose $T_u T_v = \sum_{l=1}^s x_l \otimes y_l$ for some $s \geq 1$ and x_l, y_l are in A_n^2 . By Theorem 4.9, $h\bar{g}$ is harmonic and

$$u(z)v(z) = \frac{(1 - |z|^2)^2}{n^2} \sum_{l=1}^s x_l(z) \bar{y}_l(z)$$

for $z \in \mathbb{D}$. Since $\lim_{|z| \uparrow 1} (1 - |z|^2)\varphi(z) = 0$ for all φ belonging to A_n^2 by Lemma 2.2, we conclude that $\lim_{|z| \uparrow 1} u(z)v(z) = 0$.

Since $h\bar{g}$ is harmonic, either h or g must be a constant function. This implies that either \bar{v} or u is analytic on \mathbb{D} . It then follows that either u or v is the zero function (see the proof of [7, Corollary 3.11]). \square

Corollary 4.11. *If u and v are non-constant bounded harmonic functions on \mathbb{D} such that $[T_u, T_v] = T_u T_v - T_v T_u$ is of finite rank, then either both u and v are analytic, or both \bar{u} and \bar{v} are analytic, or $u = \alpha v + \beta$ for some complex numbers α and β .*

Proof. Write $u = f + \bar{g}$ and $v = h + \bar{k}$ as in the proof of Corollary 4.10. By Theorem 4.9 part (i), $h\bar{g} - f\bar{k}$ is harmonic. The conclusion of the corollary follows from the same argument as the proof of Theorem 1 in [5, p.10]. \square

Corollary 4.12. *If u and v are bounded harmonic functions on \mathbb{D} such that the semi-commutator $[T_u, T_v] = T_u T_v - T_{uv}$ is of finite rank, then either \bar{u} or v is analytic.*

Proof. Write $u = f + \bar{g}$ and $v = h + \bar{k}$ as in the proof of Corollary 4.10. By Theorem 4.9 part (i), $h\bar{g} - uv$ is harmonic. This shows that $f\bar{k}$ is harmonic, which implies either f or k is a constant function. The conclusion of the corollary then follows. \square

Corollary 4.13. *If u and v are bounded harmonic functions and F is a function in \mathcal{S} such that $T_u T_v = T_F$, then either \bar{u} or v is analytic and $F = \bar{u}v$.*

Proof. As before, write $u = f + \bar{g}$ and $v = h + \bar{k}$. By Theorem 4.9, $F = uv$ and $h\bar{g} - F$ is harmonic, which implies that $f\bar{k}$ is harmonic. Therefore either f or k is a constant function. The conclusion of the corollary then follows. \square

Remark 4.14. In [1], Ahern obtained a version of Corollary 4.13 when $n = 1$ under a weaker assumption that F is merely bounded. We do not know whether this still holds true for $n \geq 2$.

5. OPEN QUESTIONS

In this section we list some questions that we have not been able to answer. We hope these questions will stimulate more research in this area.

Theorem 4.9 and its corollaries are restricted by the assumption that $n \leq 25$. This is due to the fact that we have not found a way to verify whether the polynomial Q has all roots outside of Ω_∞ for all n .

Question 1. Is it true that all roots of Q lie outside Ω_∞ for all $n \geq 2$?

Theorem 4.3 is restricted to Toeplitz operators with non-negative symbols. The well known Axler-Zheng theorem states that if T is a finite sum of finite products of Toeplitz operators with bounded symbols on A_1^2 such that $B(T)(z) \rightarrow 0$ as $|z| \uparrow 1$, then T is compact. (Recall that $B(T)$ is the Berezin transform of T as defined in Section 4.) Since the Berezin transform B is

no longer one-to-one on the space of bounded operators on A_n^2 when $n \geq 2$, it seems that the full version of Axler-Zheng theorem may not hold. On the other hand, the answer to the following question may be affirmative.

Question 2. Let f be bounded on \mathbb{D} . If $B(T_f)(z) \rightarrow 0$ as $|z| \uparrow 1$, does it follow that T_f is compact on A_n^2 for $n \geq 2$?

The general zero-product problem asks: if f and g are bounded functions such that $T_f T_g = 0$, does it follow that one of the functions must be zero? Corollary 4.10 answers this question in the affirmative when both functions are harmonic. In [3], Ahern and the first author also obtained the affirmative answer for operators on A_1^2 when one of the function is radial. In [11, 12], the second author generalized this result to more than two functions all of which, except possible one, are radial. While the general zero-product problem (even on A_1^2) is still far from its solution, an answer to the following question may be possible. (We note that Toeplitz operators with radial symbols are diagonal with respect to the standard orthonormal basis of A_1^2 . This, however, is not the case on A_n^2 with $n \geq 2$.)

Question 3. Let f and g be bounded functions, one of which is radial. If $T_f T_g = 0$ on A_n^2 (or more generally, $T_f T_g$ has finite rank), must one of these functions be zero?

The next question is related to Corollary 4.13. The answer was known to be affirmative on A_1^2 by Ahern [1].

Question 4. Suppose u and v are bounded harmonic functions and $T_u T_v = T_F$ on A_n^2 ($n \geq 2$), where F is a bounded function (without any assumption about smoothness). Does it imply that \bar{u} or v is analytic?

REFERENCES

- [1] Patrick Ahern, *On the range of the Berezin transform*, J. Funct. Anal. **215** (2004), no. 1, 206–216. MR 2085115 (2005e:47067)
- [2] Patrick Ahern and Željko Čučković, *A theorem of Brown-Halmos type for Bergman space Toeplitz operators*, J. Funct. Anal. **187** (2001), no. 1, 200–210. MR 1867348 (2002h:47040)
- [3] ———, *Some examples related to the Brown-Halmos theorem for the Bergman space*, Acta Sci. Math. (Szeged) **70** (2004), no. 1-2, 373–378. MR 2072710 (2005d:47046)
- [4] Patrick Ahern, Manuel Flores, and Walter Rudin, *An invariant volume-mean-value property*, J. Funct. Anal. **111** (1993), no. 2, 380–397. MR 1203459 (94b:31002)
- [5] Sheldon Axler and Željko Čučković, *Commuting Toeplitz operators with harmonic symbols*, Integral Equations Operator Theory **14** (1991), no. 1, 1–12. MR 1079815 (92f:47018)
- [6] Mark Benevich Balk, *Polyanalytic functions*, Mathematical Research, vol. 63, Akademie-Verlag, Berlin, 1991. MR 1184141 (93k:30076)
- [7] Boo Rim Choe, Hyungwoon Koo, and Young Joo Lee, *Sums of Toeplitz products with harmonic symbols*, Rev. Mat. Iberoam. **24** (2008), no. 1, 43–70. MR 2435966 (2009h:47052)
- [8] Željko Čučković, *Finite rank perturbations of Toeplitz operators*, Integral Equations Operator Theory **59** (2007), no. 3, 345–353. MR 2363013 (2008h:47056)

- [9] Kunyu Guo, Shunhua Sun, and Dechao Zheng, *Finite rank commutators and semi-commutators of Toeplitz operators with harmonic symbols*, Illinois J. Math. **51** (2007), no. 2, 583–596 (electronic). MR 2342676
- [10] A. D. Košelev, *The kernel function of a Hilbert space of functions that are polyanalytic in the disc*, Dokl. Akad. Nauk SSSR **232** (1977), no. 2, 277–279. MR 0427648 (55 #679)
- [11] Trieu Le, *Finite-rank products of Toeplitz operators in several complex variables*, Integral Equations Operator Theory **63** (2009), no. 4, 547–555.
- [12] ———, *A refined Luecking’s theorem and finite rank products of Toeplitz operators*, Complex Anal. Oper. Theory **4** (2010), no. 2, 391–399.
- [13] Miroslav Pavlović, *Decompositions of L^p and Hardy spaces of polyharmonic functions*, J. Math. Anal. Appl. **216** (1997), no. 2, 499–509. MR 1489593 (99b:46029)
- [14] A. K. Ramazanov, *On the structure of spaces of polyanalytic functions*, Mat. Zametki **72** (2002), no. 5, 750–764. MR 1963139 (2003m:30094)
- [15] ———, *An estimate for the norm of a polyanalytic function in terms of the norm of its polyharmonic component*, Mat. Zametki **75** (2004), no. 4, 608–613. MR 2068770 (2005f:30079)
- [16] N.V. Rao, *Range of Berezin transform*, preprint, arXiv:1003.3939.
- [17] Walter Rudin, *Function theory in the unit ball of \mathbf{C}^n* , Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science], vol. 241, Springer-Verlag, New York, 1980. MR 601594 (82i:32002)
- [18] Hartmut Wolf, *Toeplitz operators on polyanalytic functions and Klee’s combinatorial identity*, Math. Nachr. **166** (1994), 5–15. MR 1273319 (95m:47043)
- [19] Kehe Zhu, *Operator theory in function spaces*, second ed., Mathematical Surveys and Monographs, vol. 138, American Mathematical Society, Providence, RI, 2007. MR 2311536

DEPARTMENT OF MATHEMATICS, MAIL STOP 942, UNIVERSITY OF TOLEDO, TOLEDO, OH 43606

E-mail address: `zcuckovi@math.utoledo.edu`

DEPARTMENT OF MATHEMATICS, MAIL STOP 942, UNIVERSITY OF TOLEDO, TOLEDO, OH 43606

E-mail address: `trieu.le2@utoledo.edu`