

The KCAL VERA 22 GHz calibrator survey

L. Petrov¹

ADNET Systems Inc./NASA GSFC, Code 610.2, Greenbelt, 20771, USA

Leonid.Petrov@lpetrov.net

M. Honma² and S. M. Shibata²

National Astronomical Observatory of Japan, Mitaka, Japan

ABSTRACT

We observed at 22 GHz with the VLBI array VERA a sample of 1536 sources with correlated flux densities brighter than 200 mJy at 8 GHz. One half of target sources has been detected. The detection limit was around 200 mJy. We derived the correlated flux densities of 877 detected sources in three ranges of projected baseline lengths. The objective of these observations was to determine the suitability of given sources as phase calibrators for dual-beam and phase-referencing observations at high frequencies. Preliminary results indicate that the number of compact extragalactic sources at 22 GHz brighter than a given correlated flux density level is twice less than at 8 GHz.

Subject headings: Quasars and Active Galactic Nuclei

1. introduction

Currently, VLBI astrometry is the best tool to measure distances and motions of sources located at kpc scale and hence to explore structure of the Milky Way in the Galactic scale. For instance, Japanese VERA project (VLBI Exploration of Radio Astrometry; (Honma, Kawaguchi & Sasao 2000)) have been conducting astrometric monitoring of positions of Galactic maser sources with respect to reference compact extragalactic objects, yielding handful measurements of parallaxes and proper motions of maser sources (e.g., see recent PASJ special issue for VERA, such as Honma et al. (2011), Nagayama et al. (2011) and others). The VLBA is actively used for astrometry of Galactic maser sources (e.g., Reid et al. (2009)) and the European VLBI Network (EVN) conducts astrometric observations of methanol maser sources (e.g., Rygl et al. (2010)).

In order to measure parallaxes and proper motions of target sources at kpc scales, one has to achieve a positional uncertainty as small as $10\mu\text{as}$, which requires accurate phase calibration

based on phase-referencing. In conducting phase-referencing VLBI, nearby calibrator sources are necessary to trace and significantly reduce phase variations caused by tropospheric fluctuations. To do this effectively, calibrators must be located close to target sources, typically within $1\text{--}2^\circ$ separation. This requires a high density of calibrator sources in the sky, and hence, there is still a strong demand for finding many calibrator sources.

To date, there have been several massive surveys of compact calibrators such as VCS (VLBA Calibrator Surveys, Petrov et al. (2008) and references therein), LBA Calibrator Survey for the southern hemisphere (Petrov et al. 2011e), VIPS (Helmboldt et al. 2007, Petrov & Taylor 2011c), and several ongoing programs: the program of study the *Fermi* active galaxy nuclei (AGNs) at parsec scales¹ (Kovalev & Petrov (2011), paper in preparation), the program of observing radio-loud 2MASS galaxies² (Petrov et al. (2011), paper in preparation), the program of observing optically

¹<http://astrogeo.org/faps>

²<http://astrogeo.org/v2m>

bright quasars (Bourda et al. 2008, 2011, Petrov 2011a), and the recent VLBA calibrator search for the BeSSeL survey (Immer et al. 2011).

Together with regular geodetic VLBA observations of 1000 sources (the RDV program (Petrov et al. 2009)), by June 2011 positions of 6455 sources at a milliarcsecond level of accuracy were derived from analysis of these massive surveys. The sources turned out compact enough to be detected with VLBI, i.e. they have a core of mas scale, mostly with correlated flux larger than 70 mJy. However, these surveys were in most cases conducted in relatively low frequencies such as 2 (S-band), 5 (C-band) or 8 GHz (X-band), at which the telescope performance is the best. On the other hand, recent VLBI maser astrometry is often done at higher frequencies beyond 10 GHz. For instance, VERA’s main bands are 22 (K-band) and 43 GHz (Q-band) for H₂O and SiO maser sources. Maser astrometry with VLBA is mainly conducted at 12 GHz for CH₃OH masers and 22 GHz for H₂O masers. Therefore, calibrator information at high frequencies (such as K and higher bands) is of great importance for on-going and future astrometric observations. Compact calibrators which are cores of radio bright AGNs have a wide variety of their spectra: for the majority of sources the correlated flux densities falls with the frequency, although some sources may have spectra growing with frequency or peaked in a GHz regime. Hence, the extrapolation of the correlated flux densities from S and X band to 22 or 43 GHz is highly unreliable. Therefore, it is necessary to conduct a systematic survey of K-band flux densities for the compact sources which were already detected in S and X bands.

We have identified ~2000 sources previously observed with VLBI with $\delta > -30^\circ$ with correlated flux densities > 200 mJy at X-band at baselines longer than 900 km. According to the log N –log S curve, this sample is complete at the 95% level (Kovalev 2010, private communication). Of these sources, 511 have been previously observed in large K-band surveys: VERA Fringe Search Survey (Petrov et al. 2007), KQ survey (Lanyi et al. 2010), VLBI Galactic plane survey (Petrov et al. 2011d) and in the EVN Galactic plane survey (Petrov 2011b), and their correlated flux densities at 22 GHz have been measured. The brightness at the K-band of other objects was not known.

We conducted a dedicated survey of remaining 1536 sources at 22 GHz with VERA in the K-band Calibrator Survey (KCAL) campaign. The goal of these observations was to check their detectability at K-band and to measure the correlated flux densities of detected sources at baselines 1000–2000 km.

The first objective of this campaign was to provide a complete list of calibrators suitable for VERA observations of faint targets. According to our prior observations, the detection limit of the VERA network for 2 minutes of integration time is around 200 mJy, depending on weather conditions. Therefore, the list of sources observed in this and the previous K-band surveys is expected to approach the completeness at the 200 mJy level, provided the spectra of compact cores are flat or falling. According to Massardi et al. (2010) who analyzed simultaneous ATCA spectra at 4.8, 8.4 and 20 GHz, the share of sources with growing spectra that may be missed in our sample does not exceeded 8%.

The second objective of this campaign is to collect information for a population analysis of a large complete sample. In particular, the analysis of the dataset that combines existing and new data will help to answer the question what is the distribution of spectral indexes of the core regions and the source compactness at high frequencies, whether the spectral index at parsec scales is systematically different than the spectral index at kiloparsec scales, whether the compactness at K-band is systematically different than the compactness at X and S bands.

In this paper we present results of the survey. In section 2 we describe the observations, their design and scheduling. In section 3 we discuss analysis technique. The catalogue of correlated flux densities of detected sources accompanied with analysis of flux density uncertainties is presented in section 4 followed by concluding remarks that are given in section 5.

2. Observations

Observations were carried out at the VERA four-station network. The primary task of the array is to perform parallax measurement of maser sources. In order to maximize the throughput of the instrument, observing time for the KCAL ex-

periments was allotted in blocks that fill gaps between parallax measurement observing sessions or during periods of time when one of the antenna was under maintenance.

A monthly observing plan for VERA parallax measurements was usually finalized by at least one week before the beginning of the month. When there were suitable gaps for KCAL experiments and there were enough magnetic tapes in the Mitaka correlation center, we ran calibrator survey experiments during these gaps. The parallax measurement requires all four stations of VERA to achieve its astrometric accuracy. If any station, other than Ogasawara, could not join regular observations because of maintenance or instrumental problems, KCAL experiments also scheduled during that time with three stations. Ogasawara station was thought as indispensable station because the array are almost aligned as a line without Ogasawara station.

2.1. Scheduling

Scheduling software `sur_sked` selected sources from the pool of candidate objects in a sequence that minimizes slewing time. At a given experiment, each source was observed in one scan of 120 seconds long. Every 30 minutes a scan of a strong source with the brightness distribution map produced from VLBA observations under the KQ observing campaign (Lanyi et al. 2010) was inserted in the schedule. The purpose of including these scans in the schedule was to compare our measurements of the correlated flux densities of sources with known images considered as the ground truth in order to evaluate gain corrections. The target sources that were observed in one scan were returned in the pool for scheduling in the second scan in following experiments.

In total, 36 experiments were scheduled. However, six experiments were canceled for various reasons, in three observing sessions two stations either failed or did not observed and these experiments were excluded from analysis. The dates and durations of the 27 VLBI experiments under the KCAL program over the period 2007–2009 that were used in the analysis are shown in Table 1.

The scheduling goal was to have each target source observed in two scans. Due to the nature of scheduling in a fill-in mode, it turned out dif-

Table 1: Dates and durations of experiments. Only those experiments that were used in the final analysis are shown. Station abbreviations: Ir for Iriki, Is for Ishigakijima, Mz for Mizusawa, Og for Ogasawara.

Exp ID	Date	Dur (h)	Network
kcal_01	2007.05.28	5.3	Ir Is Mz Og
kcal_02	2007.05.30	5.8	Ir Is Mz Og
kcal_03	2007.05.31	3.9	Ir Mz Og
kcal_04	2007.08.24	3.8	Ir Is Mz Og
kcal_05	2007.08.24	14.3	Ir Is Mz Og
kcal_06	2007.08.25	6.8	Ir Is Mz Og
kcal_07	2007.11.18	4.8	Ir Is Mz Og
kcal_09	2007.11.23	4.5	Ir Is Mz Og
kcal_10	2007.12.10	5.8	Ir Is Mz Og
kcal_11	2007.12.12	2.6	Ir Mz Og
kcal_12	2007.12.12	2.8	Ir Mz Og
kcal_15	2007.12.19	5.8	Ir Is Mz Og
kcal_16	2007.12.20	3.8	Ir Is Mz Og
kcal_17	2007.12.21	3.8	Ir Is Og
kcal_18	2007.12.22	2.5	Ir Is Og
kcal_19	2007.12.22	3.8	Ir Is Mz Og
kcal_23	2008.02.29	6.8	Ir Is Mz Og
kcal_24	2008.06.03	2.1	Ir Is Mz Og
kcal_25	2008.06.11	5.2	Ir Is Mz Og
kcal_27	2008.10.06	15.8	Ir Is Mz Og
kcal_29	2008.10.12	3.6	Ir Is Mz Og
kcal_30	2008.11.11	2.3	Ir Is Mz Og
kcal_31	2008.11.16	2.1	Ir Is Mz Og
kcal_32	2008.11.14	4.0	Ir Is Mz Og
kcal_33a	2009.03.13	7.9	Ir Is Mz
kcal_33c	2009.03.21	3.9	Ir Is Mz
kcal_33d	2009.03.22	8.8	Ir Is Mz

ficult to reach this goal. As it seen from Table 2, 1/3 of the sources were observed in one scan. In total, 1536 target sources were observed for 143 hours. The antennas spent 71% time on target sources. Remaining time was spent for observing the amplitude calibrators and for slewing.

The left circular polarization in the 21.97–22.47 GHz band was received, sampled with 2 bit quantization, and filtered using the VERA digital filter (Iguchi et al. 2005) before being recorded onto magnetic tapes. The digital filter split the data within the 500 MHz band into 16 frequency channels of 16 MHz width each, equally spaced with 16 MHz wide gaps.

Table 2: Statistics of the number of scans per observed source. The first column shows the number of scans, the second table shows the number of target sources which had that number of scans. The last column shows the share of sources from the target list which had that number of scans.

# scans	# obs	Share
1	530	35%
2	633	41%
3	267	17%
4	102	5%
5	4	0.2%

3. Data analysis

3.1. Fringe fitting

The data were correlated on the Mitaka FX correlator (Chikada et al. 1991). Correlation output was written in the FITS-IDI format. Consecutive analysis was performed with computer program *PLMA*³. The procedure of data analysis is described in details in Petrov et al. (2011d). Here only a brief outline is given. After applying correction of fringe amplitude for digitization, the spectrum of the cross-correlation function was presented as a two-dimensional array with the first dimension running over frequency channels and the second dimension running over time. The two-dimensional Fourier-transform of the spectrum over frequency and time cast the spectrum of the cross-correlation function into the domain of group delay and phase delay rate. In the presence of the signal in the data, the Fourier-transform of the cross-spectrum exhibits a sequence of peaks. The amplitude of the major peak is proportional to the fringe amplitude of the signal. The fringe fitting process locates the peaks and determines the group delay, delay rate and fringe amplitude that correspond to the main maximum of the Fourier-transform of the cross-correlation spectrum.

In order to determine the detection threshold, we computed the ratios of fringe amplitudes to mean amplitudes of the Fourier-transform of the


cross-correlation spectrum. That mean amplitude was computed by averaging 32768 randomly selected samples of the cross-spectrum Fourier-transform after excluding the amplitudes that are greater than 3.5 times of the variance of amplitudes in the sample.

Even in the absence of the signal, the fringe fitting procedure will find a peak, but the amplitude of this peak will not be related to the fringe amplitude. In order to determine the detection threshold, we computed the histogram of the achieved signal-to-noise ratios SNR (SNR) in the KCAL experiments in the range of [3.8, 6.5] (see Figure 1) and fitted it with the theoretical curve $p(s)$ of the fringe amplitude distribution in the absence of the signal (Petrov et al. 2011d):

$$p(s) = \frac{2}{\pi} \frac{n_{\text{eff}}}{\sigma_{\text{eff}}} s e^{-\frac{s^2}{\pi}} \left(1 - e^{-\frac{s^2}{\pi}}\right)^{n_{\text{eff}}-1}, \quad (1)$$

where n_{eff} is the effective number of independent samples and σ_{eff} is the effective noise variance.

Fig. 1.— The left tail of the empirical distribution of the achieved SNR from results of fringe fitting VERA data (filled circles) and the fitted curve (thin line) of the theoretical distribution for the case of no signal.



After determining n_{eff} and σ_{eff} , we can find the probability of false detection at a given SNR by integrating expression 1 over s , which can be easily done analytically. Specifically, we found that the probability of false detection is less than 0.001 when the SNR > 6.03. We considered a source as detected if the SNR in at least two observations at different baselines of the same scan was above the detection limit 6.03. In the absence of the signal, the probability of finding two peaks exceeding the

³Available at <http://astrogeo.org/pima>

threshold limit in data of different observations is in the range of 10^{-3} to 10^{-6} depending whether the errors are completely correlated or completely uncorrelated. In practical terms, this means that our catalogue may have no more than one or two falsely detected objects.

3.2. Amplitude calibration

System temperatures including atmospheric attenuation were measured with the chopper-wheel method (Ulich & Haas 1976). At the beginning of each scan a microwave absorber at ambient temperature was inserted just in front of the feed horn and the received total power was measured with a power meter. Using the measured total power for the blank sky and the absorber, the temperature scale automatically corrected for the atmospheric attenuation was determined. We estimate the uncertainty in the temperature scale around 10%, mainly due to the assumption that the ambient temperature is the same as the air temperature.

The initial amplitude calibration was made by scaling fringe amplitudes determined with the fringe fitting process by measured system temperature and dividing it by the antenna gain. Then antenna gains were adjusted by using results of comparison of the calibrated fringe amplitude of observed calibrator sources with the correlated flux densities predicated on the basis of K-band brightness distributions⁴ produced from analysis of observations from the KQ (Lanyi et al. 2010) and VLBA Galactic Plane Surveys (VGaPS) (Petrov et al. 2011d):

$$F_{corr} = \left| \sum_j c_j(x, y) e^{\frac{2\pi i f}{c}(u x + v y)} \right|, \quad (2)$$

where c_i is the correlated flux density of the j th CLEAN component with coordinates x and y with respect to the center of the image; u and v are the projections of the baseline vectors onto the tangential plane of the source.

Then we built a system of equations for all observations of calibrators:

$$F_{corr} = \sqrt{g_i g_j} A_{corr} \quad (3)$$

that relates the calibrated amplitude A_{corr} , gain corrections g for stations i and j of the baseline,

and the predicted correlated flux of the amplitude calibrator. After taking logarithms from left and right hand sides, we solved for average gains corrections for all stations using least squares (LSQ). An iterative procedure of outliers elimination was performed: we searched for an observation with the maximum by module logarithm of the ratio of observed and predicted correlated flux density and removed it from consecutive computations. The LSQ solution was updated and the process was repeated till the largest by module outlier was under 3.5σ .

The number of calibrators in each individual experiment varied. On average, 9 calibrators were used for gain correction adjustment in each experiment. If the model brightness distributions were perfect, and gain corrections were stable over an experiment, calibration errors would have been below the noise level. Several factors degrades the quality of calibration using this method. First, the images of calibrator sources were produced using observations at different sampling of spatial frequencies than the analyzed observations. Computation of the predicted correlated flux densities is equivalent to an interpolation of visibilities measured in KQ and VGaPS VLBA projects to u and v baseline projections in the KCAL experiments. Errors of this interpolation may be significant, except for sources with very simple structure. Second, both source structure and the peak brightness evolve with time. Since the time difference in epochs between KQ, VGaPS and KCAL experiments is 2–6 years, the changes in source brightness distribution may be significant. The sampling bias and the source variability are expected to cause only random errors in gain, but not a systematic bias. Some calibrator sources may become brighter, some dimmer, but the average flux density of the ensemble should be rather stable. Third, we assumed that gain corrections are constant over time of an individual experiment since we do not have enough information for modeling their time variability.

4. The correlator flux density catalogue

Since the data are too sparse to produce meaningful images, we computed average correlated flux densities for detected sources in three ranges of the projected baseline lengths: 0–70 megawave-

⁴Available at <http://astrogeo.org>

lengths, 70–100 megawavelengths, and 100–250 megawavelengths, which corresponds to lengths 0–955 km, 955–1365 km and longer than 1365 km respectively. The amplitudes were calibrated for gain corrections using the method described in the previous section. This simplified method of correlated flux density evaluation is an alternative to a rigorous imaging procedure in the case when there are too few measurements.

The catalogue of estimates of correlated flux densities of 877 observed sources, including 750 targets and 127 calibrators, is presented in table 3. Objects with at least two detections are put in the catalogue. Columns 1 and 2 show IAU and IVS source names. Column 3 shows the source status: C stands for an amplitude calibrator, blank stands for a target object. Column 4 shows the number of experiments in which a source was detected and column 5 shows the total number of detections. Columns 6, 7 and 8 present the estimates of the average correlated flux density in three ranges of the projected baseline lengths. Columns 9, 10, and 11 show the estimates of the correlated flux density uncertainty: $\sigma(F_{\text{corr}}) = A_{\text{corr}} \cdot \sqrt{0.2^2 + \frac{2}{\pi} \frac{1}{\text{SNR}^2}}$. Columns 12 and 13 show right ascensions and declinations.

Of 1536 observed sources, including both targets and calibrators, 407 were not detected at all and 252 were detected only in one observation. The detections from the latter group were considered unreliable and were not included in the catalogue.

4.1. Error analysis

Errors in correlated flux density estimates are due to 1) the thermal noise in estimates of fringe amplitude; 2) the uncertainties in system temperature measurements; 3) the uncertainties in antenna gain measurement; 4) the sampling bias in a predicted correlated flux density of calibrators; 5) the variability of calibrator sources.

The uncertainty due to the thermal noise can be easily evaluated as $\sqrt{2/\pi} \langle a_n \rangle / a$, where $\langle a_n \rangle$ is the average amplitude of the noise, and a is the fringe amplitude. As we already mentioned, the uncertainty in system temperature measurement is around 10%. The aperture efficiency of VERA antenna is measured every year and known within

10% accuracy (see VERA status report⁵). Since these two uncertainties are assumed uncorrelated, these two factors would introduce an uncertainty of the a priori gain calibration at $\sim 14\%$ level.

Since on average, nine amplitude calibrators were used for gain adjustments, this redundancy can be exploited for evaluation the gain correction uncertainties. We computed the average and the root mean square (rms) of the residual mismatches between observed correlated flux density of calibrators after applying gain corrections A_{corr}^p from the LSQ fit and the predicted correlated flux densities from the brightness distributions:

$$\begin{aligned} \text{Avr} &= \left(\prod_i \frac{F_{\text{corr},i} \sqrt{g_1 g_2}}{A_{\text{corr},i}^p} \right)^{1/n} \\ \text{Rms} &= \sqrt{\frac{\sum_i \left(\frac{F_{\text{corr},i} \sqrt{g_1 g_2}}{A_{\text{corr},i}^p} - 1 \right)^2}{n}}. \end{aligned} \quad (4)$$

We found $\text{Avr} = 0.994$ and $\text{Rms} = 0.21$. The first statistics describes the systematic bias and the second statistics is the measure of the contribution of uncertainties in gain adjustments on the uncertainty of our estimate of the correlated flux density.

In order to evaluate the representativeness of this statistics, we computed the median correlated flux densities in three ranges of projected baseline lengths of two experiments of the 24 GHz VLBA VGaPS campaign using two methods: 1) rigorous self-calibration imaging and 2) the same simplified method used for processing KCAL experiments. In order to closely mimic analysis of the KCAL experiments, we used for our tests the brightness distributions from the KQ campaign made at epochs at least one year prior to observations. We got $\text{Avr} = 0.996$ and $\text{Rms} = 0.24$. Then we computed the rms of the scatter of the ratios of the correlated flux density F_{corr}^s determined by the simplified method to the flux density F_{corr}^r determined by the rigorous method: $\text{rms} = \sqrt{\sum_i (F_{\text{corr},i}^s / F_{\text{corr},i}^r - 1)^2}$. We found the rms equal to 0.15. Considering the brightness distributions from the self-calibration analysis procedure as the ground truth, we conclude that the accuracy

⁵Available at <http://veraserver.mtk.nao.ac.jp/>

TABLE 3

THE CORRELATED FLUX DENSITIES OF 877 SOURCES THAT HAVE AT LEAST TWO DETECTIONS IN VERA KCAL OBSERVING CAMPAIGN.

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
IAU name	IVS name	flag	#Exp	#Det	$F_{<70}$ Jy	F_{70-100} Jy	$F_{>100}$ Jy	$E_{<70}$ Jy	E_{70-100} Jy	$E_{>100}$ Jy	Right ascen h m s	Declination ° ' "
J0001+1914	2358+189		1	4	0.221	0.322	0.216	0.055	0.070	0.051	00 01 08.62	+19 14 33.8
J0005+3820	0003+380		2	8	-1.000	0.608	0.526	-1.000	0.136	0.116	00 05 57.17	+38 20 15.1
J0006-0623	0003-066		2	9	1.027	1.120	1.212	0.104	0.205	0.176	00 06 13.89	-06 23 35.3
J0008+6837	0005+683		1	2	-1.000	0.353	-1.000	-1.000	0.080	-1.000	00 08 33.47	+68 37 22.0
J0010+1058	IIIZW2	C	2	6	-1.000	1.193	1.442	-1.000	0.239	0.289	00 10 31.00	+10 58 29.5
J0010+1724	0007+171		1	2	-1.000	0.340	0.266	-1.000	0.078	0.060	00 10 33.99	+17 24 18.7
J0010-2157	0008-222		1	2	-1.000	0.236	-1.000	-1.000	0.052	-1.000	00 10 53.64	-21 57 04.2
J0011+7045	0008+704		2	8	0.440	0.405	0.542	0.043	0.088	0.064	00 11 31.90	+70 45 31.6
J0012-3954	0010-401		1	3	0.505	0.494	-1.000	0.120	0.103	-1.000	00 12 59.90	-39 54 26.0
J0013+4051	0010+405		2	7	0.531	0.534	0.492	0.065	0.119	0.115	00 13 31.13	+40 51 37.1
J0013-0423	0011-046		2	7	0.552	0.406	0.445	0.071	0.094	0.101	00 13 54.13	-04 23 52.2
J0017+5312	0015+529		1	2	-1.000	0.270	-1.000	-1.000	0.067	-1.000	00 17 51.75	+53 12 19.1

NOTE.—Table 3 is presented in its entirety in the electronic edition of the Publications of the Astronomical Society of the Pacific. A portion is shown here for guidance regarding its form and contents.


of the median correlated flux density obtained by the simplified method is at a level of 15% for the VGaPS campaign. Thus, the Rms statistics give us rather an upper limit of gain errors.

Another way to evaluate the uncertainty of correlated flux densities is to compute the rms of the scatter of ratios of the correlated flux densities of all the KCAL sources which have three or more observations. We got the value of the rms 0.20, which is close to the Rms statistics. Therefore, we conclude that the average uncertainty of calibration error is 20%. Since the uncertainty in fringe amplitude caused by the thermal noise and calibration errors are independent, we compute the multiplicative uncertainty of reported correlated flux density as a sum of these two contributions in quadrature: 0.2 and $\sqrt{\frac{2}{\pi}} \frac{1}{\text{SNR}}$.

5. Summary

We observed with VERA at 22 GHz a subset of the complete sample of continuum compact extragalactic sources with correlated flux densities > 200 mJy at X-band at declinations $> -30^\circ$. The subset excluded the sources previously detected at K-band at large VLBA and VERA surveys. Of 1536 target sources, approximately one half has been detected. The errors of the correlated flux densities are a level of 20%.

Fig. 2.— The distribution of the correlated flux densities at baseline projection lengths longer than 100 megawavelengths. The last bin of the histogram has all the sources with correlated flux density > 2 Jy.



The distribution of the KCAL correlated flux densities (Figure 2) suggests that the catalogue is incomplete at flux densities less than 200 mJy because weaker sources are not reliably detected with VERA. This result is in agreement with our analysis of previous VERA observations (Petrov et al. 2007) where we estimated the probability of detection of a source with the correlated flux density 200 mJy at a level of 70%.

We reserve a rigorous population analysis to a

future publication. Preliminary results indicate that the number of compact extragalactic sources at K-band brighter than a given correlated flux density level is twice less than at the X-band.

We would like to thank Alan Fey for making publicly available not only contour plots of images from the KQ survey, but brightness distribution and calibrated flux densities in the FITS-format. The availability of this information was crucial for our project.

REFERENCES

- Bourda, G., Charlot, P., & Le Campion, J.-F. 2008, *A&A*, 490, 403
- Bourda, G., Collioud, A., Charlot, P., Porcas, R. W., & Garrington, S. T. 2011, *A&A*, 526, A102
- Chikada, Y., et al. 1991, in *Frontiers of VLBI*, ed. H. Hirabayashi, M. Inoue, & H. Kobayashi, Tokyo: Universal Academy Press, 79
- Helmboldt, J. F., Taylor, G. B., Tremblay, S., Fassnacht, C. D., Walker, R. C., Myers, S. T., Sjouwerman, L. O., Pearson, T. J., Readhead, A. C. S., Weintraub, L., Gehrels, N., Romani, R. W., Healey, S., Michelson, P. F., Blandford, R. D., Cotter, G. 2007, *ApJ*, 658, 203
- Honma, M., Kawaguchi, N., Sasao, T. 2000, in *Proc. SPIE*, 4015, Radio Telescope, ed. H. R. Buthcer, 624
- Honma, M., Hirota, T., Kan-Ya, Y., Kawaguchi, N., Kobayashi, H., Kurayama, T., Sato, K. 2011, *PASJ*, 63, 17
- Iguchi, S., Kurayama, T., Kawaguchi, N., and Kawakami, K. 2005, *PASJ*, 57, 259
- Immer, K. et al., 2011, *ApJS*, 194, 25
- Lanyi G. E. et al. 2010, *AJ*, 139, 1695
- Massardi M. et al., 2011, *MNRAS*, 412, 318
- Nagayama, T., Omodaka, T., Nakagawa, A., Handa, T., Honma, M., Kobayashi, H., Kawaguchi, N., Miyaji, T. 2011, *PASJ*, 63, 23
- Petrov L., Hirota T., Honma M., Shibata S. M., Jike T., Kobayashi H. 2007, *AJ*, 133, 2487
- Petrov L., Kovalev Y. Y., Fomalont E., Gordon D., 2008, *AJ*, 136, 580
- Petrov L., Gordon D., Gipson J., MacMillan D., Ma C., Fomalont E., Walker R. C., Carabjal C., 2009, *Jour. Geodesy*, 83(9), 859
- Petrov L. 2011a, *AJ*, submitted, preprint (<http://arxiv.org/abs/1103.2840>)
- Petrov L., 2011b, *MNRAS*, submitted, preprint (<http://arxiv.org/abs/1106.4883>)
- Petrov, L., Taylor, G. B., submitted to *AJ*, 2011c, preprint (<http://arxiv.org/abs/1106.2382>)
- Petrov L., Kovalev Y. Y., Fomalont E., Gordon D., 2011d, *AJ*, 142:35(23pp)
- Petrov L., Phillips, C., Bertarini, A., Murphy, T., Sadler, E. M., 2011e, *MNRAS*, 414(3), 2528
- Reid, M. J., Menten, K. M., Zheng, X. W., Brunthaler, A., Moscadelli, L., Xu, Y., Zhang, B., Sato, M., Honma, M., Hirota, T., Hachisuka, K., Choi, Y. K., Moellenbrock, G. A., Bartkiewicz, A. 2009, *ApJ*, 700, 137
- Rygl, K. L. J., Brunthaler, A., Reid, M. J., Menten, K. M., van Langevelde, H. J., & Xu, Y. 2010, *A&A*, 511, A2
- Ulich, B. L., & Haas, R. W. 1976, *ApJS*, 30, 247