

## A MODEL FOR THE MECHANISM OF Fe<sup>3+</sup> TO Fe<sup>2+</sup> REDUCTION IN DIOCTAHEDRAL SMECTITES

V.A. DRITS<sup>1,2</sup> AND A. MANCEAU<sup>1</sup>

<sup>1</sup> Environmental Geochemistry Group, LGIT-IRIGM, University Joseph Fourier and CNRS, 38041 Grenoble Cedex 9, France

<sup>2</sup> Geological Institute of the Russian Academy of Sciences, 7 Pyzhevsky Street, 109017 Moscow, Russia

**Abstract**—A model to compensate the 2:1 layer having excess negative charge owing to the reduction of Fe<sup>3+</sup> to Fe<sup>2+</sup> by sodium dithionite buffered with citrate-bicarbonate in nontronite, beidellite, and montmorillonite is proposed. This model is based on reassessing published experimental data for Fe-containing smectites and on a recently published structural model for reduced Garfield nontronite. In the reduced state, Fe<sup>2+</sup> cations remain six-fold coordinated, and increases of negative charge in the 2:1 layer are compensated by the sorption of Na<sup>+</sup> and H<sup>+</sup> from solution. Some of the incorporated protons react with structural OH groups to cause dehydroxylation. Also, some protons bond with undersaturated oxygen atoms of the octahedral sheet. The amount of Na<sup>+</sup> ( $p$ ) and H<sup>+</sup> ( $n_i$ ) cations incorporated in the structure as a function of the amount of Fe reduction can be described quantitatively by two equations:  $p = m/(1 + K_0 m_{\text{rel}})$  and  $n_i = K_0 m m_{\text{rel}}/(1 + K_0 m_{\text{rel}})$ ; with  $K_0 = \text{CEC} (9.32 - 1.06 m_{\text{tot}} + 0.02 m_{\text{tot}}^2)$ , where  $m_{\text{tot}}$  is the total Fe content in the smectite,  $m$  is the Fe<sup>2+</sup> content,  $m_{\text{rel}}$  is the reduction level ( $m/m_{\text{tot}}$ ), CEC is the cation-exchange capacity, and  $K_0$  is a constant specific to the smectite. The model can predict, from the chemical composition of a smectite, the modifications of its properties as a function of reduction level. Based on this model, the structural mechanism of Fe<sup>3+</sup> reduction in montmorillonite differs from that determined in nontronite and beidellite owing to differences in the distribution of cations over *trans*- and *cis*-octahedral sites.

**Key Words**—CMS Source Clay SWa-1, Dioctahedral Smectite, Iron Reduction, Structural Formula.

### INTRODUCTION

One of the most important factors governing the geochemical behavior of Fe in soils and sediments is the ability of Fe to change oxidation state. Therefore, oxido-reduction reactions of layer silicates often play an important role in environmental processes including weathering, microbial activity, and diagenetic transformations (Egashira and Ohtsubo, 1983; Stucki *et al.*, 1987, 1996). Redox reactions modify the chemical and physical properties of Fe-containing smectites, such as cation-exchange capacity (CEC), specific surface area, swelling behavior, and ability to fix interlayer cations (Khaled and Stucki, 1991; Lear and Stucki, 1985, 1989; Stucki *et al.*, 1984, 1996). Consequently, the texture (Gates *et al.*, 1998; Stucki and Tessier, 1991), permeability (Shen *et al.*, 1992), and fertility of formations in which Fe-containing smectites are present (Stucki, 1988) are affected. The potential for modification of smectite properties for industrial, engineering, and environmental applications by controlling the oxidation state of Fe is great (Chen *et al.*, 1987; Low *et al.*, 1983). The knowledge of the reduction mechanism of Fe<sup>3+</sup> to Fe<sup>2+</sup> is, therefore, essential for understanding a number of natural weathering processes as well as for purposely modifying specific properties of smectite (Ernstsen, 1998). This subject has been extensively studied in the past two decades, and also largely reviewed in the literature (Heller-Kallai, 1997; Rozenson and Heller-Kallai, 1976; Russell *et al.*,

1979; Stucki, 1988; Stucki *et al.*, 1976, 1996; Stucki and Roth, 1976, 1977).

Reliable results describing the reduction of Fe were obtained in dioctahedral smectites with Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> in a citrate-bicarbonate medium. Although parameters controlling Fe<sup>3+</sup>–Fe<sup>2+</sup> redox reactions in smectites were identified, an atomic-level understanding of these transformations is lacking, essentially because the crystal structure of reduced smectites was unknown until recently. An atomic-level understanding is crucial because revealing structural changes, which occur during the reduction of ferric iron, may provide insight, not only on the mechanism of the Fe<sup>3+</sup> reduction, but also how structural Fe<sup>2+</sup> modifies structural and surface properties of clays. Progress towards describing the structure of reduced smectites was achieved recently by Manceau *et al.* (2000), who investigated by polarized extended X-ray absorption fine structure (P-EXAFS) spectroscopy and powder X-ray diffraction (XRD) the modifications of the local structure of Fe atoms in reduced nontronite from Garfield, Washington. They proposed a model for the reduction mechanism of Fe<sup>3+</sup> to Fe<sup>2+</sup>.

Published models on redox reactions in Fe<sup>3+</sup>-containing dioctahedral smectites are first examined here. It is shown that they fail to explain several important experimental results, specifically, the dependence of the layer charge on the Fe<sup>2+</sup> content in reduced smectites. Stucki (1988) noted that “the source of this discrepancy is intriguing and may offer clues for the spe-




Figure 1. Structural transformations corresponding to the  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$  reduction mechanism proposed by Stucki and Roth (1977) in Equation (2).


cific reaction involved in the redox processes". A model describing the structural modifications of dioctahedral smectites as a function of  $\text{Fe}^{2+}$  content is proposed, and the capacity to explain experimental data is demonstrated. This model can be used to calculate, at least semi-quantitatively, the cationic and anionic composition of dioctahedral smectites for any amount of  $\text{Fe}^{3+}$  reduction.

#### MODELS FOR THE REDUCTION MECHANISM OF $\text{Fe}^{3+}$ TO $\text{Fe}^{2+}$

Solution-chemistry experiments have shown that the reduction mechanism of iron-containing dioctahedral smectites depends partly on the reducing agent, and on the physico-chemical characteristics of the solution where the redox reaction takes place (Stucki, 1988). For this reason, the following descriptions of previously proposed mechanistic models for the reduction of  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$  in dioctahedral ferruginous smectites is restricted to those involving sodium-dithionite reductant buffered with citrate-bicarbonate solution (CBD). This review shows those physico-chemical problems that have been solved and those that warrant further examination.

##### *The model of Roth and Tullock (1973)*

According to the model of Roth and Tullock (1973), the reducing agent transfers an electron to structural  $\text{Fe}^{3+}$  to produce  $\text{Fe}^{2+}$ . This process occurs with dehydroxylation of the octahedral sheets, followed by protonation of oxygen atoms that are undersaturated. The sequence of these reactions is described as follows:




where subscripts c and s denote species from the crystal structure and from solution, respectively. Water molecules formed by dehydroxylation migrate from

the clay framework, and the residual oxygen atoms incorporate protons from solution to form OH groups. This mechanism accounts for the decrease of structural OH groups observed experimentally during the reduction of  $\text{Fe}^{3+}$  and implies that ferrous atoms have five-fold coordination. Equation (1) indicates also that the layer charge should be invariant with the amount of iron reduction.

##### *The model of Stucki and Roth (1977)*

Measurements obtained for the first time under controlled atmospheric conditions showed that the layer charge of reduced smectites increased with  $\text{Fe}^{2+}$  content, in contrast to the prediction of the model of Roth and Tullock. To explain this result, Stucki and Roth (1977) proposed the following reaction mechanism:



where  $m$  and  $n$  are stoichiometric parameters. Equation (2) suggests that dehydroxylation occurs by a coalescence of edge-sharing OH groups during reduction to form  $\text{H}_2\text{O}$  molecules that diffuse from the structure, leaving behind oxygen ions. These oxygen ions are subsequently reprotated by hydrogen ions from solution. The stoichiometry coefficient,  $n$ , refers to sorbed  $\text{H}^+$  which remains in the clay structure after reduction. A schematic representation of the reaction mechanism corresponding to this model is shown in Figure 1. The main difference between Equations (1) and (2) is that the layer charge in the model of Stucki and Roth depends on the number,  $n$ , of OH groups that undergo dehydroxylation. By using tritium as a label element, Lear and Stucki (1985) showed that the initial loss of structural protons and subsequent sorption of protons from the solution to structural positions occurred near the ratio of 2:1. This result is in agreement with Equation (2) and with the structural model of Figure 1 because the loss of  $2\text{H}^+$  from the structure is partly balanced by the sorption of one  $\text{H}^+$  from the solution. The agreement between theory and experiment was considered by Stucki and Roth as strong support in favor of this mechanism. However, according to Equation (2) and Figure 1,  $\text{Fe}^{2+}$  ions are in five-fold coordination in reduced dioctahedral ferruginous smectites. This model was recently discounted by Manceau *et al.* (2000), who showed by P-EXAFS and XRD that  $\text{Fe}^{2+}$  cations remain in six-fold coordination in the reduced state.


From titration experiments on nontronite (Clay Minerals Society Source Clay, SWa-1), Lear and Stucki (1985) determined that the  $n/m$  ratio of 0.32 was invariant over the range of  $\text{Fe}^{2+}$  content explored. Based on this result, the following relationship between the layer charge,  $w$ , and the  $\text{Fe}^{2+}$  content was calculated:

$$w = w_0 + Km(1 - n/m) = w_0 + 0.68Km \quad (3)$$

where  $w_0$  is the layer charge of the oxidized sample,  $m$  is the Fe<sup>2+</sup> content, and  $K$  is a constant, the value of which depends on the units used for the terms in Equation (3). If  $w$  and  $w_0$  are expressed in meq/g, and  $m$  in mmole/g of clay, then  $K = 1$  meq/mmole. Stucki *et al.* (1984) and Lear and Stucki (1985) measured the dependence of layer charge on the amount of reduced iron present. The layer charge was estimated from the total Na content in reduced samples. Figure 2 shows that the linear relationship between  $w$  and  $n$  based on titration results (dotted lines) differs substantially from experimental data (curvilinear solid lines). Since theory [Equations (2) and (3)] and experiment disagree, this model should be reconsidered.

#### The model of Gan *et al.* (1992)

Electron spin resonance (ESR) spectroscopy was used to determine the reductant species responsible for the transfer of electrons during reduction. In solution, sodium dithionite (Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub>) dissociates to Na<sup>+</sup> and (S<sub>2</sub>O<sub>4</sub>)<sup>2-</sup> ions. The disulfite anion also dissociates in water, partly according to 2(S<sub>2</sub>O<sub>4</sub>)<sup>2-</sup> + H<sub>2</sub>O → (S<sub>2</sub>O<sub>3</sub>)<sup>2-</sup> + 2(HSO<sub>3</sub>)<sup>-</sup> → (S<sub>2</sub>O<sub>3</sub>)<sup>2-</sup> + 2(SO<sub>3</sub>)<sup>2-</sup> + 2H<sup>+</sup>, and partly into the paramagnetic free radical species, SO<sub>2</sub><sup>-•</sup>, which contains unpaired electrons as indicated here by “•” (Dunitz, 1956; Lynn *et al.*, 1964; Rinker *et al.*, 1958). Gan *et al.* (1992) showed that in Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> solutions, these free radicals are responsible for the reduction of structural Fe<sup>3+</sup>. They observed a correlation between the concentration of SO<sub>2</sub><sup>-•</sup> in solution and the reducing ability of the reagent. In particular, the reductive capacity of the Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> solution decreased with time as a result of the progressive diminution of the number of free radicals in the Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> solution. However, surprisingly, the addition of Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> to a nontronite suspension did not decrease the free-radical content, and the concentration of paramagnetic species was preserved or enhanced. To explain this result, Gan *et al.* (1992) assumed that, at a certain stage of Fe<sup>3+</sup> reduction, unpaired electrons were produced within the nontronite structure, and that Al sites may serve as sources of these unpaired electrons. Gan *et al.* (1992) emphasized that the migration of the internal high-potential electrons to structural Fe<sup>3+</sup> occurs only after sufficient Fe<sup>3+</sup> is reduced owing to the incorporation of electrons from the Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> solution. Based on these results, Gan *et al.* (1992) and Stucki *et al.* (1996) adapted the reaction sequence in (2) to:



where Z is an unidentified electron donor of electrical


charge  $-q$  located within the nontronite structure, which may be Al<sup>3+</sup> substituted for Si<sup>4+</sup> in tetrahedral sites; e<sup>-</sup> represents the reducing agent SO<sub>2</sub><sup>-•</sup> in the Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> solution; and  $w$ ,  $m$ ,  $n$ , and  $\alpha$  are stoichiometry coefficients. Equation (4) can not be used to simulate experimental observations because the value of  $\alpha$  is unknown.

Gan *et al.* (1992) and Stucki *et al.* (1996) proposed a conceptual model where they assumed the existence of a relationship between the energy of reducing electrons, the structural Fe<sup>2+</sup> content, and the progression of the reduction process. Three steps were distinguished. At first, reducing agents with modest reductive capability (RC) act on a small number of Fe<sup>3+</sup>, and the charge deficit in the reduced structure provokes a concomitant increase of layer charge. Then, the reduction of Fe<sup>3+</sup> by SO<sub>2</sub><sup>-•</sup> free radicals with higher RC results in the partial dehydroxylation of the nontronite structure. Finally, corresponding to the highest reducing electron energy of free radicals, internal electrons of the nontronite structure become delocalized, thereby increasing further the rate of reduction.


Thus, according to this model, the reduction mechanism of Fe<sup>3+</sup> in dioctahedral ferruginous smectites includes two predominant processes. The first is the reduction of Fe<sup>3+</sup> by electrons produced by free radicals. The second process is the initial elimination of structural OH and eventual incorporation of protons from solution to compensate, *pro parte*, the negative layer charge resulting from further reduction of iron. This model requires also a five-fold coordination for reduced Fe and an internal source of electrons with a high reducing energy. In addition, this model does not account for the measured variation of the layer charge with Fe reduction (Figure 2).

#### PROPOSED MODEL

Let us consider a nontronite model with the ideal structural formula:



The increase of negative charge of the layer resulting from the reduction of Fe<sup>3+</sup> is compensated two ways: (1) by an increase in the amount of positive interlayer charge through the adsorption of Na from Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> in solution (Stucki *et al.*, 1984), and (2) by a decrease in total negative charge of the anionic framework of the 2:1 layer through a loss of structural hydroxyl groups (Roth and Tullock, 1973). Both processes may occur simultaneously. Based on these observations, the reduction of Fe<sup>3+</sup> proceeds as:



and is accompanied by:







Figure 2. Relationships between the experimental and theoretical surface charge and the fraction of reduced Fe<sup>3+</sup> in various dioctahedral smectite samples. Observed data are from Stucki *et al.* (1984) and Lear and Stucki (1985), calculated data are from Equation (9). Dotted lines correspond to Equation (3) and curvilinear solid lines to Equation (10).



$$m = p + n_i \quad (6d)$$

where  $m$ ,  $n_i$ , and  $p$  are stoichiometric parameters, and  $w$  and  $w_0$  correspond to interlayer Na content in the reduced and unreduced state, respectively. Equation (6a) corresponds to the reduction step by electrons produced by free radicals. Equations (6b) and (6c) represent changes in composition of the interlayer and anionic framework. Equation (6d) indicates that the charge deficit created by the Fe<sup>3+</sup> to Fe<sup>2+</sup> reduction is balanced by the sorption of Na<sup>+</sup> and H<sup>+</sup>. Note that Equations (6a)–(6d) are based on the structural model proposed by Manceau *et al.* (2000) for reduced nontronite. According to this model, each pair of edge-forming OH groups coordinated to reduced Fe<sup>2+</sup> ions receives two protons, thus forming two H<sub>2</sub>O molecules, which diffuse from the structure to the solution. This reaction is accompanied by the migration of Fe<sup>2+</sup> ions from *cis*-sites to nearest *trans*-sites along the [010] direction. In accordance with this model, the stoichiometric coefficient  $n_i$  refers to the amount of sorbed H<sup>+</sup> and of structural OH groups, which both leave the structure after protonation. Unfortunately, the concentration of structural OH groups in reduced smectites is unknown, and the bonding mechanism of the incorporated H<sup>+</sup> cannot be easily derived (discussed in more detailed below). Initially, we assume that all incorporated H<sup>+</sup> ions are consumed in the process of dehydroxylation of the 2:1 layer (6c). Based on this model, the structural formula (5) is rewritten as a function of the amount of iron reduction (“reduction level”):



To evaluate if this model can account for the observed variation of the layer charge of Figure 2, the relationship between  $m$ ,  $n_i$ , and  $p$  must be obtained.

Based on crystal chemical considerations, the relative values of  $p$  and  $n_i$  are assumed to vary with the reduction level. The reduction of small amounts of ferrous iron will not appreciably disturb the crystal-lattice stabilization energy because dioctahedral smectites can accommodate a limited amount of divalent octahedral species in their structure (Güven, 1991). Therefore, when the content of Fe<sup>2+</sup> is relatively low, the layer charge is compensated predominantly by the sorption of Na. In this case,  $p > n_i$ . When the reduction level increases, the proportion of Na sorbed from solution will decrease progressively, and the crystal-lattice energy will stabilize by the hydration of structural OH groups. Thus, at a certain stage of the reduction process,  $n_i > p$ . The compensation of the layer charge as a function of the reduction level can thus be written:

$$n_i/p = K_0 m/m_{\text{tot}} = K_0 m_{\text{rel}} \quad (8)$$

where  $m$  and  $m_{\text{tot}}$  are the amount of reduced Fe<sup>3+</sup>, and the total amount of Fe per unit cell, respectively,  $m_{\text{rel}}$  equals the reduction level ( $m_{\text{rel}} \leq 1$ ),  $K_0$  is a sample-dependent constant ( $K_0 > 0$ ). The physical meaning of  $K_0$  is discussed below. Equation (8) shows that when  $K_0 m_{\text{rel}} < 1$ , then  $p > n_i$ , and *vice versa*. Combining of Equations (6d) and (8) leads to:

$$p = m/(1 + K_0 m_{\text{rel}}) \quad (9a)$$

$$n_i = K_0 m m_{\text{rel}}/(1 + K_0 m_{\text{rel}}). \quad (9b)$$

Because  $p$  and  $n_i$  depend only on  $m$  and  $m_{\text{tot}}$ , Equations (9a) and (9b) can be applied to any Fe<sup>3+</sup>-containing dioctahedral smectite. Indeed, a structural formula of a partially reduced dioctahedral smectite can be written:  $\text{Na}_{a_x+z+p}(\text{Si}_{8-x}\text{Al}_x)(\text{Al}_{4-m_{\text{tot}}-z}\text{Fe}^{3+}_{m_{\text{tot}}-m}\text{Fe}^{2+}_m\text{Mg}_z)\text{O}_{20}(\text{OH})_{4-n_i}$ .

In contrast to Equation (3), Equations (9a) and (9b) show that the relationships between  $n_i$  and  $m$ , and between  $p$  and  $m$ , are not constant, but they depend on the reduction level and on the value of  $K_0$ . For a given reduction level,  $p$  and  $n_i$  depend on the amount of Fe<sup>2+</sup>, and thus on the total Fe content in the smectite: The higher the  $m_{\text{tot}}$ , the higher the  $m$  and, consequently, the higher  $p$  and  $n_i$ . However, the ratio of  $p/n_i$  is independent of  $m_{\text{tot}}$ . It is convenient to exclude the amount of Fe<sup>2+</sup> from Equations (9a) and (9b):

$$p = m_{\text{rel}} m_{\text{tot}}/(1 + K_0 m_{\text{rel}}) \quad (9c)$$

$$n_i = K_0 m_{\text{rel}}^2 m_{\text{tot}}/(1 + K_0 m_{\text{rel}}). \quad (9d)$$

Thus,  $p$  and  $n_i$  may now be calculated as a function of  $m_{\text{rel}}$  and  $m_{\text{tot}}$  if  $K_0$  is known. If this model is valid, we can provide a rationale to explain quantitatively the change of surface-charge properties of smectites as a function of the reduction level. Thus, then Equations (6) and (9) may predict from the initial chemical composition, the evolution of the structural and chemical modifications of the layer and interlayer during Fe<sup>3+</sup> reduction.

## RESULTS AND DISCUSSION

### *Comparison between the observed and predicted layer charge as a function of reduction level*

Stucki *et al.* (1984) and Lear and Stucki (1985) studied the influence of Fe<sup>2+</sup> concentration on the layer charge for five dioctahedral smectites of different chemical compositions: Nontronite from Garfield, Washington (GAN) and Grant County, Washington (SWa-1), beidellite from Czechoslovakia #650 (CZM) and New Zealand (NZM), and montmorillonite from Upton, Wyoming (UPM). CEC and total Fe contents measured by Stucki *et al.* (1984) and Lear and Stucki (1985) are listed in Table 1. The layer charge of reduced samples ( $w_{\text{exp}}$ ) as a function of Fe<sup>2+</sup> ( $m_{\text{exp}}$ ) was determined by Stucki *et al.* (1984) on the basis of the

Table 1. CEC ( $w_0$ ), Fe concentration, and  $K_0$ , and  $K_r$  values for the smectite specimens studied by Stucki *et al.* (1984) and Lear and Stucki (1985).

Sample	$w_0$ (meq/g)	Total Fe <sup>3+</sup> (mmole/g clay)	$K_0$	$K_r$ (g/meq)
GAN	1.046	4.201	5.50	5.26
SWa-1	0.806	3.545	4.73	5.83
NZB	0.799	1.502	6.25	7.81
CZB	0.496	1.257	3.93	8.00
UPM	0.876	0.539	0.32	0.37

total Na concentration, and their results are shown in Figure 2. The trend of layer charge with Fe<sup>2+</sup>, as predicted by Equation (3), is represented by straight lines. As noted above, experimental results deviate from straight lines, and Stucki *et al.* (1984) found that the empirical relationship between  $w_{\text{exp}}$  and  $m_{\text{exp}}$  is described by:

$$w_{\text{exp}} = C_0 + C_1 m_{\text{exp}} + C_2 m_{\text{exp}}^2 \quad (10)$$

where  $C_0$ ,  $C_1$ , and  $C_2$  are empirical coefficients of unknown physical meaning. These values were determined by the best fit of experimental data (Stucki *et al.*, 1984). According to our model,  $w$  is expressed as a function of  $m$  by:

$$w = w_0 + p = w_0 + m/(1 + K_0 m_{\text{rel}}) \quad (11)$$

where  $w_0$  and  $m$  are expressed in mmoles of Na and of Fe<sup>2+</sup> per gram of clay. The layer charge,  $w$ , for each sample was calculated as a function of Fe<sup>2+</sup> by optimizing  $K_0$  values to superpose  $w$  functions and  $w_{\text{exp}}$  (Table 1). Note that Figure 2 shows that satisfactory fits of  $w$  values to experimental data were obtained, which provides good support of the validity of our model. As shown below,  $K_0$  values can be determined by an additional and independent method. Thus, Equation (11) may be used to predict layer charge of an iron-containing smectite as a function of its reduction level, provided that  $w_0$ ,  $K_0$ , and  $m_{\text{tot}}$  are known.

#### Comparison between the observed and predicted losses of structural OH groups as a function of reduction rate

The loss of structural OH groups as a function of the reduction level is difficult to measure. Lear and Stucki (1985) used tritium (<sup>3</sup>H) as a label element where 26% of structural H<sup>+</sup> in SWa-1 was replaced by <sup>3</sup>H<sup>+</sup> by suspending the nontronite sample in a <sup>3</sup>H solution heated at 125°C for 24 h. Then, the labeled sample was reduced by Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub>, and <sup>3</sup>H was measured against Fe<sup>2+</sup>. Lear and Stucki (1985) assumed that the loss of <sup>3</sup>H is proportional to the loss of structural OH groups in reduced samples and, based on this assumption, they determined the amount of structural protons lost as a function of the reduction level (Figure 3). The experimental relationship was fitted by the empirical linear equation:  $n_i = 0.77m$ .




Figure 3. Loss of H<sup>+</sup> by dehydroxylation as determined by tritium experiments (<sup>3</sup>H) (after Lear and Stucki, 1985) and calculated by Equation (9) ( $n_i$ ). The solid line was obtained by least-squares fit of observed data.

The loss of structural OH groups in SWa-1 ( $n_i$ ) can be calculated from Equation (9b), which supposes that the increase of the negative charge of the layer resulting from Fe<sup>3+</sup> reduction is entirely compensated by the sorption of Na and H<sup>+</sup>, and that all H<sup>+</sup> ions in the structure are consumed for the dehydroxylation of OH groups. The calculated  $n_i$  values are compared to the observed data in Figure 3. The match between the predicted and observed values is good, which supports our theoretical approach. However, the loss of structural H determined from the <sup>3</sup>H experiment is higher than that calculated in an amount equal to 0.2 mmole H<sup>+</sup>/g of clay where the Fe<sup>2+</sup> content is equal to 0.6–1.3 mmole/g; the differences between calculated and observed values gradually decreases to zero at greater reduction levels. The reason for this small discrepancy is unclear.

#### Determination of $K_0$

It follows from Equation (8) that the value of  $K_0$  determines the  $n_i/p$  ratio during the reduction process. The value of  $K_0$  of a smectite may be assumed to depend on the initial Fe<sup>3+</sup> content. This assumption is substantiated by the fact that the structural modification of *trans*-vacant (tv) Fe<sup>3+</sup>-containing dioctahedral 2:1 layer silicates during dehydroxylation depends on the amount of Al and Fe in the octahedral sheet. Indeed, the dehydroxylation of tv 2:1 layer silicates by heating results in the migration of 50% of the octahedral cations from *cis*- to *trans*-sites where Fe<sup>3+</sup> > Al. In contrast, octahedral cations preserve the original *cis*-site occupancy where Al is higher than Fe<sup>3+</sup> (Drits *et al.*, 1993; Muller *et al.*, 2000; Tshipursky *et al.*, 1985). Although these results were obtained for thermally treated samples, cation migration does not depend on the way the dehydroxylation is achieved (*i.e.*, either thermally or by reduction). Instead, it is deter-




Figure 4. Relationship between  $K_r$  and total content of Fe<sup>3+</sup> ( $m_{tot}$ ) for different beidellite and nontronite samples.

mined by the crystal chemistry because Manceau *et al.* (2000) showed that in the fully reduced Garfield nontronite, Fe<sup>2+</sup> cations occupy *trans*- and *cis*-sites to form trioctahedral domains. Cation migration and dehydroxylation of the octahedral sheet probably destabilize the crystal-lattice energy of reduced smectites, in an amount dependent on the total Fe content. To minimize this effect, Fe-rich smectites will tend to maximize the number of Na<sup>+</sup> ions in the interlayer and to minimize the loss of structural OH groups. Thus, the incorporation of Na<sup>+</sup> ions in the interlayer *vs.* cation migration and loss of structural OH groups are stabilizing and destabilizing factors, respectively.

Let us write Equation (9c) in the form:  $p/m_{rel} = m_{tot}/(1 + K_0 m_{rel})$ . This equation establishes that Fe-rich nontronites have a small  $K_0$  value to maximize the amount of sorbed Na ( $p$ ) at a given reduction rate. Conversely, the lower the Fe<sup>3+</sup> concentration, the lower the destabilization of the crystal-lattice energy is upon reduction and, accordingly,  $K_0$  is higher. However, Table 1 shows that there is no significant dependence between  $K_0$  and  $m_{tot}$ . In contrast to Equation (9c), both  $m_{tot}$  and  $K_0$  decrease in the order GAN, SWa-1, CZB, and UPM. Moreover, samples NZB and CZB have a similar  $m_{tot}$  but different values of  $K_0$ . However, the amount of Na sorbed from solution is controlled by the CEC of the oxidized smectite. Therefore, smectites with the same  $m_{tot}$ , but a different CEC value, have a different capacity to sorb Na in the interlayer for a given reduction rate. Based on electrostatic reasons, the higher the CEC, the lower the number of Na<sup>+</sup> ions can be assumed to be sorbed in the structure upon reduction because the interlayer has initially much Na.

To test this hypothesis, we normalize  $K_0$  values to the CEC by

$$K_r = K_0/CEC. \quad (12)$$

Note in Figure 4 that  $K_r$  and  $m_{tot}$  values are almost proportional for beidellite and nontronite samples. This relationship is expressed by:

$$K_r = 9.32 - 1.06m_{tot} + 0.02m_{tot}^2. \quad (13)$$

The dependence between  $K_r$  and  $m_{tot}$  of Equation (13) is in full agreement with the theoretical prediction: as the value of  $m_{tot}$  increases, the value of  $K_r$  decreases. Thus, the structural formula of a smectite sample is sufficient to determine  $K_r$  and  $K_0$  from Equations (12) and (13). However, Equations (12) and (13) are not universal and do not apply to montmorillonite. The  $K_r$  value for the UPM montmorillonite predicted from Equation (13) is equal to 8.75 g/meq and not 0.37 g/meq (Table 1). This large difference between predicted and experimental  $K_r$  values for UPM probably occurs because of a difference in the Fe<sup>3+</sup> reduction mechanism between montmorillonite and beidellite-nontronite. This is addressed below.

#### Relationship between $p$ and $n_i$ as a function of $m_{tot}$ , $m_{rel}$ , $K_0$ , and CEC

From the relationships between  $m_{tot}$ ,  $K_0$ , and CEC [Equations (12) and (13)], Equation (9) may be used to predict the amount of sorbed Na ( $p$ ) and H<sup>+</sup> ( $n_i$ ) from solution during the reduction of Fe<sup>3+</sup>. The  $p = f(m_{rel})$  and  $n_i = f(m_{rel})$  functions for three hypothetical smectites containing 4.0, 2.5, and 1.0 Fe<sup>3+</sup> atoms per unit cell, and having the same CEC of 1 meq/g, are plotted in Figure 5a.  $K_0$  values are calculated from Equations (12) and (13), and are equal to 5.4, 6.8, and 8.28, respectively.

Figure 5a shows the following: (1) For each sample, there is a value of  $m_{rel} = 1/K_0$  (or  $m_{rel}^{eq}$  where eq equals equilibrium) for which  $p = n_i$ . Below  $m_{rel}^{eq}$ ,  $p > n_i$ , which means that the increase in the negative charge of the layer is compensated predominantly by the sorption of Na<sup>+</sup> from solution. In contrast, above  $m_{rel}^{eq}$ , the increase in the negative charge of the layer is compensated by the sorption of H<sup>+</sup>. (2) The  $n_i$  function is nearly linear whereas  $p$  has a logarithm-like shape. Consequently, over most of the range of Fe reduction, H<sup>+</sup> is regularly sorbed, whereas most Na<sup>+</sup> is sorbed below  $m_{rel}^{eq}$ . For instance, for nontronite ( $m_{tot} = 4.0$ ), an increase of  $m_{rel}$  from 0.2 to 0.35 corresponds to an increase of sorbed Na<sup>+</sup> from 0.38 to 0.48 atoms per unit cell, whereas the H<sup>+</sup> content more than doubles from 0.42 to 0.92 atoms per unit cell. The variations of  $p$  and  $n_i$  functions with the total Fe concentration ( $m_{tot}$ ) for a given  $m_{rel}$  above the  $m_{rel}^{eq}$  value indicates a proton-exchange reaction. For example, at  $m_{rel} = 0.5$ ,  $p$  values for smectite samples with  $m_{tot} = 4.0, 2.5$ , and 1.0 are equal to 0.54, 0.28, and 0.09 Na, whereas  $n_i$  values for the same samples are equal to 1.46, 0.96, and 0.40 H<sup>+</sup> per unit cell. Therefore, a change of  $m_{tot}$  from 4.0 to 1.0 decreases  $p$  by only 0.45 Na, whereas  $n$  decreases by 1.06 H<sup>+</sup>.




Figure 5. a) Variations in the predicted amount of sorbed Na ( $p$ ) and loss of structural OH groups ( $n_i$ ) as a function of the  $\text{Fe}^{3+}$  reduction level ( $m_{\text{rel}}$ ) for three hypothetical smectites having a total Fe content ( $m_{\text{tot}}$ ) of 4.0, 2.5, and 1.0 atoms per unit cell, and an identical CEC of 1 meq/g ( $5.4 \leq K_0 \leq 8.28$ ). b) Same representation as a) for  $m_{\text{tot}} = 4.0$  and 2.0 and  $K_0 = 3.5$  and 6.5 ( $0.48 \leq \text{CEC} \leq 1.20$  meq/g).

Figure 5b represents  $p = f(m_{\text{rel}})$  and  $n_i = f(m_{\text{rel}})$  functions for two hypothetical smectites with identical  $K_0$  values (alternatively, either 3.5 or 6.5), but different total Fe contents ( $m_{\text{tot}}$  equals either 4.0 or 2.0). Thus, four  $m_{\text{rel}}^{\text{eq}}$  points shown in Figure 5b correspond to the following physico-chemical parameters: point 1:  $m_{\text{rel}}^{\text{eq}}$




Figure 6. Different lengths of OH-OH edges in tv (a) and cv (b) octahedral sheets.

$= 0.15$ ,  $m_{\text{tot}} = 4.0$ ,  $K_0 = 6.5$ ,  $\text{CEC} = 1.20$ ; point 2:  $m_{\text{rel}}^{\text{eq}} = 0.28$ ,  $m_{\text{tot}} = 4.0$ ,  $K_0 = 3.5$ ,  $\text{CEC} = 0.65$ ; point 3:  $m_{\text{rel}}^{\text{eq}} = 0.15$ ,  $m_{\text{tot}} = 2.0$ ,  $K_0 = 6.5$ ,  $\text{CEC} = 0.89$ ; and point 4:  $m_{\text{rel}}^{\text{eq}} = 0.28$ ,  $m_{\text{tot}} = 2.0$ ,  $K_0 = 3.5$ ,  $\text{CEC} = 0.48$ . Points 1 and 3, and 2 and 4 are aligned vertically because they have equal  $m_{\text{rel}}^{\text{eq}}$  values. Samples belonging to a vertical line have different  $m_{\text{tot}}$  and CEC values but equal  $K_0$  values. Points 1 and 2, and 3 and 4 are linear with the origin. Each line indicates an equal Fe content ( $m_{\text{tot}}$ ), but a different CEC. CEC increases towards the origin, and this trend is indicated on Figure 5b by the location of point 2 above point 1, and point 4 above point 3. This variation at constant  $m_{\text{tot}}$  value indicates that the layer charge is preferably compensated by the sorption of  $\text{H}^+$  rather than Na when CEC increases. This interesting finding is illustrated in Figure 5b by the line at  $m_{\text{rel}} = 0.24$ . At this reduction level, for a smectite with a total Fe content of 2.0, an increase in the CEC from 0.48 (point 4) to 0.89 meq/g (point 3) leads to a decrease in the amount of sorbed Na from 0.26 to 0.19 atoms per unit cell ( $\Delta p = 0.07$ ), and a concomitant increase of  $\text{H}^+$  from 0.22 to 0.29 ( $\Delta n_i = 0.07$ ).

#### Structural changes in reduced trans- and cis-vacant 2:1 layers

As shown above, Equations (12) and (13) successfully apply to Fe-rich beidellite and nontronite samples, but not to the UPM montmorillonite, and this is now discussed. According to Tsipursky and Drits (1984), montmorillonite has cis-vacant (cv) 2:1 layers, whereas beidellites and nontronites possess tv 2:1 layers. The octahedral edge formed by the two adjacent OH groups is short in tv sheets (2.40–2.45 Å), and longer in cv sheets (2.85–2.90 Å) because of the different electrostatic interactions owing to the configurations of octahedral cations in the two structures (Drits *et al.*, 1984, 1995) (Figure 6). This structural






Figure 7. Proposed structural mechanisms for the reduction of Fe<sup>3+</sup> to Fe<sup>2+</sup> in tv (a,b,c) and cv (d,e,f) octahedral sheets. OH, Fe, and vacant sites are represented by empty circles, filled circles, and squares, respectively. In a, b, d, and e, H<sub>2</sub>O molecules are in grey. See also Manceau *et al.* (2000).

difference is of key importance and it probably determines the behavior of tv and cv layers during the reduction of Fe<sup>3+</sup>.

Now consider the successive structural changes in the octahedral sheet of reduced tv and cv 2:1 layers. Structural studies of nontronite by P-EXAFS and XRD (Manceau *et al.*, 2000) and by Mössbauer spectroscopy (Russell *et al.*, 1979) showed that Fe atoms preserve their octahedral coordination in the reduced state. To maintain six-fold coordination of iron and to allow dehydroxylation, Fe<sup>2+</sup> must migrate to nearest vacant octahedral sites, otherwise Fe<sup>2+</sup> will be coordinated to five, or even four, oxygens (Manceau *et al.*, 2000). As seen in Figure 7, the most preferable local structural environment for the migration of Fe<sup>2+</sup> is an adjacent empty *trans*-site. Manceau *et al.* (2000) showed that in the Garfield nontronite, Fe<sup>2+</sup> migrates along the [010] direction to adjacent empty *trans*-sites to form trioctahedral-like Fe<sup>2+</sup> clusters. H<sub>2</sub>O molecules are formed owing to the protonation of edge-forming OH groups. A structural study of reduced SWa-1 (Manceau *et al.*, unpublished) showed that the migration of Fe<sup>2+</sup> cations also occurs despite the presence of Al and Mg in the octahedral sheet. Thus, Fe<sup>2+</sup> atoms migrate in tv 2:1 layers of nontronite and Fe-rich beidellite.

In comparison with tv layers, structural changes in reduced cv 2:1 layers are probably more complex. In cv octahedral sheets, OH pairs are bridged to three octahedral cations, and only two octahedral cations in tv sheets (Figure 7). Therefore, dehydroxylation in a cv framework, and the creation of local trioctahedral clusters as in a tv framework, require the simultaneous


migration of three neighboring Fe<sup>2+</sup> cations in [010], [310], and [310] directions. Such a significant structural modification requires high energy. This energy expenditure is probably even greater for low Fe<sup>3+</sup>-containing montmorillonite because the reduction of Fe<sup>3+</sup> requires that two Al, Mg cations migrate. Thus, in comparison with beidellites and nontronites, the greater activation energy of cv layers of montmorillonites will limit Fe<sup>2+</sup> migration. Therefore, it is unlikely that layers dehydroxylate in montmorillonites, and the deficit of positive charge created by the Fe<sup>3+</sup> to Fe<sup>2+</sup> reduction is probably compensated entirely by sorption of Na. This interpretation explains the low value of K<sub>0</sub> for UPM (0.32, Table 1). Thus, for this sample,  $p/n_i > 1$  [Equation (9)]. Further investigations are clearly needed to refine the local structure around Fe<sup>2+</sup> atoms in Fe-poor montmorillonite and beidellite.

#### Calculation of structural formulae


From the chemical composition of nontronite or Fe-rich beidellite samples, Equation (13) permits the determination of K<sub>r</sub> and, thereafter, K<sub>0</sub> is obtained from Equation (12). Then, the value of K<sub>0</sub> can estimate the amount of Na and H<sup>+</sup> sorbed from solution using Equation (9). If all the incorporated H<sup>+</sup> ions are consumed in dehydroxylation, then the structural formula of the reduced sample containing  $m_{\text{tot}}$  Fe<sup>2+</sup> ions per unit cell can be written in the form given by Equation (7). For example, let us consider the Garfield nontronite sample GO studied by Manceau *et al.* (2000) with a structural formula:




in which 50% of structural Fe<sup>3+</sup> was reduced. From Equations (9), (12), and (13),  $K_0 = 4.92$ ,  $p = 0.53$ ,  $n_i = 1.29$ , and  $w = w_0 + p = 0.81 + 0.54 = 1.35$  atoms of Na per unit cell. Then, the structural formula of this reduced sample is:




However, to preserve the octahedral coordination of reduced Fe<sup>2+</sup>, limited dehydroxylation occurs (see also Manceau *et al.*, 2000). To illustrate, let us calculate a structural formula for a 100% reduction level, assuming that all initially sorbed H<sup>+</sup> participates in subsequent dehydroxylation. In this case  $p = 0.63$ ,  $n_i = 3.02$ , and the resulting structural formula is:



With such a low amount of OH groups, Fe<sup>2+</sup> cannot be six-fold coordinated. If all H<sup>+</sup> ions in the structure are consumed by dehydroxylation, then Fe<sup>2+</sup> will be five-fold, or even four-fold coordinated. This conclusion contradicts the structural results obtained by Russell *et al.* (1979) and Manceau *et al.* (2000), who showed that Fe<sup>2+</sup> remains coordinated to six oxygens in reduced nontronite. Thus, the total OH of reduced nontronite will be much higher than that calculated in Equation (15). The structural model proposed by Manceau *et al.* (2000) for reduced Garfield contains 2.93 OH per unit cell. Based on this structural model, and on the reduction mechanism proposed here, the predicted structural formula is:




In this formula, the layer charge is compensated by the sorption of 1.44 Na and 1.95 H<sup>+</sup>. This interlayer Na (1.44) is close to that determined by Manceau *et al.* (2000) for reduced Garfield ( $1.3 \pm 0.1$ ) by chemical analysis and P-EXAFS:




The agreement between these two structural formulae provides additional support to the validity of our proposed reduction mechanism of Fe<sup>3+</sup> to Fe<sup>2+</sup> in dioctahedral smectites.

Equation (17) indicates that a limited number of OH groups dehydroxylate, and that the negative layer charge is partly compensated by H<sup>+</sup> sorbed from solution. This conclusion contrasts with tritium measurements, which showed that in SWa-1, the loss of structural OH occurs to  $\leq 80\%$  of Fe<sup>3+</sup> reduction. Figure 3 also shows that the measured loss of hydroxyls is in quantitative agreement with the prediction of Equations (6) and (9).

The high number of hydroxyl groups in reduced nontronite [Equation (18)] despite the massive loss of hydroxyl groups observed in tritium measurements can be explained by considering that the reduction process occurs as two distinct steps. In the first step, reactions occur according to Equations (6), and all sorbed H<sup>+</sup> ions are used in dehydroxylation. This reaction produces an unstable environment around Fe<sup>2+</sup> owing to the lower-coordination number, and also reduces the charge compensation of oxygen atoms, so they become strongly undersaturated (Manceau *et al.*, 2000). The stability of the local structure around Fe<sup>2+</sup> atoms are then reestablished by the sorption of two H<sub>2</sub>O molecules, which dissociate (H<sub>2</sub>O → OH<sup>-</sup> + H<sup>+</sup>). Newly formed hydroxyl groups then serve to rebuild the six-fold coordination of Fe<sup>2+</sup> atoms, and H<sup>+</sup> protons balance the local charge. These reactions probably occur simultaneously with, or immediately after, the migration of Fe<sup>2+</sup> from *cis* to *trans* sites. In this case, the reaction of the anionic framework to produce the transformation of (16) to (17) can be written:



This two-step reduction mechanism accounts for the continuous loss of structural OH with increasing Fe<sup>2+</sup> content as determined experimentally. This mechanism also accounts for the incorporation in the structure of aqueous H<sup>+</sup> ions measured by titration experiments for SWa-1 nontronite. As for reduced Garfield nontronite, this increase in H<sup>+</sup> may result from the partial or complete dissociation of sorbed H<sub>2</sub>O molecules during the second stage of the reduction process. To determine this quantitatively, the content of structural OH groups per unit cell must be known at a given reduction level. Unfortunately, at present this value is known only for 100%-reduced Garfield nontronite (Manceau *et al.*, 2000). However, the structural formula for SWa-1 involving the Fe<sup>2+</sup> content can be predicted from the initial composition (Si<sub>7.30</sub>Al<sub>0.70</sub>)(Al<sub>1.06</sub>Fe<sub>2.73</sub><sup>3+</sup>Mg<sub>0.26</sub>)O<sub>20</sub>(OH)<sub>4</sub>, Equation (9), and titration data (Figure 2; Lear and Stucki, 1985). For example, for  $m = 2.32$ ,  $m_{\text{tot}} = 2.73$ , and  $m_{\text{rel}} = 0.85$ , the expected structural formula is:



assuming that 0.40 H<sub>2</sub>O was incorporated into the structure.

In conclusion, the proposed reduction mechanism of Fe<sup>3+</sup> to Fe<sup>2+</sup> in dioctahedral smectites satisfactorily accounts for the modifications of the physico-chemical properties of 2:1 layers as measured by solution chem-

istry by various investigators. However, the absence of experimental data for OH content in smectites as a function of the reduction level prevents the establishment of a structural formula for dioctahedral beidelite-nontronite samples containing a variable amount of Fe<sup>2+</sup>.

#### ACKNOWLEDGMENTS

The authors acknowledge W.P. Gates for his constructive scientific comments and English corrections on a preliminary version of the paper, and S. Guggenheim is thanked for his editorial assistance. V.A. Drits acknowledges the Russian Science Foundation for partial funding of this study.

#### REFERENCES

- Chen, S.Z., Low, P.F., and Roth, C.B. (1987) Relation between potassium fixation and the oxidation state of octahedral iron. *Soil Science Society of America Journal*, **51**, 82–86.
- Drits, V.A., Plançon, A., Sakharov, B.A., Besson, G., Tshipursky, S.I., and Tchoubar, C. (1984) Diffraction effects calculated for structural models of K-saturated montmorillonite containing different types of defects. *Clay Minerals*, **19**, 541–561.
- Drits, V.A., Weber, A.I., Salyn, A.L., and Tshipursky, S.I. (1993) X-ray identification of one-layer illite varieties: Application to the study of illites around uranium deposits of Canada. *Clays and Clay Minerals*, **41**, 389–398.
- Drits, V.A., Besson, G., and Muller, F. (1995) An improved model for structural transformation of heat-treated aluminous dioctahedral 2:1 layer silicates. *Clays and Clay Minerals*, **43**, 718–731.
- Dunitz, J.D. (1956) The structure of sodium dithionite and the nature of the dithionite ion. *Acta Crystallographica*, **9**, 579–586.
- Egashira, K. and Ohtsubo, M. (1983) Swelling and mineralogy of smectites in paddy soils derived from marine alluvium, Japan. *Geoderma*, **29**, 119–127.
- Ernstsen, V. (1998) Reduction of nitrate by Fe<sup>2+</sup> in clay minerals. *Clays and Clay Minerals*, **44**, 599–608.
- Gan, H., Stucki, J.W., and Bailey, G.W. (1992) Reduction of structural iron in feruginous smectites by free radicals. *Clays and Clay Minerals*, **40**, 659–665.
- Gates, W.P., Jaunet, A.M., Tessier, D., Cole, M.A., Wilkinson, H.T., and Stucki, J.W. (1998) Swelling and texture of iron-bearing smectites reduced by bacteria. *Clays and Clay Minerals*, **46**, 487–497.
- Güven, N. (1991) Smectites. In *Hydrous Phyllosilicates (Exclusive of Micas)*, *Reviews in Mineralogy, Volume 19*, S.W. Bailey, ed., Mineralogical Society of America, Washington, D.C., 497–560.
- Heller-Kallai, L. (1997) Reduction and reoxidation of nontronite. The data reassessed. *Clays and Clay Minerals*, **45**, 476–479.
- Khaled, E.M. and Stucki, J.W. (1991) Effect of iron oxidation state on cation fixation in smectites. *Soil Science Society of America Journal*, **55**, 550–554.
- Lear, P.R. and Stucki, J.W. (1985) The role of structural hydrogen in the reduction and reoxidation of iron in nontronite. *Clays and Clay Minerals*, **33**, 539–545.
- Lear, P.R. and Stucki, J.W. (1989) Effects of iron oxidation state on the specific area of nontronites. *Clays and Clay Minerals*, **37**, 547–552.
- Low, P.F., Roth, C.B., and Stucki, J.W. (1983) System and method for rapid beneficiation of bentonite clay. U.S. Patent 4,411,530.
- Lynn, S., Rinker, R.G., and Corcoran, W.H. (1964) The monomerization rate of dithionite ion in aqueous solution. *Journal of Physical Chemistry*, **68**, 2363.
- Manceau, A., Drits, V.A., Lanson, B., Chateigner, D., Wu, J., Huo, D.F., Gates, W.P., and Stucki, J.W. (2000) Oxidation-reduction mechanism of iron in dioctahedral smectites. 2. Structural chemistry of reduced Garfield nontronite. *American Mineralogist*, **85**, 153–172.
- Muller, F., Drits, V.A., Plançon, A., Besson, G., and Tshipursky, S. (2000) Structural transformation of heat-treated Fe<sup>3+</sup>-rich dioctahedral micas. Part I: Characterization of the structural transformation. *Clay Minerals*, in press.
- Rinker, R.G., Gordon, T.P., Mason, D.M., and Corcoran, W.H. (1958) The presence of the SO<sub>2</sub><sup>-</sup> radical ion in aqueous solutions of sodium dithionite. *Journal of Physical Chemistry*, **63**, 302.
- Roth, C.B. and Tullock, R.J. (1973) Deprotonation of nontronite resulting from chemical reduction of structural ferric iron. In *Proceeding of the International Clay Conference*, J.M. Serratosa, ed., Division Ciencias C.S.I.C., Madrid, 107–114.
- Rozenon, I. and Heller-Kallai, L. (1976) Reduction and oxidation of Fe<sup>3+</sup> in dioctahedral smectite. 1: Reduction with hydrazine and dithionite. *Clays and Clay Minerals*, **24**, 271–282.
- Russell, J.D., Goodman, B.A., and Fraser, A.R. (1979) Infrared and Mössbauer studies of reduced nontronites. *Clays and Clay Minerals*, **27**, 63–71.
- Shen, S., Stucki, J.W., and Boast, C.W. (1992) Effects of structural iron reduction on the hydraulic conductivity of Na-smectite. *Clays and Clay Minerals*, **40**, 381–386.
- Stucki, J.W. (1988) Structural iron in smectites. In *Iron in Soils and Clay Minerals*, J.W. Stucki, B.A. Goodman, and U. Schwertmann, eds., Riedel Publishing Company, Dordrecht, 625–676.
- Stucki, J.W. and Roth, C.B. (1976) Interpretation of infrared spectra of oxidized and reduced nontronite. *Clays and Clay Minerals*, **24**, 293–296.
- Stucki, J.W. and Roth, C.B. (1977) Oxidation-reduction mechanism for structural iron in nontronite. *Soil Science Society of America Journal*, **41**, 808–814.
- Stucki, J.W. and Tessier, D. (1991) Effects of iron oxidation state on the structural order of Na-nontronite. *Clays and Clay Minerals*, **39**, 137–143.
- Stucki, J.W., Roth, C.B., and Baitinger, W.E. (1976) Analysis of iron-bearing clay minerals by electron spectroscopy for chemical analysis (ESCA). *Clays and Clay Minerals*, **24**, 289–292.
- Stucki, J.W., Golden, D.C., and Roth, C.B. (1984) Effects of reduction and reoxidation of structural iron on the surface charge and dissolution of dioctahedral smectites. *Clays and Clay Minerals*, **32**, 350–356.
- Stucki, J.W., Komadel, P., and Wilkinson, H.T. (1987) The microbial reduction of structural iron<sup>3+</sup> in smectites. *Soil Science Society of America Journal*, **51**, 1663–1665.
- Stucki, J.W., Bailey, G.W., and Gan, H. (1996) Oxidation-reduction mechanisms in iron-bearing phyllosilicates. *Applied Clay Science*, **10**, 417–430.
- Tshipursky, S.I. and Drits, V.A. (1984) The distribution of cations in the 2:1 layers of dioctahedral smectites studied by oblique-texture electron diffraction. *Clay Minerals*, **19**, 177–193.
- Tshipursky, S.I., Kamenova, M.Y., and Drits, V.A. (1985) Structural transformation of Fe<sup>3+</sup>-containing 2:1 dioctahedral phyllosilicates in the course of dehydroxylation. In *European Clay Conference*, J. Konta, ed., Univerzita Karlova Praha, Praha, 564–577.

E-mail of corresponding author: manceau@ujf-grenoble.fr

(Received 16 July 1998; accepted 28 September 1999; Ms. 98-099)