

AN INVERSE PROBLEM OF CALDERÓN TYPE WITH PARTIAL DATA

JUSSI BEHRNDT AND JONATHAN ROHLEDER

ABSTRACT. A generalized variant of the Calderón problem from electrical impedance tomography with partial data for anisotropic Lipschitz conductivities is considered in an arbitrary space dimension $n \geq 2$. The following two results are shown: (i) The selfadjoint Dirichlet operator associated with an elliptic differential expression on a bounded Lipschitz domain is determined uniquely up to unitary equivalence by the knowledge of the Dirichlet-to-Neumann map on an open subset of the boundary, and (ii) the Dirichlet operator can be reconstructed from the residuals of the Dirichlet-to-Neumann map on this subset.

1. INTRODUCTION AND MAIN RESULTS

Let \mathcal{L} be a uniformly elliptic and formally symmetric second order differential expression of the form

$$(1.1) \quad \mathcal{L} = - \sum_{j,k=1}^n \partial_j a_{jk} \partial_k + \sum_{j=1}^n (a_j \partial_j - \partial_j \bar{a}_j) + a$$

with variable coefficients on a bounded Lipschitz domain Ω in \mathbb{R}^n , $n \geq 2$. The main objective of the present paper is to show that the selfadjoint *Dirichlet operator*

$$(1.2) \quad Au = \mathcal{L}u, \quad \text{dom } A = \{u \in H_0^1(\Omega) : \mathcal{L}u \in L^2(\Omega)\},$$

associated with \mathcal{L} in $L^2(\Omega)$ is uniquely determined by a local variant of the Dirichlet-to-Neumann map on some open subset ω of the boundary $\partial\Omega$, and that A can be reconstructed from the residuals of this partial Dirichlet-to-Neumann map. Here the *Dirichlet-to-Neumann map* M is defined by

$$(1.3) \quad M(\lambda) : H^{1/2}(\partial\Omega) \rightarrow H^{-1/2}(\partial\Omega), \quad u_\lambda|_{\partial\Omega} \mapsto \partial_{\mathcal{L}} u_\lambda|_{\partial\Omega},$$

where $u_\lambda \in H^1(\Omega)$ is a solution of $\mathcal{L}u = \lambda u$ and $\partial_{\mathcal{L}} u_\lambda|_{\partial\Omega}$ denotes the conormal derivative of u at $\partial\Omega$. The mapping $M(\lambda)$ is well-defined for all λ in the resolvent set $\rho(A)$ of A ; see Section 2 for further details.

The above inverse problem is closely connected to and inspired by the famous Calderón problem from electrical impedance tomography, where the aim is to determine the isotropic conductivity γ of an inhomogeneous body from current and voltage data measured on the surface or on parts of it. The classical Calderón problem corresponds to the special case $a_{jk} = \gamma \delta_{jk}$, $a_j = a = 0$, $1 \leq j, k \leq n$, in (1.1), and the knowledge of $M(\lambda)$ on $\partial\Omega$ or $\omega \subset \partial\Omega$ for some λ or at $\lambda = 0$; see [17].

The Calderón problem has been a major challenge in the field of inverse problems for PDEs in the last three decades. The first positive results were obtained for $\omega = \partial\Omega$ using only Dirichlet and Neumann data for $\lambda = 0$ in the pioneering work [60] in dimension $n \geq 3$ and smooth γ , see also [52] for $\gamma \in C^{1,1}(\bar{\Omega})$ and [49] for the reconstruction of γ from the boundary measurements. In the two-dimensional case the first main contribution was the solution of the problem for $\gamma \in W^{2,p}(\Omega)$ in [50]; later in [10] less regular conductivities in $L^\infty(\Omega)$ were allowed. For partial data

given only on special subsets of $\partial\Omega$ uniqueness was shown in the recent works [16, 39] for a C^2 -function γ and a reconstruction method was provided in [51]. Also the more general case of an anisotropic conductivity $(a_{jk})_{j,k=1}^n$ has been investigated; in this situation, the single coefficients are in general not uniquely determined. Nevertheless, uniqueness up to diffeomorphisms was shown for smooth coefficients assuming knowledge of $M(0)$ in [44] in dimension $n \geq 3$ and in [59, 61] for $n = 2$. Less regular coefficients were treated in dimension $n = 2$ in [9, 58]. In the publications [42, 43] the related problem of determining a real-analytic Riemannian manifold from the given Dirichlet and Neumann boundary data on arbitrary open subsets of $\partial\Omega$ was considered. There uniqueness up to isometry in $n \geq 3$ and uniqueness of the conformal class in $n = 2$ was shown for partial data supported in an open subset of $\partial\Omega$; see [24] for further recent results of this type and [38] for results on smooth Riemannian manifolds and elliptic operators with smooth coefficients as well as related problems as, e.g., the Gelfand problem for the wave equation.

The aim of the present paper is to prove somewhat different, milder type of uniqueness and reconstruction results in space dimension $n \geq 2$ for partial data given on an open subset ω of $\partial\Omega$. Since the coefficients of \mathcal{L} are not uniquely determined in general we focus on the selfadjoint Dirichlet operator (1.2) associated with \mathcal{L} on Ω . In return this point of view onto the problem allows us to consider the more general differential expression (1.1) and to impose, in an arbitrary space dimension $n \geq 2$, the following comparatively weak assumptions on the coefficients of \mathcal{L} .

Assumption 1.1. The coefficients a_{jk} and a_j are bounded Lipschitz functions on $\bar{\Omega}$ satisfying $a_{kj} = \overline{a_{jk}}$, $1 \leq j, k \leq n$, and $a \in L^\infty(\Omega)$ is real-valued. Moreover, \mathcal{L} is uniformly elliptic, i.e., the matrix $(a_{jk}(x))_{j,k=1}^n$ is uniformly positive for all $x \in \bar{\Omega}$.

Furthermore, we impose the following conditions on the domain Ω and the subset ω of the boundary $\partial\Omega$ where Dirichlet and Neumann data is assumed to be given.

Assumption 1.2. Ω is a bounded Lipschitz domain in \mathbb{R}^n , $n \geq 2$, and ω is a nonempty, open subset of the boundary $\partial\Omega$.

The following two theorems on uniqueness and reconstruction of the Dirichlet operator A in (1.2) from the knowledge of the Dirichlet-to-Neumann map on $\omega \subseteq \partial\Omega$ are the main results in the present paper. The usual duality between $H^{1/2}(\partial\Omega)$ and $H^{-1/2}(\partial\Omega)$ is denoted by (\cdot, \cdot) .

Theorem 1.3. *Let Ω and ω be as in Assumption 1.2, and let \mathcal{L}_1 and \mathcal{L}_2 be differential expressions on Ω of the form (1.1) with coefficients $a_{jk,1}$, $a_{j,1}$, a_1 and $a_{jk,2}$, $a_{j,2}$, a_2 satisfying Assumption 1.1. Denote by A_1 , A_2 , and M_1 , M_2 the corresponding Dirichlet operators and Dirichlet-to-Neumann maps, respectively. Assume that*

$$(M_1(\lambda)\varphi, \psi) = (M_2(\lambda)\varphi, \psi) \quad \varphi, \psi \in H^{1/2}(\partial\Omega), \text{ supp } \varphi, \psi \subset \omega,$$

holds for all $\lambda \in \mathcal{D}$, where $\mathcal{D} \subseteq \rho(A_1) \cap \rho(A_2)$ is a set of points which has an accumulation point in $\rho(A_1) \cap \rho(A_2)$. Then A_1 and A_2 are unitarily equivalent.

We point out that Theorem 1.3 is a ‘‘mild’’ uniqueness result in the sense that even in the special case $a_{jk} = \gamma\delta_{jk}$, $a_j = a = 0$, $1 \leq j, k \leq n$ and $\omega = \partial\Omega$ in the classical Calderón problem it does not imply uniqueness of γ as shown in [10, 16, 39, 49, 50, 51, 52, 60].

In order to state our second main result we remark that $\lambda \mapsto M(\lambda)$ is a meromorphic operator function; the residual of the Dirichlet-to-Neumann map M at a pole μ is a linear operator from $H^{1/2}(\partial\Omega)$ to $H^{-1/2}(\partial\Omega)$ which is denoted by $\text{Res}_\mu M$. Since the aim in the next theorem is to use only knowledge of M and its residuals

on ω , also the operator $\text{Res}_\mu M$ is replaced by a local version. The residual $\text{Res}_\mu^\omega M$ on ω is defined on the linear subspace $H_\omega^{1/2}(\partial\Omega)$ of functions $\varphi \in H^{1/2}(\partial\Omega)$ with $\text{supp } \varphi \subset \omega$ by

$$\text{Res}_\mu^\omega M : H_\omega^{1/2}(\partial\Omega) \rightarrow (H_\omega^{1/2}(\partial\Omega))', \quad (\text{Res}_\mu^\omega M \varphi)(\psi) := (\psi, \text{Res}_\mu M \varphi),$$

where $\varphi, \psi \in H_\omega^{1/2}(\partial\Omega)$. The next theorem provides a one-to-one correspondence between the eigenfunctions of the Dirichlet operator A and the residuals of the Dirichlet-to-Neumann map M on ω and, hence, yields a representation of A in terms of these residuals.

Theorem 1.4. *Let Assumption 1.1 and Assumption 1.2 be satisfied and let M be the Dirichlet-to-Neumann map associated with the elliptic differential expression \mathcal{L} . Then the poles of M coincide with the eigenvalues of the Dirichlet operator A and for each eigenvalue λ_k of A the corresponding eigenspace is given by*

$$\ker(A - \lambda_k) = \text{ran}(\tau_k^{-1} \circ \text{Res}_{\lambda_k}^\omega M),$$

where τ_k denotes the restriction of the Neumann trace operator $u \mapsto \partial_{\mathcal{L}} u|_\omega$ onto $\ker(A - \lambda_k)$. In particular, there exist $\varphi_1^{(k)}, \dots, \varphi_{n(k)}^{(k)} \in H^{1/2}(\partial\Omega)$ with support in ω , such that the functions

$$e_i^{(k)} := \tau_k^{-1}(\text{Res}_{\lambda_k}^\omega M) \varphi_i^{(k)}, \quad i = 1, \dots, n(k),$$

form an orthonormal basis of $\ker(A - \lambda_k)$ and A can be represented in the form

$$Au = \sum_{k=1}^{\infty} \lambda_k \sum_{i=1}^{n(k)} (u, e_i^{(k)}) e_i^{(k)}, \quad u \in \text{dom } A.$$

The proofs of our uniqueness and reconstruction results are based on the powerful interplay of modern operator theory with classical PDE techniques, as, e.g., unique continuation results from [7, 8, 36, 62]. Some of the main ideas are inspired by abstract methods in extension and spectral theory of symmetric and selfadjoint operators as provided in [2, 11, 12, 19, 20, 21, 22, 37, 41]. Further operator theoretic approaches to elliptic boundary value problems and related questions via Dirichlet-to-Neumann maps or other analytic operator functions can also be found in the recent publications [3, 4, 5, 13, 14, 15, 28, 30, 31, 32, 56, 57]. For general methods from extension theory of symmetric operators that are applied to elliptic PDEs we also refer to, e.g., [6, 27, 29, 33, 46, 47, 53, 54], the monographs [26, 34, 45, 48], and the references therein.

The main part of the present paper is devoted to the proofs of the two main results; along the proofs we show Proposition 2.5 and Proposition 2.7 on the eigenvalues and eigenspaces of the Dirichlet operator which are of independent interest. The paper closes with a short appendix which summarizes some basic facts on unbounded operators in Hilbert spaces and on Banach space-valued analytic functions.

2. PROOFS OF THEOREM 1.3 AND THEOREM 1.4

In this section we give complete proofs of the uniqueness and reconstruction theorems from above. Instead of two single proofs the material is ordered in several smaller statements which then lead to the proofs of the main results.

We fix some notation first. By $H^s(\Omega)$ and $H^s(\partial\Omega)$ we denote the Sobolev spaces of order $s \geq 0$ on Ω and $\partial\Omega$, respectively, and by $H_0^s(\Omega)$ the closure of the set of C^∞ -functions with compact support in Ω with respect to the H^s -norm. Further, $H^{-s}(\partial\Omega)$ denotes the dual space of $H^s(\partial\Omega)$; the duality is expressed via

$$(2.1) \quad f(\varphi) = (\varphi, f) = \overline{(f, \varphi)}, \quad f \in H^{-s}(\partial\Omega), \quad \varphi \in H^s(\partial\Omega).$$

For the Lipschitz-domain Ω we write $u|_{\partial\Omega} \in H^{1/2}(\partial\Omega)$ for the trace of $u \in H^1(\Omega)$ on the boundary $\partial\Omega$ and $\partial_{\mathcal{L}}u|_{\partial\Omega} \in H^{-1/2}(\partial\Omega)$ for the conormal derivative or Neumann trace of u at $\partial\Omega$, see, e.g., [48] for more details. Recall further that $H_0^1(\Omega)$ coincides with the kernel of the trace operator $u \mapsto u|_{\partial\Omega}$ on $H^1(\Omega)$. In order to define a restriction of the conormal derivative to ω let

$$(2.2) \quad H_{\omega}^{1/2}(\partial\Omega) = \{\varphi \in H^{1/2}(\partial\Omega) : \text{supp } \varphi \subset \omega\}$$

be the linear subspace of $H^{1/2}(\partial\Omega)$ which consists of functions with support in ω . The restriction $\partial_{\mathcal{L}}u|_{\omega}$ of the Neumann trace $\partial_{\mathcal{L}}u|_{\partial\Omega}$ to ω is defined as

$$(2.3) \quad (\partial_{\mathcal{L}}u|_{\omega})(\varphi) := (\partial_{\mathcal{L}}u|_{\partial\Omega})(\varphi) = (\varphi, \partial_{\mathcal{L}}u|_{\partial\Omega}), \quad \varphi \in H_{\omega}^{1/2}(\partial\Omega).$$

Let us recall some well-known properties of the Dirichlet operator associated to \mathcal{L} , which can be found in, e.g., [26, Chapter VI] and [48, Chapter 4].

Proposition 2.1. *The Dirichlet operator A in (1.2) is a selfadjoint operator in $L^2(\Omega)$ and its spectrum $\sigma(A)$ consists of isolated (real) eigenvalues with finite-dimensional eigenspaces. The Dirichlet eigenvalues accumulate to $+\infty$ and are bounded from below.*

The next lemma shows that the Dirichlet-to-Neumann map and the Poisson operator in Definition 2.3 below are well-defined.

Lemma 2.2. *For all λ in the resolvent set $\rho(A)$ of A and all $\varphi \in H^{1/2}(\partial\Omega)$ there exists a unique solution $u_{\lambda} \in H^1(\Omega)$ of the boundary value problem*

$$(2.4) \quad \mathcal{L}u = \lambda u, \quad u|_{\partial\Omega} = \varphi.$$

In particular, for all $\lambda \in \rho(A)$ the Dirichlet-to-Neumann map $M(\lambda)$ in (1.3) is well-defined.

Proof. Let $\lambda \in \rho(A)$ and $\varphi \in H^{1/2}(\partial\Omega)$. Then the homogeneous problem

$$(\mathcal{L} - \lambda)u = 0, \quad u|_{\partial\Omega} = 0,$$

has only the trivial solution, and by [48, Theorem 4.10] it follows that the inhomogeneous problem (2.4) has a unique solution. \square

Besides the Dirichlet-to-Neumann map a Poisson operator which maps functions on $\partial\Omega$ onto the corresponding solutions of (2.4) will play an important role.

Definition 2.3. Let $\lambda \in \rho(A)$. The *Poisson operator* is defined by

$$(2.5) \quad \gamma(\lambda) : H^{1/2}(\partial\Omega) \rightarrow L^2(\Omega), \quad u_{\lambda}|_{\partial\Omega} \mapsto u_{\lambda},$$

where $u_{\lambda} \in H^1(\Omega)$ is the unique solution of (2.4) with $\varphi = u_{\lambda}|_{\partial\Omega}$. The range of the restriction of the Poisson operator to $H_{\omega}^{1/2}(\partial\Omega)$ is denoted by \mathcal{N}_{λ} ,

$$(2.6) \quad \mathcal{N}_{\lambda} = \{u \in H^1(\Omega) : \mathcal{L}u = \lambda u, \text{supp}(u|_{\partial\Omega}) \subset \omega\}.$$

The operator $\gamma(\lambda)$ is well-defined for each $\lambda \in \rho(A)$ by Lemma 2.2 and the relation $M(\lambda)\varphi = \partial_{\mathcal{L}}(\gamma(\lambda)\varphi)|_{\partial\Omega}$ holds for all $\varphi \in H^{1/2}(\partial\Omega)$. Some properties and formulas for the Poisson operator and the Dirichlet-to-Neumann map will be given in the next lemma. Its proof is essentially based on the second Green identity

$$(2.7) \quad (\mathcal{L}u, v) - (u, \mathcal{L}v) = (u|_{\partial\Omega}, \partial_{\mathcal{L}}v|_{\partial\Omega}) - (\partial_{\mathcal{L}}u|_{\partial\Omega}, v|_{\partial\Omega})$$

for $u, v \in H^1(\Omega)$ satisfying $\mathcal{L}u, \mathcal{L}v \in L^2(\Omega)$, see, e.g., [48]. Here (\cdot, \cdot) on the left hand side denotes the inner product in $L^2(\Omega)$ and on the right hand side the duality between $H^{-1/2}(\partial\Omega)$ and $H^{1/2}(\partial\Omega)$; cf. (2.1). In the following it will be clear from the context if the entries in (\cdot, \cdot) are functions on Ω or $\partial\Omega$, respectively, so that no confusion can arise. We remark that in a more abstract setting statements of

similar form as in Lemma 2.4 can be found in, e.g., [11, Proposition 2.6], [20, § 1], and [41, § 2].

Lemma 2.4. *Let $\lambda, \mu \in \rho(A)$, let $\gamma(\lambda), \gamma(\mu)$ be the Poisson operators and let $M(\lambda), M(\mu)$ be the Dirichlet-to-Neumann maps. Then the following statements (i)-(iv) hold.*

- (i) $\gamma(\lambda)$ is bounded and the adjoint operator $\gamma(\lambda)' : L^2(\Omega) \rightarrow H^{-1/2}(\partial\Omega)$ is given by

$$\gamma(\lambda)'u = -\partial_{\mathcal{L}}((A - \bar{\lambda})^{-1}u)|_{\partial\Omega}, \quad u \in L^2(\Omega).$$

- (ii) $\gamma(\lambda)$ and $\gamma(\mu)$ satisfy the identity

$$\gamma(\lambda) = (I + (\lambda - \mu)(A - \lambda)^{-1})\gamma(\mu).$$

- (iii) The Poisson operators and the Dirichlet-to-Neumann maps are connected via

$$(\bar{\mu} - \lambda)(\gamma(\lambda)\varphi, \gamma(\mu)\psi) = (M(\lambda)\varphi, \psi) - (\varphi, M(\mu)\psi),$$

and, in particular, $(M(\lambda)\varphi, \psi) = (\varphi, M(\bar{\lambda})\psi)$ holds for all $\varphi, \psi \in H^{1/2}(\partial\Omega)$.

- (iv) $M(\lambda)$ is bounded, the function $\lambda \mapsto M(\lambda)$ is holomorphic on $\rho(A)$, and the identity

$$\begin{aligned} (M(\lambda)\varphi, \psi) &= (\varphi, M(\lambda_0)\psi) \\ &\quad + (\bar{\lambda}_0 - \lambda) \left((1 + (\lambda - \lambda_0)(A - \lambda)^{-1})\gamma(\lambda_0)\varphi, \gamma(\lambda_0)\psi \right) \end{aligned}$$

holds for all $\lambda, \lambda_0 \in \rho(A)$ and $\varphi, \psi \in H^{1/2}(\partial\Omega)$. In particular, every eigenvalue of A is either a pole of order one or a removable singularity of the mapping $\lambda \mapsto M(\lambda)$.

Proof. (i) In order to verify the formula for $\gamma(\lambda)'$ we have to show

$$(\gamma(\lambda)\varphi, u) = (\varphi, -\partial_{\mathcal{L}}((A - \bar{\lambda})^{-1}u)|_{\partial\Omega}) \quad \text{for all } \varphi \in H^{1/2}(\partial\Omega), \quad u \in L^2(\Omega);$$

cf. (2.1). Let $\varphi \in H^{1/2}(\partial\Omega)$, set $u_\lambda = \gamma(\lambda)\varphi$ and let $u \in L^2(\Omega)$. One finds by (2.7)

$$\begin{aligned} (\gamma(\lambda)\varphi, u) &= (u_\lambda, (I + \bar{\lambda}(A - \bar{\lambda})^{-1})u) - (\lambda u_\lambda, (A - \bar{\lambda})^{-1}u) \\ &= (u_\lambda, \mathcal{L}(A - \bar{\lambda})^{-1}u) - (\mathcal{L}u_\lambda, (A - \bar{\lambda})^{-1}u) \\ &= (\partial_{\mathcal{L}}u_\lambda|_{\partial\Omega}, (A - \bar{\lambda})^{-1}u|_{\partial\Omega}) - (u_\lambda|_{\partial\Omega}, \partial_{\mathcal{L}}((A - \bar{\lambda})^{-1}u)|_{\partial\Omega}) \\ &= (\varphi, -\partial_{\mathcal{L}}((A - \bar{\lambda})^{-1}u)|_{\partial\Omega}), \end{aligned}$$

where we have used $(A - \bar{\lambda})^{-1}u|_{\partial\Omega} = 0$ in the last equality. It follows that $\gamma(\lambda)'$ acts as in the assertion and is defined on $L^2(\Omega)$. Moreover, the above reasoning implies that $\gamma(\lambda)$ is closed and hence bounded by the closed graph theorem.

- (ii) For $\lambda, \mu \in \rho(A)$, $\varphi \in H^{1/2}(\partial\Omega)$ and $u \in L^2(\Omega)$ we find by (i)

$$\begin{aligned} (\gamma(\lambda)\varphi, u) - (\gamma(\mu)\varphi, u) &= (\varphi, -\partial_{\mathcal{L}}((A - \bar{\mu})^{-1}(\bar{\lambda} - \bar{\mu})(A - \bar{\lambda})^{-1}u)|_{\partial\Omega}) \\ &= (\gamma(\mu)\varphi, (\bar{\lambda} - \bar{\mu})(A - \bar{\lambda})^{-1}u) \\ &= ((\lambda - \mu)(A - \lambda)^{-1}\gamma(\mu)\varphi, u). \end{aligned}$$

Hence we have $\gamma(\lambda)\varphi = \gamma(\mu)\varphi + (\lambda - \mu)(A - \lambda)^{-1}\gamma(\mu)\varphi$ which shows (ii).

(iii) Let $\lambda, \mu \in \rho(A)$ and $\varphi, \psi \in H^{1/2}(\partial\Omega)$, and set $u_\lambda = \gamma(\lambda)\varphi$ and $v_\mu = \gamma(\mu)\psi$. Again by (2.7) we find

$$(\bar{\mu} - \lambda)(\gamma(\lambda)\varphi, \gamma(\mu)\psi) = (u_\lambda, \mathcal{L}v_\mu) - (\mathcal{L}u_\lambda, v_\mu) = (M(\lambda)\varphi, \psi) - (\varphi, M(\mu)\psi).$$

(iv) From $(M(\lambda)\varphi, \psi) = (\varphi, M(\bar{\lambda})\psi)$ for $\lambda \in \rho(A)$ and $\varphi, \psi \in H^{1/2}(\partial\Omega)$ it follows that $M(\lambda) : H^{1/2}(\partial\Omega) \rightarrow H^{-1/2}(\partial\Omega)$ is closed and hence bounded by the closed graph theorem. Furthermore, by (iii) we have

$$(M(\lambda)\varphi, \psi) = (\varphi, M(\lambda_0)\psi) + (\bar{\lambda}_0 - \lambda)(\gamma(\lambda)\varphi, \gamma(\lambda_0)\psi)$$

which together with (ii) implies the formula in (iv). The remaining statements in (iv) follow from this and the corresponding properties of the resolvent of A , see also the Appendix. \square

The statement in Lemma 2.4 (iv) on the singularities of M and the eigenvalues of A will be improved later in Corollary 2.8, where it turns out that the Dirichlet eigenvalues coincide with the poles of the meromorphic operator function $\lambda \mapsto M(\lambda)$ and its restriction on ω .

The following proposition is essential for the proof of our main results. It states that the restriction of the Dirichlet operator A onto $\{u \in \text{dom } A : \partial_{\mathcal{L}}u|_{\omega} = 0\}$ is an operator without eigenvalues. In an operator theoretic language this implies that this restriction is a *simple* symmetric operator. The main ingredient in the proof of Proposition 2.5 is a classical unique continuation theorem for solutions of second order elliptic differential inequalities, see, e.g., [62] and [7, 8, 36]. Moreover, the proof makes use of the sesquilinear form $\Phi_{\mathcal{L}}(\cdot, \cdot)$ induced by \mathcal{L} on $H^1(\Omega)$,

$$\Phi_{\mathcal{L}}(u, v) := \int_{\Omega} \left(\sum_{j,k=1}^n a_{jk} \partial_k u \cdot \overline{\partial_j v} + \sum_{j=1}^n (a_j(\partial_j u) \cdot \bar{v} + \bar{a}_j u \cdot \overline{\partial_j v}) + au\bar{v} \right) dx$$

for $u, v \in H^1(\Omega)$, and the corresponding first Green identity

$$(2.8) \quad \Phi_{\mathcal{L}}(u, v) = (\mathcal{L}u, v) + (\partial_{\mathcal{L}}u|_{\partial\Omega}, v|_{\partial\Omega})$$

for all $u, v \in H^1(\Omega)$ satisfying $\mathcal{L}u \in L^2(\Omega)$; see [48].

Proposition 2.5. *There exists no eigenfunction u of the Dirichlet operator A satisfying $\partial_{\mathcal{L}}u|_{\omega} = 0$.*

Proof. Let $\tilde{\Omega} \supseteq \Omega$ be a bounded Lipschitz domain such that $\partial\Omega \setminus \omega \subset \partial\tilde{\Omega}$ and $\tilde{\Omega} \setminus \Omega$ contains an open ball \mathcal{O} . We extend the coefficients a_{jk} , a_j , and a of the differential expression \mathcal{L} from (1.1) to functions \tilde{a}_{jk} , \tilde{a}_j , and \tilde{a} on $\tilde{\Omega}$ such that Assumption 1.1 holds for the corresponding differential expression $\tilde{\mathcal{L}}$ in $\tilde{\Omega}$ defined as in (1.1).

Assume that there exists λ and $u \neq 0$ in the domain of the Dirichlet operator (1.2) with $\mathcal{L}u = \lambda u$ on Ω and $\partial_{\mathcal{L}}u|_{\omega} = 0$. Since we have $u|_{\partial\Omega} = 0$, we can extend u by zero on $\tilde{\Omega} \setminus \Omega$ to a function $\tilde{u} \in H^1(\tilde{\Omega})$. Moreover, $\tilde{\mathcal{L}}\tilde{u} \in L^2(\tilde{\Omega})$ and $\tilde{\mathcal{L}}\tilde{u} = \lambda\tilde{u}$ holds. In fact, we compute for $\tilde{\varphi} \in C_0^\infty(\tilde{\Omega})$

$$(\tilde{\mathcal{L}}\tilde{u} - \lambda\tilde{u})(\tilde{\varphi}) = \Phi_{\tilde{\mathcal{L}}}(\tilde{u}, \tilde{\varphi}) - (\lambda\tilde{u}, \tilde{\varphi}),$$

where the left hand side is understood in the sense of distributions and the right hand side consists of integrals on $\tilde{\Omega}$ with integrands vanishing outside of Ω . Denoting the restriction of $\tilde{\varphi}$ to Ω by φ , it follows with the help of the first Green identity (2.8) that

$$(\tilde{\mathcal{L}}\tilde{u} - \lambda\tilde{u})(\tilde{\varphi}) = \Phi_{\mathcal{L}}(u, \tilde{\varphi}) - \lambda(u, \tilde{\varphi}) = (\mathcal{L}u, \tilde{\varphi}) + (\partial_{\mathcal{L}}u|_{\partial\Omega}, \tilde{\varphi}|_{\partial\Omega}) - \lambda(u, \tilde{\varphi}).$$

Since $\mathcal{L}u = \lambda u$, $\text{supp}(\tilde{\varphi}|_{\partial\Omega}) \subset \omega$ and $\partial_{\mathcal{L}}u|_{\omega} = 0$ we conclude together with (2.3)

$$(\tilde{\mathcal{L}}\tilde{u} - \lambda\tilde{u})(\tilde{\varphi}) = (\partial_{\mathcal{L}}u|_{\partial\Omega}, \tilde{\varphi}|_{\partial\Omega}) = \overline{(\partial_{\mathcal{L}}u|_{\omega})(\tilde{\varphi}|_{\omega})} = 0.$$

Hence $\tilde{\mathcal{L}}\tilde{u} \in L^2(\tilde{\Omega})$ and $\tilde{\mathcal{L}}\tilde{u} = \lambda\tilde{u}$ hold; in particular, \tilde{u} is locally in H^2 , see, e.g., [48, Theorem 4.16]. Furthermore, we obtain

$$-\sum_{j,k=1}^n \tilde{a}_{jk} \partial_j \partial_k \tilde{u} = \left(\lambda - \tilde{a} + \sum_{j=1}^n \partial_j \tilde{a}_j \right) \tilde{u} + \sum_{k=1}^n \left(\sum_{j=1}^n \partial_j \tilde{a}_{jk} - \tilde{a}_k + \tilde{a}_k \right) (\partial_k \tilde{u}).$$

Since the functions \tilde{a}_{jk} and \tilde{a}_j , $1 \leq j, k \leq n$, together with their derivatives of first order as well as \tilde{a} are bounded on Ω and $\tilde{u} = 0$ on $\tilde{\Omega} \setminus \Omega$ there exist constants α and β such that

$$(2.9) \quad \left| \sum_{j,k=1}^n \tilde{a}_{jk} \partial_j \partial_k \tilde{u} \right| \leq \alpha |\tilde{u}| + \beta \sum_{k=1}^n |\partial_k \tilde{u}|$$

holds a.e. on $\tilde{\Omega}$. As $\tilde{u} = 0$ on \mathcal{O} it follows from the differential inequality (2.9) and classical unique continuation results that \tilde{u} vanishes identically on $\tilde{\Omega}$; cf. [62]. In particular, we conclude $u = 0$ on Ω , a contradiction, since u was chosen to be an eigenfunction of A . \square

Our last preparatory lemma will establish, as a consequence of Proposition 2.5, a density statement on the ranges \mathcal{N}_λ of the Poisson operators $\gamma(\lambda)$ in (2.5) restricted to $\mathcal{H}_\omega^{1/2}(\partial\Omega)$. Recall that \mathcal{N}_λ is the space of solutions u of the boundary value problem (2.4) which satisfy $\text{supp}(u|_{\partial\Omega}) \subset \omega$.

Lemma 2.6. *Let $\mathcal{O} \subseteq \mathbb{C}^+$ be an open set and let $\mathcal{O}^* = \{\lambda \in \mathbb{C} : \bar{\lambda} \in \mathcal{O}\}$. Then*

$$\text{span}\{\mathcal{N}_\lambda : \lambda \in \mathcal{O} \cup \mathcal{O}^*\} = \text{span}\{\gamma(\lambda)\varphi : \varphi \in H_\omega^{1/2}(\partial\Omega), \lambda \in \mathcal{O} \cup \mathcal{O}^*\}$$

is a dense subspace of $L^2(\Omega)$.

Proof. The proof consists of three separate steps. It makes use of two further operator realizations S and T of the differential expression \mathcal{L} . We consider the restriction

$$Su = \mathcal{L}u, \quad \text{dom } S = \{u \in H_0^1(\Omega) : \mathcal{L}u \in L^2(\Omega), \partial_{\mathcal{L}}u|_\omega = 0\},$$

of the Dirichlet operator A in $L^2(\Omega)$, which has no eigenvalues by Proposition 2.5 and we define the operator T in $L^2(\Omega)$ by

$$Tu = \mathcal{L}u, \quad \text{dom } T = \{u \in H^1(\Omega) : \mathcal{L}u \in L^2(\Omega), u|_{\partial\Omega} \in H_\omega^{1/2}(\partial\Omega)\}.$$

It follows immediately that the Dirichlet operator A is a restriction of T and that the spaces \mathcal{N}_λ coincide with $\ker(T - \lambda)$. In the first step of this proof we show that these spaces are dense in the spaces $\ker(S^* - \lambda)$. In the second step, which can also be found in a different form in [40], a selfadjoint restriction of S in the orthogonal complement of $\text{span}\{\mathcal{N}_\lambda : \lambda \in \mathcal{O} \cup \mathcal{O}^*\}$ is constructed. In the last step we then show that the spectrum of this selfadjoint operator is empty, which implies the assertion.

Step 1. In this step we show that $\mathcal{N}_\lambda = \ker(T - \lambda)$ are dense subspaces of $\ker(S^* - \lambda)$, $\lambda \in \mathcal{O} \cup \mathcal{O}^*$. For this we check first that $S = T^*$ holds. In fact, for $u \in \text{dom } S$ and $v \in \text{dom } T$ the second Green identity (2.7) together with $u|_{\partial\Omega} = 0$, $\partial_{\mathcal{L}}u|_\omega = 0$, and $\text{supp}(v|_{\partial\Omega}) \subset \omega$ implies

$$(Tv, u) = (v, \mathcal{L}u) + (v|_{\partial\Omega}, \partial_{\mathcal{L}}u|_{\partial\Omega}) - (\partial_{\mathcal{L}}v|_{\partial\Omega}, u|_{\partial\Omega}) = (v, \mathcal{L}u),$$

where we have used $(v|_{\partial\Omega}, \partial_{\mathcal{L}}u|_{\partial\Omega}) = (\partial_{\mathcal{L}}u|_\omega)(v|_{\partial\Omega}) = 0$; cf. (2.3). Hence $u \in \text{dom } T^*$ and $T^*u = \mathcal{L}u = Su$ by the definition of the adjoint operator. For the converse inclusion let $u \in \text{dom } T^*$. From $A \subseteq T$ we obtain $T^* \subseteq A^* = A$ and therefore $T^*u = \mathcal{L}u$ and $u \in \text{dom } A$. In particular, we have $u \in H^1(\Omega)$ and

$u|_{\partial\Omega} = 0$. It remains to show $\partial_{\mathcal{L}}u|_{\omega} = 0$. For $v \in \text{dom } T$ we have $\text{supp}(v|_{\partial\Omega}) \subset \omega$ and from (2.3) and (2.7) we obtain

$$(\partial_{\mathcal{L}}u|_{\omega})(v|_{\partial\Omega}) = (v|_{\partial\Omega}, \partial_{\mathcal{L}}u|_{\partial\Omega}) = (Tv, u) - (v, T^*u) + (\partial_{\mathcal{L}}v|_{\partial\Omega}, u|_{\partial\Omega}) = 0.$$

As $v|_{\partial\Omega}$ runs through $H_{\omega}^{1/2}(\partial\Omega)$ when v runs through $\text{dom } T$, it follows $\partial_{\mathcal{L}}u|_{\omega} = 0$, hence $u \in \text{dom } S$. We have shown $T^* = S$. This implies $\overline{T} = T^{**} = S^*$ and hence the spaces $\mathcal{N}_{\lambda} = \ker(T - \lambda)$ are dense in the spaces $\ker(S^* - \lambda)$.

Step 2. In this step we show that the Hilbert space

$$(2.10) \quad \mathcal{M} := (\text{span}\{\mathcal{N}_{\lambda} : \lambda \in \mathcal{O} \cup \mathcal{O}^*\})^{\perp}$$

is invariant for S and that $S_{\mathcal{M}} := S \upharpoonright (\mathcal{M} \cap \text{dom } S)$ is a selfadjoint operator in \mathcal{M} ; cf. [40]. Observe first that by step 1 we have

$$\mathcal{M} = (\text{span}\{\ker(S^* - \lambda) : \lambda \in \mathcal{O} \cup \mathcal{O}^*\})^{\perp} = \bigcap_{\lambda \in \mathcal{O} \cup \mathcal{O}^*} \text{ran}(S - \lambda).$$

Let $u \in \mathcal{M} \cap \text{dom } S$. Then $u \in \text{ran}(S - \lambda)$ for all $\lambda \in \mathcal{O} \cup \mathcal{O}^*$, i.e., for each $\lambda \in \mathcal{O} \cup \mathcal{O}^*$ there exists $u_{\lambda} \in \text{dom } S$ such that $(S - \lambda)u_{\lambda} = u$ holds. This implies

$$Su = S(S - \lambda)u_{\lambda} = (S - \lambda)Su_{\lambda} \in \text{ran}(S - \lambda)$$

for each $\lambda \in \mathcal{O} \cup \mathcal{O}^*$ and hence $Su \in \mathcal{M}$. Therefore, $S_{\mathcal{M}} = S \upharpoonright (\mathcal{M} \cap \text{dom } S)$ is an operator in \mathcal{M} and since S is symmetric we conclude that the restriction $S_{\mathcal{M}}$ is a symmetric operator in \mathcal{M} . For the selfadjointness of $S_{\mathcal{M}}$ in \mathcal{M} it is sufficient to show

$$(2.11) \quad \text{ran}(S_{\mathcal{M}} - \nu) = \mathcal{M} = \text{ran}(S_{\mathcal{M}} - \bar{\nu}) \quad \text{for some } \nu \in \mathcal{O}.$$

For this let $u \in \mathcal{M}$. We claim that $v := (S - \nu)^{-1}u \in \text{dom } S$ belongs to \mathcal{M} . In fact, for each $\lambda \in \mathcal{O} \cup \mathcal{O}^*$, $\lambda \neq \nu$, the function

$$v_{\lambda} := \frac{1}{\lambda - \nu} ((S - \lambda)^{-1} - (S - \nu)^{-1})u$$

satisfies $(S - \lambda)v_{\lambda} = v$ and hence $v \in \text{ran}(S - \lambda)$ for all $\lambda \in \mathcal{O} \cup \mathcal{O}^*$, $\lambda \neq \nu$. In order to check that also $v \in \text{ran}(S - \nu)$ holds, we choose a sequence $(\lambda_n)_n \subseteq \mathcal{O}$, $\lambda_n \neq \nu$, which converges to ν . As above it follows that $(S - \lambda_n)^{-1}u \in \text{ran}(S - \lambda)$ for all $\lambda \in \mathcal{O} \cup \mathcal{O}^*$, $\lambda \neq \lambda_n$, and, in particular, $(S - \lambda_n)^{-1}u \in \text{ran}(S - \nu)$. Since $S = T^*$ is a closed symmetric operator the estimates $\|(S - \nu)^{-1}\| \leq |\text{Im } \nu|^{-1}$ and $\|(S - \lambda_n)^{-1}\| \leq |\text{Im } \lambda_n|^{-1}$ hold, and hence

$$v - (S - \lambda_n)^{-1}u = (S - \nu)^{-1}u - (S - \lambda_n)^{-1}u = (\nu - \lambda_n)(S - \nu)^{-1}(S - \lambda_n)^{-1}u$$

implies

$$(2.12) \quad v = \lim_{n \rightarrow \infty} (S - \lambda_n)^{-1}u.$$

Since $(S - \lambda_n)^{-1}u \in \text{ran}(S - \nu)$ and $\text{ran}(S - \nu)$ is closed we conclude $v \in \text{ran}(S - \nu)$ from (2.12). We have shown $v \in \mathcal{M}$. Moreover, $(S_{\mathcal{M}} - \nu)v = (S - \nu)v = u$ and, hence, the first equality in (2.11) holds. The second equality in (2.11) follows in the same way. Therefore $S_{\mathcal{M}} = S \upharpoonright (\mathcal{M} \cap \text{dom } S)$ is a selfadjoint operator in \mathcal{M} .

Step 3. It follows from step 2 that the operator S can be written as the direct orthogonal sum $S_{\mathcal{M}} \oplus S_{\mathcal{M}^{\perp}}$ with respect to the decomposition $L^2(\Omega) = \mathcal{M} \oplus \mathcal{M}^{\perp}$, where $S_{\mathcal{M}}$ is a selfadjoint operator in \mathcal{M} and $S_{\mathcal{M}^{\perp}} = S \upharpoonright (\mathcal{M}^{\perp} \cap \text{dom } S)$ is a closed symmetric operator in \mathcal{M}^{\perp} . Since the spectrum of the Dirichlet operator A consists only of eigenvalues (see Proposition 2.1) the same holds for the spectrum of the selfadjoint part $S_{\mathcal{M}}$. Clearly, each eigenfunction of $S_{\mathcal{M}}$ is also an eigenfunction of $S = S_{\mathcal{M}} \oplus S_{\mathcal{M}^{\perp}}$, but by Proposition 2.5 this operator has no eigenfunctions.

Therefore, the spectrum of the selfadjoint operator $S_{\mathcal{M}}$ in \mathcal{M} is empty which implies $\mathcal{M} = \{0\}$, i.e.,

$$L^2(\Omega) = \mathcal{M}^\perp = \overline{\text{span}}\{\mathcal{N}_\lambda : \lambda \in \mathcal{O} \cup \mathcal{O}^*\}$$

by (2.10), where $\overline{\text{span}}$ denotes the closed linear span. This completes the proof of Lemma 2.6. \square

With these preparations we are ready to prove the uniqueness result Theorem 1.3.

Proof of Theorem 1.3. Let $\mathcal{L}_1, \mathcal{L}_2$ be elliptic differential expressions as in the theorem and let A_1, A_2 and M_1, M_2 be the corresponding selfadjoint Dirichlet operators and Dirichlet-to-Neumann maps, respectively. The associated Poisson operators from Definition 2.3 will be denoted by $\gamma_1(\lambda)$ and $\gamma_2(\lambda)$, $\lambda \in \rho(A_1) \cap \rho(A_2)$, respectively. Since $\mathcal{D} \subseteq \rho(A_1) \cap \rho(A_2)$ in Theorem 1.3 is a set with an accumulation point in the intersection of the domains of holomorphy of the functions M_1 and M_2 (see Lemma 2.4 (iv)), it follows that M_1 and M_2 coincide on ω , i.e.,

$$(2.13) \quad (M_1(\lambda)\varphi, \psi) = (M_2(\lambda)\varphi, \psi), \quad \varphi, \psi \in H_\omega^{1/2}(\partial\Omega),$$

holds for all $\lambda \in \mathbb{C} \setminus \mathbb{R}$, where $H_\omega^{1/2}(\partial\Omega)$ is the linear subspace of $H^{1/2}(\partial\Omega)$ which consists of functions with support in ω ; cf. (2.2).

Let in the following $\lambda, \mu \in \mathbb{C} \setminus \mathbb{R}$, $\lambda \neq \bar{\mu}$. Lemma 2.4 (iii) and (2.13) yield

$$(2.14) \quad \begin{aligned} (\gamma_1(\lambda)\varphi, \gamma_1(\mu)\psi) &= \frac{(M_1(\lambda)\varphi, \psi) - (\varphi, M_1(\mu)\psi)}{\bar{\mu} - \lambda} \\ &= \frac{(M_2(\lambda)\varphi, \psi) - (\varphi, M_2(\mu)\psi)}{\bar{\mu} - \lambda} = (\gamma_2(\lambda)\varphi, \gamma_2(\mu)\psi) \end{aligned}$$

for all $\varphi, \psi \in H_\omega^{1/2}(\partial\Omega)$. Next we define a linear mapping V in $L^2(\Omega)$ on the domain $\text{dom } V = \text{span}\{\gamma_1(\lambda)\varphi : \varphi \in H_\omega^{1/2}(\partial\Omega), \lambda \in \mathbb{C} \setminus \mathbb{R}\}$ by

$$V : \sum_{j=1}^{\ell} \gamma_1(\lambda_j)\varphi_j \mapsto \sum_{j=1}^{\ell} \gamma_2(\lambda_j)\varphi_j, \quad \lambda_j \in \mathbb{C} \setminus \mathbb{R}, \varphi_j \in H_\omega^{1/2}(\partial\Omega), 1 \leq j \leq \ell.$$

Formula (2.14) yields that V is a well-defined, isometric operator in $L^2(\Omega)$ with $\text{ran } V = \text{span}\{\gamma_2(\lambda)\varphi : \varphi \in H_\omega^{1/2}(\partial\Omega), \lambda \in \mathbb{C} \setminus \mathbb{R}\}$. By Lemma 2.6 the domain and range of V are both dense subspaces of $L^2(\Omega)$, and hence the closure U of V in $L^2(\Omega)$ is a unitary operator in $L^2(\Omega)$. Obviously $U\gamma_1(\lambda)\varphi = \gamma_2(\lambda)\varphi$ holds for all $\lambda \in \mathbb{C} \setminus \mathbb{R}$ and $\varphi \in H_\omega^{1/2}(\partial\Omega)$. With the help of Lemma 2.4 (ii) one computes for $\lambda \neq \mu$

$$U(A_1 - \lambda)^{-1}\gamma_1(\mu)\varphi = U \frac{\gamma_1(\lambda)\varphi - \gamma_1(\mu)\varphi}{\lambda - \mu} = (A_2 - \lambda)^{-1}U\gamma_1(\mu)\varphi$$

for $\varphi \in H_\omega^{1/2}(\partial\Omega)$. From Lemma 2.6 we then conclude

$$U(A_1 - \lambda)^{-1} = (A_2 - \lambda)^{-1}U$$

and therefore $UA_1u = A_2Uu$ for all $u \in \text{dom } A_1$, that is, A_1 and A_2 are unitarily equivalent. \square

The next proposition is the key ingredient in the proof of the reconstruction result Theorem 1.4. Here the residual $\text{Res}_\lambda M : H^{1/2}(\partial\Omega) \rightarrow H^{-1/2}(\partial\Omega)$ of the Dirichlet-to-Neumann map at a pole λ and a local version of it play the main role. Define the operator

$$(2.15) \quad \text{Res}_\lambda^\omega M : H_\omega^{1/2}(\partial\Omega) \rightarrow (H_\omega^{1/2}(\partial\Omega))', \quad \varphi \mapsto \text{Res}_\lambda^\omega M\varphi,$$

where $\text{Res}_\lambda^\omega M\varphi$ is the restriction of $\text{Res}_\lambda M\varphi \in H^{-1/2}(\partial\Omega)$ onto $H_\omega^{1/2}(\partial\Omega)$, i.e.,

$$(2.16) \quad (\text{Res}_\lambda^\omega M\varphi)(\psi) := (\text{Res}_\lambda M\varphi)(\psi) = (\psi, \text{Res}_\lambda M\varphi), \quad \psi \in H_\omega^{1/2}(\partial\Omega);$$

cf. the definition above Theorem 1.4 in the introduction. Note that in the special case $\omega = \partial\Omega$ the operators $\text{Res}_\mu M$ and $\text{Res}_\mu^\omega M$ coincide. It turns out that the Neumann trace operator on ω maps each eigenspace of the Dirichlet operator bijectively onto the range of the corresponding residual of the Dirichlet-to-Neumann map on ω .

Proposition 2.7. *For each λ the mapping*

$$\tau_\lambda : \ker(A - \lambda) \rightarrow \text{ran Res}_\lambda^\omega M, \quad u \mapsto \partial_{\mathcal{L}} u|_\omega,$$

is an isomorphism and, in particular,

$$\text{ran Res}_\lambda^\omega M = \{\psi : \text{there exists } u \in \ker(A - \lambda) \text{ such that } \partial_{\mathcal{L}} u|_\omega = \psi\}.$$

As an immediate consequence of this proposition we obtain the following statement which complements Lemma 2.4 (iv).

Corollary 2.8. *A point λ_0 is an eigenvalue of A if and only if λ_0 is a pole of the Dirichlet-to-Neumann map on ω . In this case the dimension of the eigenspace $\ker(A - \lambda_0)$ coincides with the dimension of the range of the operator $\text{Res}_{\lambda_0}^\omega M$.*

Proof of Proposition 2.7. Note first that if λ is not an eigenvalue of A , then the function M is holomorphic in λ by Lemma 2.4 (iv) and hence the residual $\text{Res}_\lambda M$ is zero. Therefore the assertion holds in this case.

In the following let $\lambda \in \mathbb{R}$ be an eigenvalue of A and let $\tau_\lambda u = \partial_{\mathcal{L}} u|_\omega$ be the Neumann trace operator on ω on $\ker(A - \lambda)$ defined in (2.3). For $u \in \ker(A - \lambda)$ and some fixed $\mu \in \mathbb{C} \setminus \mathbb{R}$ we obtain for all $\psi \in H_\omega^{1/2}(\partial\Omega)$ by Lemma 2.4 (i)

$$(2.17) \quad \begin{aligned} (\tau_\lambda u)(\psi) &= (\psi, \partial_{\mathcal{L}} u|_{\partial\Omega}) = (\psi, \partial_{\mathcal{L}} ((A - \bar{\mu})^{-1} Au - (A - \bar{\mu})^{-1} \bar{\mu} u)|_{\partial\Omega}) \\ &= (\psi, (\lambda - \bar{\mu}) \partial_{\mathcal{L}} ((A - \bar{\mu})^{-1} u)|_{\partial\Omega}) = (\psi, (\bar{\mu} - \lambda) \gamma(\mu)' u). \end{aligned}$$

Let us show that τ_λ is injective. Assume there exists $u \in \ker(A - \lambda)$, $u \neq 0$, such that $\tau_\lambda u = 0$, that is,

$$(2.18) \quad (\partial_{\mathcal{L}} ((A - \bar{\mu})^{-1} u)|_\omega)(\psi) = (\psi, \partial_{\mathcal{L}} ((A - \bar{\mu})^{-1} u)|_{\partial\Omega}) = 0$$

for all $\psi \in H_\omega^{1/2}(\partial\Omega)$ by (2.17). Since $u \neq 0$, also $(A - \bar{\mu})^{-1} u \neq 0$ holds, but

$$(A - \lambda)(A - \bar{\mu})^{-1} u = (A - \bar{\mu})^{-1} (A - \lambda) u = 0$$

implies that $(A - \bar{\mu})^{-1} u$ is an eigenfunction of A with vanishing Neumann trace on ω by (2.18). This contradicts Proposition 2.5. Hence τ_λ is injective.

Next we show that τ_λ maps onto $\text{ran Res}_\lambda^\omega M$, which by (2.17) is equivalent to

$$(2.19) \quad \text{ran Res}_\lambda^\omega M = \text{ran} (\gamma(\mu)' \upharpoonright \ker(A - \lambda)).$$

The inclusion \subseteq in (2.19) will be shown first. For this denote by E_λ the spectral projection onto the eigenspace $\ker(A - \lambda)$ of A . Let $\eta \in \mathbb{C} \setminus \mathbb{R}$, $\eta \neq \bar{\mu}$, and \mathcal{O} be an open ball centered in λ such that \mathcal{O} does not contain any further eigenvalues of A and $\mu, \eta \notin \bar{\mathcal{O}}$. Denote by Γ_λ the boundary of \mathcal{O} and write the projection E_λ as a Cauchy integral; cf. Appendix. For $\varphi, \psi \in H^{1/2}(\partial\Omega)$ we obtain with the help of Lemma 2.4 (ii) and (iii)

$$\begin{aligned} (E_\lambda \gamma(\eta) \varphi, \gamma(\mu) \psi) &= -\frac{1}{2\pi i} \int_{\Gamma_\lambda} ((A - \zeta)^{-1} \gamma(\eta) \varphi, \gamma(\mu) \psi) d\zeta \\ &= -\frac{1}{2\pi i} \int_{\Gamma_\lambda} \left(\frac{1}{\zeta - \eta} (\gamma(\zeta) \varphi, \gamma(\mu) \psi) - \frac{1}{\zeta - \eta} (\gamma(\eta) \varphi, \gamma(\mu) \psi) \right) d\zeta \\ &= \frac{1}{2\pi i} \int_{\Gamma_\lambda} \left(-\frac{(M(\zeta) \varphi, \psi)}{(\zeta - \eta)(\bar{\mu} - \zeta)} + \frac{(\varphi, M(\mu) \psi)}{(\bar{\mu} - \zeta)(\bar{\mu} - \eta)} + \frac{(M(\eta) \varphi, \psi)}{(\zeta - \eta)(\bar{\mu} - \eta)} \right) d\zeta; \end{aligned}$$

cf. [23, §I.1]. Note that the second and third fraction in the integral are holomorphic in \mathcal{O} as functions of ζ and hence can be neglected. In the remaining fraction, we develop the function $\zeta \mapsto M(\zeta)$ into a Laurent series at λ . Since this function has a pole of, at most, order one in λ , see Lemma 2.4 (iv) and the Appendix, we obtain

$$(E_\lambda \gamma(\eta)\varphi, \gamma(\mu)\psi) = \frac{1}{2\pi i} \int_{\Gamma_\lambda} \frac{(\operatorname{Res}_\lambda M\varphi, \psi)}{(\zeta - \lambda)(\eta - \zeta)(\bar{\mu} - \zeta)} d\zeta = \frac{(\operatorname{Res}_\lambda M\varphi, \psi)}{(\eta - \lambda)(\bar{\mu} - \lambda)},$$

where Cauchy's integral formula was used in the second equality. Therefore we have

$$(\psi, \operatorname{Res}_\lambda M\varphi) = (\bar{\eta} - \bar{\lambda})(\mu - \bar{\lambda})(\gamma(\mu)\psi, E_\lambda \gamma(\eta)\varphi)$$

and hence, in particular,

$$(2.20) \quad \operatorname{Res}_\lambda^\omega M\varphi = (\bar{\eta} - \bar{\lambda})(\mu - \bar{\lambda})\gamma(\mu)'E_\lambda \gamma(\eta)\varphi, \quad \varphi \in H_\omega^{1/2}(\partial\Omega);$$

cf. (2.15) and (2.16). This shows the inclusion \subseteq in (2.19).

In order to show the inclusion \supseteq in (2.19) let $u \in \ker(A - \lambda)$ and $\varepsilon > 0$. Since $\gamma(\mu)'E_\lambda$ is bounded by Lemma 2.4 (i), there exists $\delta > 0$ such that $\|u - v\| < \delta$ implies $\|\gamma(\mu)'E_\lambda u - \gamma(\mu)'E_\lambda v\| < \varepsilon$. Note that $E_\lambda u = u$ as $u \in \ker(A - \lambda)$. By Lemma 2.6 we find $\ell \in \mathbb{N}$, $\eta_j \in \mathbb{C} \setminus \mathbb{R}$, and $\varphi_j \in H_\omega^{1/2}(\partial\Omega)$, $1 \leq j \leq \ell$, such that

$$\left\| u - \sum_{j=1}^{\ell} \gamma(\eta_j)\varphi_j \right\| < \delta.$$

Then we have

$$(2.21) \quad \left\| \gamma(\mu)'E_\lambda u - \gamma(\mu)'E_\lambda \sum_{j=1}^{\ell} \gamma(\eta_j)\varphi_j \right\| < \varepsilon$$

and from (2.20) and (2.21) we conclude

$$\left\| \gamma(\mu)'u - \sum_{j=1}^{\ell} \frac{\operatorname{Res}_\lambda^\omega M\varphi_j}{(\bar{\lambda} - \bar{\eta}_j)(\bar{\lambda} - \mu)} \right\| < \varepsilon,$$

that is, $\gamma(\mu)'u \in \overline{\operatorname{ran} \operatorname{Res}_\lambda^\omega M}$. The fact that $\ker(A - \lambda)$ is finite-dimensional (see Proposition 2.1) shows (2.19). Thus τ_λ is bijective and maps the finite-dimensional space $\ker(A - \lambda)$ onto $\operatorname{ran} \operatorname{Res}_\lambda^\omega M$, i.e., τ_λ is an isomorphism. \square

Theorem 1.4 is now essentially a consequence of Proposition 2.7.

Proof of Theorem 1.4. For the statement on the poles of M see Corollary 2.8. In order to prove the representation of A let us denote by $(\lambda_k)_k$ the sequence of eigenvalues of A and by τ_k the restriction of the Neumann trace operator onto the eigenspace $\ker(A - \lambda_k)$ from Proposition 2.7,

$$\tau_k = \tau_{\lambda_k} : \ker(A - \lambda_k) \rightarrow \operatorname{ran} \operatorname{Res}_{\lambda_k}^\omega M, \quad u \mapsto \partial_{\mathcal{L}} u|_\omega.$$

Since by Proposition 2.7 τ_k is an isomorphism, the formula

$$\ker(A - \lambda_k) = \operatorname{ran} (\tau_k^{-1} \circ \operatorname{Res}_{\lambda_k}^\omega M)$$

for the eigenspace corresponding to the eigenvalue λ_k follows immediately. In particular, we can choose $\varphi_1^{(k)}, \dots, \varphi_{n(k)}^{(k)} \in H_\omega^{1/2}(\partial\Omega)$, $n(k) = \dim \ker(A - \lambda_k)$, such that the functions

$$e_i^{(k)} = \tau_k^{-1} (\operatorname{Res}_{\lambda_k}^\omega M) \varphi_i^{(k)}, \quad 1 \leq i \leq n(k),$$

form an orthonormal basis in $\ker(A - \lambda_k)$. Then the orthogonal projection E_k in $L^2(\Omega)$ onto $\ker(A - \lambda_k)$ is given by

$$E_k u = \sum_{i=1}^{n(k)} (u, e_i^{(k)}) e_i^{(k)}, \quad u \in L^2(\Omega).$$

Since the spectrum of A consists only of eigenvalues with finite-dimensional eigenspaces we conclude that A can be represented in the form

$$Au = \sum_{k=1}^{\infty} \lambda_k \sum_{i=1}^{n(k)} (u, e_i^{(k)}) e_i^{(k)}, \quad u \in \text{dom } A.$$

□

Remark 2.9. Since M is a holomorphic operator function whose values are bounded operators from $H^{1/2}(\partial\Omega) \rightarrow H^{-1/2}(\partial\Omega)$ it is sufficient to assume knowledge of $M(\lambda)\varphi$ in Theorem 1.3 and Theorem 1.4 for φ in an arbitrary dense subspace of $H^{1/2}(\partial\Omega)$ and λ in a discrete set of points \mathcal{D} with an accumulation point in $\rho(A)$. In particular, since the spectrum of A is discrete, it is possible to choose \mathcal{D} as a sequence in any nonempty, open subset of \mathbb{R} .

APPENDIX

In this appendix we recall some definitions and basic facts on (unbounded) operators in Hilbert spaces and Banach space-valued mappings which can be found in, e.g., the monographs [1, 18, 25, 35, 55].

Linear operators in Hilbert spaces. Let \mathcal{H} and \mathcal{K} be Hilbert spaces with scalar products $(\cdot, \cdot)_{\mathcal{H}}$ and $(\cdot, \cdot)_{\mathcal{K}}$, and let T be a linear operator from \mathcal{H} to \mathcal{K} . We write $\text{dom } T$, $\ker T$ and $\text{ran } T$ for the domain, kernel and range of T , respectively. The operator T is said to be *closed* if its graph is a closed subspace of $\mathcal{H} \times \mathcal{K}$. If $\text{dom } T$ is dense in \mathcal{H} , then the *adjoint operator* T^* is defined by $T^*g := \tilde{g}$, where

$$\text{dom } T^* = \{g \in \mathcal{K} : \text{exists } \tilde{g} \in \mathcal{H} \text{ such that } (Tf, g)_{\mathcal{K}} = (f, \tilde{g})_{\mathcal{H}} \text{ for all } f \in \text{dom } T\}.$$

Observe that T^* is a closed linear operator from \mathcal{K} to \mathcal{H} . Moreover, $\text{dom } T^*$ is dense in \mathcal{K} if and only if the closure \overline{T} of (the graph of) T is an operator, and in this case T^{**} coincides with \overline{T} .

Let A be a linear operator in \mathcal{H} . Then A is said to be *symmetric* if the relation $(Af, g)_{\mathcal{H}} = (f, Ag)_{\mathcal{H}}$ holds for all $f, g \in \text{dom } A$. If A is densely defined, then A is symmetric if and only if the adjoint operator A^* is an extension of A , that is, $\text{dom } A \subseteq \text{dom } A^*$ and $A^*f = Af$ holds for all $f \in \text{dom } A$; in short $A \subseteq A^*$. If $A = A^*$ holds, then the operator A is called *selfadjoint*. Recall that a symmetric operator A is selfadjoint if and only if $\text{ran}(A - \lambda_{\pm}) = \mathcal{H}$ holds for some $\lambda_{\pm} \in \mathbb{C}^{\pm}$ and that for a closed symmetric operator A and $\lambda \in \mathbb{C} \setminus \mathbb{R}$ the operator $(A - \lambda)^{-1}$ defined on the closed subspace $\text{ran}(A - \lambda)$ is bounded by $|\text{Im } \lambda|^{-1}$.

Let S be a closed linear operator in \mathcal{H} . The *resolvent set* $\rho(S)$ consists of all $\lambda \in \mathbb{C}$ such that $(S - \lambda)^{-1}$ is a bounded operator defined on \mathcal{H} . Note that the set $\rho(S)$ is open in \mathbb{C} . The *spectrum* $\sigma(S)$ of S is the complement of $\rho(S)$ in \mathbb{C} , in particular, $\sigma(S)$ contains the eigenvalues of S , i.e., the points $\lambda \in \mathbb{C}$ such that $\ker(S - \lambda) \neq \{0\}$ holds. Recall that for a symmetric operator A in \mathcal{H} the eigenvalues are real and that for a selfadjoint operator A the whole spectrum $\sigma(A)$ is contained in \mathbb{R} .

Holomorphic functions with values in Banach spaces. Let X be a Banach space and let $D \subseteq \mathbb{C}$ be an open set. If the function $m : D \rightarrow X$ is holomorphic on D and $\mu \in \mathbb{C}$ is a pole of m , then the *residual* of m at μ is

$$\operatorname{Res}_\mu m = \frac{1}{2\pi i} \int_{\Gamma_\mu} m(\zeta) d\zeta,$$

where Γ_μ denotes a closed Jordan curve with interior \mathcal{O} which contains μ , such that $\mathcal{O} \setminus \{\mu\} \subseteq D$. Equivalently, $\operatorname{Res}_\mu m$ is the first coefficient of negative order in the Laurent series expansion of m at μ . If A is a selfadjoint operator in a Hilbert space \mathcal{H} , then $\lambda \mapsto R_A(\lambda) = (A - \lambda)^{-1}$ is a holomorphic function on $\rho(A)$ with values in the space of bounded linear operators in \mathcal{H} . Here each isolated point μ in $\sigma(A)$ is a pole of R_A of order one and the orthogonal projection E_μ in \mathcal{H} onto the eigenspace $\ker(A - \mu)$ is

$$E_\mu = -\operatorname{Res}_\mu R_A = -\frac{1}{2\pi i} \int_{\Gamma_\mu} (A - \zeta)^{-1} d\zeta.$$

REFERENCES

- [1] N. I. Akhiezer and I. M. Glazman, *Theory of Linear Operators in Hilbert Space*, Dover publications, 1993.
- [2] D. Alpay and J. Behrndt, *Generalized Q -functions and Dirichlet-to-Neumann maps for elliptic differential operators*, *J. Funct. Anal.* 257 (2009), 1666–1694.
- [3] W. O. Amrein and D. B. Pearson, *M -operators: a generalisation of Weyl–Titchmarsh theory*, *J. Comput. Appl. Math.* 171 (2004), 1–26.
- [4] W. Arendt and A. F. M. ter Elst, *The Dirichlet-to-Neumann operator on rough domains*, Preprint, arXiv:1005.0875.
- [5] W. Arendt and R. Mazzeo, *Spectral properties of the Dirichlet-to-Neumann operator on Lipschitz domains*, *Ulmer Seminare* 12 (2007), 23–38.
- [6] Yu. Arlinskiĭ, *Abstract boundary conditions for maximal sectorial extensions of sectorial operators*, *Math. Nachr.* 209 (2000), 5–36.
- [7] N. Aronszajn, *A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order*, *J. Math. Pures Appl.* 36 (1957), 235–249.
- [8] N. Aronszajn, A. Krzywicki, and J. Szarski, *A unique continuation theorem for exterior differential forms on Riemannian manifolds*, *Ark. Mat.* 4 (1962), 417–453.
- [9] K. Astala, M. Lassas, and L. Päivärinta, *Calderón’s inverse problem for anisotropic conductivity in the plane*, *Comm. Partial Differential Equations* 30 (2005), 207–224.
- [10] K. Astala and L. Päivärinta, *Calderón’s inverse conductivity problem in the plane*, *Ann. of Math.* (2) 163 (2006), 265–299.
- [11] J. Behrndt and M. Langer, *Boundary value problems for elliptic partial differential operators on bounded domains*, *J. Funct. Anal.* 243 (2007), 536–565.
- [12] J. Behrndt and M. Langer, *Elliptic operators, Dirichlet-to-Neumann maps and quasi boundary triples*, submitted.
- [13] M. Brown, G. Grubb, and I. Wood, *M -functions for closed extensions of adjoint pairs of operators with applications to elliptic boundary problems*, *Math. Nachr.* 282 (2009), 314–347.
- [14] M. Brown, J. Hinchcliffe, M. Marletta, S. Naboko and I. Wood, *The abstract Titchmarsh–Weyl M -function for adjoint operator pairs and its relation to the spectrum*, *Integral Equations Operator Theory* 63 (2009), 297–320.
- [15] M. Brown, M. Marletta, S. Naboko, and I. Wood, *Boundary triplets and M -functions for non-selfadjoint operators, with applications to elliptic PDEs and block operator matrices*, *J. Lond. Math. Soc.* (2) 77 (2008), 700–718.
- [16] A. L. Bukhgeim and G. Uhlmann, *Recovering a potential from partial Cauchy data*, *Comm. Partial Differential Equations* 27 (2002), 653–668.
- [17] A. P. Calderón, *On an inverse boundary value problem*, *Seminar on Numerical Analysis and its Applications to Continuum Physics* (Rio de Janeiro, 1980), pp. 65–73, *Soc. Brasil. Mat.*, Rio de Janeiro, 1980.
- [18] J. B. Conway, *A Course in Operator Theory*, *Grad. Stud. Math.* 21, Amer. Math. Soc., Providence, 1991.
- [19] V. A. Derkach, S. Hassi, M. M. Malamud, and H. S. V. de Snoo, *Boundary relations and their Weyl families*, *Trans. Amer. Math. Soc.* 358 (2006), 5351–5400.
- [20] V. A. Derkach and M. M. Malamud, *Generalized resolvents and the boundary value problems for Hermitian operators with gaps*, *J. Funct. Anal.* 95 (1991), 1–95.

- [21] V. A. Derkach and M. M. Malamud, *The extension theory of Hermitian operators and the moment problem*, J. Math. Sciences 73 (1995), 141–242.
- [22] A. Dijksma, H. Langer, and H. de Snoo, *Representations of holomorphic operator functions by means of resolvents of unitary or selfadjoint operators in Krein spaces*, Oper. Theory Adv. Appl. 24 (1987), 123–143.
- [23] A. Dijksma, H. Langer, and H. de Snoo, *Eigenvalues and pole functions of Hamiltonian systems with eigenvalue depending boundary conditions*, Math. Nachr. 161 (1993), 107–154.
- [24] D. Dos Santos Ferreira, C. E. Kenig, M. Salo, and G. Uhlmann, *Limiting Carleman weights and anisotropic inverse problems*, Invent. Math. 178 (2009), 119–171.
- [25] N. Dunford and J. T. Schwartz, *Linear Operators I*, Wiley, New York, 1958.
- [26] D. E. Edmunds and W. D. Evans, *Spectral Theory and Differential Operators*, Clarendon Press, Oxford, 1987.
- [27] W. N. Everitt and L. Markus, *Elliptic partial differential operators and symplectic algebra*, Mem. Amer. Math. Soc. 162 (2003), no. 770, 111 pp.
- [28] F. Gesztesy and M. Mitrea, *Generalized Robin boundary conditions, Robin-to-Dirichlet maps, and Krein-type resolvent formulas for Schrödinger operators on bounded Lipschitz domains*, Proc. Sympos. Pure Math. 79, 105–173, Amer. Math. Soc., 2008.
- [29] F. Gesztesy and M. Mitrea, *Nonlocal Robin Laplacians and some remarks on a paper by Filonov on eigenvalue inequalities*, J. Differential Equations 247 (2009), 2871–2896.
- [30] F. Gesztesy and M. Mitrea, *Robin-to-Robin maps and Krein-type resolvent formulas for Schrödinger operators on bounded Lipschitz domains*, Oper. Theory Adv. Appl. 191 (2009), 81–113.
- [31] F. Gesztesy, M. Mitrea, and M. Zinchenko, *Multi-dimensional versions of a determinant formula due to Jost and Pais*, Rep. Math. Phys. 59 (2007), 365–377.
- [32] F. Gesztesy, M. Mitrea, and M. Zinchenko, *Variations on a theme of Jost and Pais*, J. Funct. Anal. 253 (2007), 399–448.
- [33] G. Grubb, *A characterization of the non-local boundary value problems associated with an elliptic operator*, Ann. Scuola Norm. Sup. Pisa (3) 22 (1968), 425–513.
- [34] G. Grubb, *Distributions and Operators*, Graduate Texts in Mathematics 252, Springer, New York, 2009.
- [35] H. Heuser, *Functional Analysis*, John Wiley & Sons, Ltd., Chichester, 1982.
- [36] L. Hörmander, *Uniqueness theorems for second order elliptic differential equations*, Comm. Partial Differential Equations 8 (1) (1983), 21–64.
- [37] P. Jonas, *Operator representations of definitizable functions*, Ann. Acad. Sci. Fenn. Math. 25 (2000), 41–72.
- [38] A. Katchalov, Y. Kurylev, and M. Lassas, *Inverse Boundary Spectral Problems*, Chapman and Hall/CRC Monogr. and Surv. in Pure and Appl. Math., 123, Chapman and Hall/CRC, 2001.
- [39] C. E. Kenig, J. Sjöstrand, and G. Uhlmann, *The Calderón problem with partial data*, Ann. of Math. (2) 165 (2007), 567–591.
- [40] M. G. Krein, *Basic propositions of the theory of representation of Hermitian operators with deficiency index (m, m)* , Ukrain. Mat. Zh. 1 (1949), 3–66.
- [41] H. Langer and B. Textorius, *On generalized resolvents and Q -functions of symmetric linear relations (subspaces) in Hilbert space*, Pacific J. Math. 72 (1977), 135–165.
- [42] M. Lassas, M. Taylor, and G. Uhlmann, *The Dirichlet-to-Neumann map for complete Riemannian manifolds with boundary*, Comm. Anal. Geom. 11 (2003), 207–221.
- [43] M. Lassas and G. Uhlmann, *Determining Riemannian manifold from boundary measurements*, Ann. Sci. Ec. Norm. Super. 34 (2001), 771–787.
- [44] J. Lee and G. Uhlmann, *Determining anisotropic real-analytic conductivities by boundary measurements*, Comm. Pure Appl. Math. 42 (1989), 1097–1112.
- [45] J. L. Lions and E. Magenes, *Non-Homogenous Boundary Value Problems and Applications*, Volume 1, Springer, 1972.
- [46] M. M. Malamud, *Spectral theory of elliptic operators in exterior domains*, Russ. J. Math. Phys. 17 (2010), 96–125.
- [47] J. Malinen and O. J. Staffans, *Impedance passive and conservative boundary control systems*, Complex Anal. Oper. Theory 1 (2007), 279–300.
- [48] W. McLean, *Strongly Elliptic Systems and Boundary Integral Equations*, Cambridge University Press, 2000.
- [49] A. Nachman, *Reconstructions from boundary measurements*, Ann. of Math. (2) 128 (1988), 531–576.
- [50] A. Nachman, *Global uniqueness for a two-dimensional inverse boundary value problem*, Ann. of Math. (2) 143 (1996), 71–96.

- [51] A. Nachman and B. Street, *Reconstruction in the Calderón problem with partial data*, Comm. Partial Differential Equations 35 (2010), 375–390.
- [52] A. Nachman, J. Sylvester, and G. Uhlmann, *An n -dimensional Borg-Levinson theorem*, Comm. Math. Phys. 115 (1988), 595–605.
- [53] A. Posilicano, *Self-adjoint extensions of restrictions*, Oper. Matrices 2 (2008), 483–506.
- [54] O. Post, *First-order operators and boundary triples*, Russ. J. Math. Phys. 14 (2007), 482–492.
- [55] M. Reed and B. Simon, *Methods of Modern Mathematical Physics. I. Functional Analysis*, Academic Press, New York, 1980.
- [56] V. Ryzhov, *A general boundary value problem and its Weyl function*, Opuscula Math. 27 (2007), 305–331.
- [57] V. Ryzhov, *Weyl-Titchmarsh function of an abstract boundary value problem, operator collocations, and linear systems with boundary control*, Complex Anal. Oper. Theory 3 (2009), 289–322.
- [58] Z. Sun and G. Uhlmann, *Anisotropic inverse problems in two dimensions*, Inverse Problems 19 (2003), 1001–1010.
- [59] J. Sylvester, *An anisotropic inverse boundary value problem*, Comm. Pure Appl. Math. 43 (1990), 201–232.
- [60] J. Sylvester and G. Uhlmann, *A global uniqueness theorem for an inverse boundary value problem*, Ann. of Math. (2) 125 (1987), 153–169.
- [61] J. Sylvester and G. Uhlmann, *Inverse problems in anisotropic media*, Contemp. Math. 122 (1991), 105–117.
- [62] T. H. Wolff, *Recent work on sharp estimates in second-order elliptic unique continuation problems*, J. Geom. Anal. 3 (1993), 621–650.

INSTITUT FÜR MATHEMATIK, MA 6-4, TECHNISCHE UNIVERSITÄT BERLIN, STRASSE DES 17. JUNI
136, 10623 BERLIN, DEUTSCHLAND
E-mail address: behrndt@math.tu-berlin.de and rohleder@math.tu-berlin.de