

DDH-like Assumptions Based on Extension Rings

Ronald Cramer*, Ivan Damgård†, Eike Kiltz‡, Sarah Zakarias†, Angela Zottarel†

* CWI and Leiden University,
† Aarhus University and CTIC,
‡ RU Bochum University

Abstract. We introduce and study a new type of DDH-like assumptions based on groups of prime order q . Whereas standard DDH is based on encoding elements of \mathbb{F}_q “in the exponent” of elements in the group, we ask what happens if instead we put in the exponent elements of the extension ring $R_f = \mathbb{F}_q[X]/(f)$ where f can be any degree- d polynomial. We show that solving the decision problem that follows naturally reduces to the case where f is irreducible. This variant is called the d -DDH problem, where 1-DDH is standard DDH. Essentially any known cryptographic construction based on DDH can be immediately generalized to use instead d -DDH, and we show in the generic group model that d -DDH is harder than DDH. This means that virtually any application of DDH can now be realized with the same (amortized) efficiency, but under a potentially weaker assumption. On the negative side, we also show that d -DDH, just like DDH, is easy in bilinear groups. This motivates our suggestion of a different type of assumption, the d -vector DDH problems (d -VDDH), which are based on $f(X) = X^d$, but with a twist to avoid the problems with reducible polynomials. We show in the generic group model that d -VDDH is hard in bilinear groups and that in fact the problems become harder with increasing d and hence form an infinite hierarchy. We show that hardness of d -VDDH implies CCA-secure encryption, efficient Naor-Reingold style pseudorandom functions, and auxiliary input secure encryption, a strong form of leakage resilience. This can be seen as an alternative to the known family of k -linear assumptions.

Keywords: DDH, Public Key Encryption, PRF, Leakage Resilient Encryption

1 Introduction

The computational Diffie-Hellman assumption (proposed by Diffie and Hellman in [DH76]), says that if one chooses random g in a finite group \mathbb{G} and random exponents a, b , then given g, g^a, g^b it is hard to compute g^{ab} . The assumption was introduced as basis for the well-known Diffie-Hellman key exchange, but to get efficient public-key encryption one needs the stronger Decisional Diffie-Hellman assumption (DDH, studied by Naor and Reingold in [NR97]). It says that given g, g^a, g^b , the group element g^{ab} is pseudorandom, i.e., cannot be efficiently distinguished from g^c for a random c . In some groups, the DDH assumption is clearly false, but it is widely conjectured to hold when \mathbb{G} is, for instance, a large prime order subgroup of \mathbb{F}_p^* .

DDH has been used as the basis for a very wide range of cryptographic primitives, such as pseudorandom functions (PRF) [NR97], hash-proof systems and CCA-secure public-key Encryption [CS98], leakage resilient cryptography (in particular, auxiliary input security [DGK⁺10]), and circular secure encryption [BHHO08].

Clearly, the DDH-assumption is false in groups where discrete log is easy. However, in so-called bilinear groups (that can be constructed from certain elliptic curves), DDH is easy, even though discrete log is still conjectured to be hard. This has led to a large body of research on weaker variants of the DDH assumption of which some can be used in bilinear groups. A well-known example is a family of assumptions called the k -linear assumptions (where $k = 1$ is simply

DDH) [BBS04, HK07, Kil07, Sha07]. In the generic group model, these assumptions are shown to become progressively weaker for increasing k . Most of the primitives we mentioned above can be based on the k -linear assumptions, but with a loss of efficiency that increases with k . Several other assumptions related to discrete log in bilinear groups have been proposed, for instance to support variants of identity-based encryption or signature schemes. These assumptions sometimes seem to be conceived specifically to support a particular scheme. While this is not necessarily a bad idea, we believe there is good motivation to complement this by taking the opposite approach, namely start from a large class of assumptions and study which of them are useful and plausible.

In this paper we initiate such a study of a new family of assumptions that form natural generalizations of DDH in prime order groups: if \mathbb{G} has prime order q , and we fix a generator h , then an element $g \in \mathbb{G}$ “encodes” an element $a \in \mathbb{F}_q$ namely the a for which $g = h^a$. Intuitively we can think of a copy of \mathbb{F}_q sitting in the exponent, and we can add field elements by multiplying in \mathbb{G} , and multiply by known constants by doing exponentiation. However, if CDH is hard, we cannot do general multiplication, i.e., compute g^{ab} from g^a, g^b . If DDH is hard, we cannot even distinguish the correct result from random. Now, let us instead consider the extension ring $R_f = \mathbb{F}_q[X]/(f)$ where f is a degree- d polynomial. It is well-known that an element $\mathbf{w} \in R_f$ can be represented as a vector $(w_0, \dots, w_{d-1}) \in \mathbb{F}_q^d$. We can therefore represent \mathbf{w} by a tuple of d group elements $(h^{w_0}, \dots, h^{w_{d-1}}) \in \mathbb{G}^d$. Addition in R_f now becomes multiplication in \mathbb{G}^d , and multiplication by a known constant $\mathbf{a} \in R_f$ can be done (as we shall see) by applying a linear function in the exponent. This is simply because in R_f multiplication by a constant \mathbf{a} acts as a linear mapping on the vector (w_0, \dots, w_{d-1}) . More details will be given below, but the essence is that if we set $\mathbf{g} = (h^{w_0}, \dots, h^{w_{d-1}}) \in \mathbb{G}^d$ and take any $\mathbf{a} \in R_f$, we can define $\mathbf{g}^{\mathbf{a}}$ in a completely natural way, namely as the d -tuple of elements in \mathbb{G} that represent $\mathbf{w}\mathbf{a}$. This leads to defining the f -DDH problem as follows: given $(\mathbf{g}, \mathbf{g}^{\mathbf{a}}, \mathbf{g}^{\mathbf{b}}, \mathbf{g}^{\mathbf{c}})$, where $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{F}_{q^d}, \mathbf{g} \in \mathbb{G}^d$, decide if \mathbf{c} is random or $\mathbf{c} = \mathbf{ab}$.

We show that f -DDH and f' -DDH are equivalent whenever R_f is isomorphic to $R_{f'}$, and also that f -DDH is no harder than f_0 -DDH where f_0 is an irreducible factor in f . So it is natural to consider only the case where f is irreducible of degree d , in which case $R_f = \mathbb{F}_{q^d}$. This variant is called d -DDH. We show that if d_2 divides d_1 , so that $\mathbb{F}_{q^{d_2}}$ is a subfield of $\mathbb{F}_{q^{d_1}}$, then d_1 -DDH is at least as hard as d_2 -DDH. We also show in the generic group model that d -DDH for $d > 1$ is in fact *harder* than DDH. The proof of this is interesting from a technical point of view: proofs in the generic group model usually work by arguing that the adversary fails because he cannot compute expressions “in the exponent” of sufficiently high degree. This approach completely fails in our case, instead we have to solve a much harder task, namely we show that the ability to verify whether certain degree-2 equations are satisfied, does not allow verification of a different class of degree-2 equations.

The result also has a number of interesting consequences: since d -DDH is syntactically so similar to DDH, it can be directly plugged into virtually any cryptographic construction based on DDH, and security would now be reducible to d -DDH. An important point is that using d -DDH, you can produce several pseudorandom elements at essentially the same cost per element as with DDH. Therefore all applications we are aware of can now be realized under the weaker d -DDH assumption, but - in an amortized sense - with the same efficiency. We explain this in more detail in Section 5.

Note that the d -DDH assumption should not be confused with the previously known k -DDH assumption which is completely different and is *stronger* than DDH (see, e.g., [BB04, DY05, BMR10] for details on and applications of this assumption).

On the negative side, however, we also show that d -DDH, just like DDH, is easy in bilinear groups. This fact motivates our suggestion of the following alternative family of problems: we observe that by omitting some group elements from an instance of f -DDH, one can obtain a problem that is hard, even if f is reducible. Based on this, we propose the d -vector DDH (d -VDDH) assumptions, based on $f(X) = X^d$. We show in the generic group model that the d -VDDH assumption holds even in bilinear groups. In fact, it holds even given a d -linear map, which can be thought of as an oracle allowing the adversary to compute expressions of degree d in the exponent. This means that the d -VDDH assumptions become progressively weaker for increasing d . We show that the d -VDDH assumption implies CCA-secure encryption and efficient Naor-Reingold style pseudorandom functions. We also construct another cryptosystem based on the d -VDDH assumption, very similar to the BHHO scheme [BHHO08]. We show that this scheme is auxiliary input secure, a strong form of leakage resilience where full information on the secret key can be leaked, as long as the key remains hard to compute.

The family of d -VDDH assumptions can therefore be seen as an alternative to the (incomparable) family of k -linear assumptions.

A final related work that should be mentioned is [HYZX08] in which an assumption called EDDH is proposed, which is our 2-DDH assumption. This is the only prior work we know of that mentions a DDH variant based on ring extensions. It is claimed in [HYZX08] that DDH reduces to EDDH and that in the generic group model EDDH is hard, even in bilinear groups. The first result is correct, but we could not verify the proof. In this paper, we give a different proof of a more general statement. The second claim is false, and is refuted by our result that d -DDH for any d is easy in bilinear groups.

2 Preliminaries

2.1 Notation

If S is a set, we write $x \leftarrow S$ meaning that x is sampled uniformly from S . If $\mathbf{x} \in \mathbb{F}_q^m$ is a vector, we write $x[i]$ for the i th entry of \mathbf{x} . We say that a function $f : \mathbb{N} \rightarrow \mathbb{R}$ is negligible if, for every polynomial p , there exists an integer $n_p \in \mathbb{N}$ such that $f(n) < 1/p(n)$ for every $n > n_p$. If X and Y are two random variables, we say that X and Y are computationally indistinguishable ($X \stackrel{c}{\approx} Y$) if their computational distance is negligible. Furthermore, throughout the paper, vectors are denoted by bold lowercase letters.

A d -linear map $e : \mathbb{G}^d \rightarrow \mathbb{G}_T$ is an efficiently computable map such that $e(g, \dots, g) \neq 1$ and $e(g_1^{a_1}, \dots, g_d^{a_d}) = e(g_1, \dots, g_d)^{\prod a_i}$, for all g_i in \mathbb{G} and for all a_i in \mathbb{F}_q . A d -linear group \mathbb{G} is a group \mathbb{G} together with a d -linear map.

3 Extension Rings and DDH

We consider here a finite field \mathbb{F}_q of prime order q and its extension with a polynomial f of degree d . By this we obtain the ring $R_f = \mathbb{F}_q[X]/(f)$, where an element \mathbf{v} can be written as $v_0 + \dots + v_{d-1}X^{d-1} + (f)$. However, we can also represent \mathbf{v} by the matrix $V = v_0\mathbf{I}_d + v_1A_f + \dots + v_{d-1}A_f^{d-1}$, where \mathbf{I}_d is the d -dimensional identity matrix and A_f is the so-called companion matrix of f . The companion matrix of a monic polynomial $f = X^d + \alpha_{d-1}X^{d-1} + \dots + \alpha_1X + \alpha_0$ is given by the

$d \times d$ matrix

$$A_f = \begin{pmatrix} 0 & 0 & \cdots & 0 & -\alpha_0 \\ 1 & 0 & \cdots & 0 & -\alpha_1 \\ 0 & 1 & \cdots & 0 & -\alpha_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & -\alpha_{d-1} \end{pmatrix}.$$

Action of matrices on \mathbb{G}^d Given a group \mathbb{G} of order q and a tuple of elements $\mathbf{g} = (g_0, \dots, g_{d-1}) \in \mathbb{G}^d$, any matrix $M = (m_{ij})$ of dimension $n \times d$ defines a mapping $\mathbb{G}^d \rightarrow \mathbb{G}^n$ as follows:

$$\mathbf{g}^M := \left(\prod_j g_j^{m_{1j}}, \dots, \prod_j g_j^{m_{nj}} \right). \quad (1)$$

In particular this means that R_f can act on \mathbb{G}^d : we write the element $\mathbf{v} \in R_f$ in its matrix representation V and compute $\mathbf{g}^{\mathbf{v}} := \mathbf{g}^V$ as above. It is straightforward to verify that this map behaves according to the standard rules for exponentiation:

$$(\mathbf{g}^{\mathbf{a}})^{\mathbf{b}} = \mathbf{g}^{\mathbf{ab}}, \quad \mathbf{g}^{\mathbf{a}} \mathbf{g}^{\mathbf{b}} = \mathbf{g}^{\mathbf{a+b}}.$$

Note that this action can also be understood as implementing a product in R_f in a slightly different way: if we choose a generator h of \mathbb{G} , then we can write any \mathbf{g} as $(g_0, \dots, g_{d-1}) = (h^{w_0}, \dots, h^{w_{d-1}})$. Once we fix h , we can therefore think of \mathbf{g} as representing an element \mathbf{w} in R_f , namely $\mathbf{w} = w_{d-1}X^{d-1} + \dots + w_0 + (f)$. We will write this as $\mathbf{g} = h(\mathbf{w})$. It now turns out that we have

$$\mathbf{g}^{\mathbf{v}} = h(\mathbf{w})^{\mathbf{v}} = h(\mathbf{wv}).$$

This follows because we can think of R_f as a d -dimensional vector space over \mathbb{F}_q . In that interpretation, multiplication by \mathbf{v} is a linear mapping which has a matrix, namely V . Since the action $\mathbf{g}^{\mathbf{v}}$ is defined to be multiplication by V “in the exponent”, it follows that by computing $\mathbf{g}^{\mathbf{v}} = (h^{w_0}, \dots, h^{w_{d-1}})^{\mathbf{v}}$, we are in fact multiplying \mathbf{w} by \mathbf{v} .

3.1 The f -DDH Problem

Given the above, we can now define an new variant of the DDH problem:

Definition 1 (The f -DDH Problem). Let f be a d -degree polynomial. Let \mathcal{G} be a PPT algorithm, which given the security parameter λ , outputs the description of a group \mathbb{G} of order $q = q(1^\lambda)$. Let \mathcal{A} be a probabilistic algorithm that takes as input (a description of) \mathbb{G} and a 4-tuple of elements in \mathbb{G}^d , and outputs 0 or 1. We say that \mathcal{A} solves the f -DDH problem with advantage $\varepsilon_{\mathcal{A}}(\lambda)$, where

$$\varepsilon_{\mathcal{A}}(\lambda) = |\Pr[\mathcal{A}(\mathbb{G}, (\mathbf{g}, \mathbf{g}^{\mathbf{a}}, \mathbf{g}^{\mathbf{b}}, \mathbf{g}^{\mathbf{c}})) = 1] - \Pr[\mathcal{A}(\mathbb{G}, (\mathbf{g}, \mathbf{g}^{\mathbf{a}}, \mathbf{g}^{\mathbf{b}}, \mathbf{g}^{\mathbf{ab}})) = 1]|$$

where $\mathbf{g} \leftarrow \mathbb{G}^d$ and $\mathbf{a} \leftarrow R_f, \mathbf{b} \leftarrow R_f, \mathbf{c} \leftarrow R_f$. In other words, given $(\mathbf{g}, \mathbf{g}^{\mathbf{a}}, \mathbf{g}^{\mathbf{b}}, \mathbf{g}^{\mathbf{c}})$, the problem is to decide whether $\mathbf{c} = \mathbf{ab}$ or \mathbf{c} is a random element in R_f .

Equivalently, we can think of the problem instance as being given in the alternative representation $(h(\mathbf{w}), h(\mathbf{wa}), h(\mathbf{wb}), h(\mathbf{wc}))$. This makes no difference to the adversary, as he would not be given \mathbf{w} – but he knows that such a \mathbf{w} exists. From the above we construct the following assumption.

Definition 2 (The f -DDH Assumption). For any probabilistic polynomial time algorithm \mathcal{A} as in Definition 1, it holds that $\varepsilon_{\mathcal{A}}(\lambda)$ is negligible as a function of λ .

Note that this is a generalization of the DDH problem: for a polynomial f of degree 1, $R_f = \mathbb{F}_q$ and f -DDH is just the standard DDH problem in \mathbb{G} .

Now we look a bit closer at the polynomial f . We can distinguish between two different cases: one where f is reducible and one where f is irreducible. For the first case we have the following theorem:

Theorem 1 (f -DDH for reducible f). Let f be a d -degree reducible polynomial and suppose f_0 divides f , then solving f -DDH is polynomial time reducible to solving f_0 -DDH.

Proof. Let d_0 and d be the degrees of f_0 and f respectively. Let us consider an element \mathbf{w} in R_f . We know that \mathbf{w} can be written as $w_{d-1}x^{d-1} + \dots + w_0 + (f)$. If we map \mathbf{w} to R_{f_0} by reducing modulo f_0 we get an element $\mathbf{v} = v_{d_0-1}x^{d_0-1} + \dots + v_0 + (f_0)$. In fact, reduction modulo f_0 is a ring homomorphism $\phi : R_f \rightarrow R_{f_0}$. In particular, it is linear and therefore has a matrix M . By (1) we can let M act on \mathbf{w} , so we get $h(\mathbf{w})^M = h(\phi(\mathbf{w})) = h(\mathbf{v})$. Hence, M can be used to efficiently map an f -DDH instance $(h(\mathbf{w}), h(\mathbf{w}\mathbf{a}), h(\mathbf{w}\mathbf{b}), h(\mathbf{w}\mathbf{c}))$ to an f_0 -DDH instance $(h(\phi(\mathbf{w})), h(\phi(\mathbf{w}\mathbf{a})), h(\phi(\mathbf{w}\mathbf{b})), h(\phi(\mathbf{w}\mathbf{c}))) = (h(\phi(\mathbf{w})), h(\phi(\mathbf{w})\phi(\mathbf{a})), h(\phi(\mathbf{w})\phi(\mathbf{b})), h(\phi(\mathbf{w})\phi(\mathbf{c})))$. If $\mathbf{c} = \mathbf{a}\mathbf{b}$, then $\phi(\mathbf{c}) = \phi(\mathbf{a})\phi(\mathbf{b})$, while if \mathbf{c} is uniform in R_f , then $\phi(\mathbf{c})$ is uniformly chosen in R_{f_0} . Thus, if we can solve f_0 -DDH, we can solve f -DDH with the same advantage.

4 The d -DDH Problem

Theorem 1 implies that f -DDH is no harder than f_0 -DDH, where f_0 is the smallest irreducible factor in f . The natural conclusion is therefore that we should only look at the irreducible polynomials. In this case we know that our ring R_f is a field, namely the extension field \mathbb{F}_{q^d} where d is the degree of f . In fact, since all fields with q^d elements are isomorphic, f -DDH is equivalent f' -DDH for any f' which is also irreducible and of the same degree as f . This is because the isomorphism can be implemented as a linear mapping in the same fashion as in the proof of Theorem 1. We can thus efficiently map an f -DDH instance to an f' -DDH instance and hence the only thing that may matter to the hardness of the problem is the degree of the extension. In the following, we will talk about d -DDH. In this definition we do not fix f ; we can use any d -degree irreducible polynomial and otherwise the game is the same as in Definition 1.

Theorem 2. Let d' divide d , so $\mathbb{F}_{q^{d'}}$ is a subfield of \mathbb{F}_{q^d} , then d' -DDH is no harder than d -DDH.

We refer to Appendix D for the proof of this theorem.

We now show that d -DDH for $d > 1$ is, in fact, harder than DDH in the generic group model. For this, we need two auxiliary results. The first is a standard result, known as the Schwartz-Zippel lemma [Sch80, Zip79]:

Theorem 3. For a non-zero multivariate polynomial over a finite field K of degree at most t , if uniformly random and independent values are assigned to the variables, the probability that this produces a root is at most $t/|K|$.

The second is our main technical result supporting the hardness of d -DDH:

Theorem 4. Let f_1, f_2, f_3, f_4 be affine functions, where $f_i : (\mathbb{F}_{q^d})^3 \mapsto \mathbb{F}_q$, where $d > 1$. Call the inputs $\mathbf{x}, \mathbf{y}, \mathbf{z}$, and assume that $\mathbf{xy} = \mathbf{z}$ implies $(f_1 f_2 - f_3 f_4)(\mathbf{x}, \mathbf{y}, \mathbf{z}) = 0$. Then $(f_1 f_2 - f_3 f_4)(\mathbf{x}, \mathbf{y}, \mathbf{z}) = 0$ for all inputs.

The proof can be found in Appendix A.

Theorem 5. In the generic group model, the d -DDH assumption holds for $d > 1$, even when the adversary is given an oracle allowing him to solve the DDH problem.

Proof. Recall that an instance to the d -DDH problem can be written as $(h(\mathbf{w}), h(\mathbf{wa}), h(\mathbf{wb}), h(\mathbf{wc}))$ for a fixed generator h of \mathbb{G} . A way to phrase the problem that will be convenient here is to assume that a random bit b has been chosen, if $b = 0$ then $\mathbf{c} = \mathbf{ab}$ and otherwise \mathbf{c} is uniform. It will be sufficient to show that a polynomial-time adversary cannot guess b with non-negligible advantage over $1/2$.

We will show in the generic group model that the problem remains hard even if the adversary is given \mathbf{w} . From \mathbf{w} , it is easy to compute \mathbf{w}^{-1} . As we have seen, it is easy to multiply a known field element “into the exponent” since this is a linear operation, so we can equivalently think of the problem as being given instead as $(h(\mathbf{x}), h(\mathbf{y}), h(\mathbf{z}))$, where the adversary now has to decide whether $\mathbf{z} = \mathbf{xy}$.

The adversary \mathcal{A} will be a polynomial-time generic group adversary, which (as is standard) is formalized by giving him random strings representing group elements and access to computing the group operation and inversion via oracles. In our case, we also give \mathcal{A} access to an oracle for solving DDH. In fact we will provide something a bit stronger: Given $g, g^i, g^j, g^k \in \mathbb{G}$, the oracle will say whether $k = ij \pmod q$. Since we are representing all elements as powers of the fixed generator h , it is easy to see that we can equivalently state this as follows: given $h^{\alpha_1}, h^{\alpha_2}, h^{\alpha_3}, h^{\alpha_4}$, is it the case that $\alpha_1 \alpha_2 = \alpha_3 \alpha_4 \pmod q$? We will provide \mathcal{A} with an oracle for this question in the following.

We now consider an algorithm \mathcal{B} playing the following game with the adversary \mathcal{A} . \mathcal{B} chooses $3d + 1$ bit strings $\sigma_0, \dots, \sigma_{3d+1}$ uniformly in $\{0, 1\}^m$, for a sufficiently large m . These strings represent the encoded elements which algorithm \mathcal{A} will work with. Internally, \mathcal{B} keeps track of the encoded elements using polynomials in the ring $\mathbb{F}_q[X_1, \dots, X_{d-1}, Y_0, \dots, Y_{d-1}, Z_0, \dots, Z_{d-1}]$. Externally, the elements that \mathcal{B} gives to \mathcal{A} are just bit strings in $\{0, 1\}^m$. To maintain consistency, \mathcal{B} creates a list L of pairs (F, σ) where F is a polynomial in the ring specified above and σ is a bit string. List L represents elements in \mathbb{G} . Initially, L is populated by the elements $\{(1, \sigma_0), (X_1, \sigma_1), \dots, (X_{d-1}, \sigma_{d-1}), (Y_0, \sigma_d), \dots, (Y_{d-1}, \sigma_{2d-1}), (Z_0, \sigma_{2d}), \dots, (Z_{d-1}, \sigma_{3d-1})\}$. The algorithm \mathcal{B} starts the game providing \mathcal{A} with $\sigma_0, \dots, \sigma_{3d+1}$. The simulation of the oracles goes as follows:

Group action: Given two elements σ_i, σ_j in \mathbb{G} , \mathcal{B} recovers the corresponding polynomials F_i and F_j and computes $F_i + F_j$. If $F_i + F_j$ is already in L , \mathcal{B} returns to \mathcal{A} the corresponding bit string; otherwise it returns a uniform element σ in $\{0, 1\}^m$ and stores $(F_i + F_j, \sigma)$ in L .

Inversion: Given an element σ in \mathbb{G} , \mathcal{B} recovers its internal representation F and computes $-F$. If the polynomial $-F$ is already in L , \mathcal{B} returns the corresponding bit string; otherwise it returns a uniform string σ and stores $(-F, \sigma)$ in L .

DDH oracle: Given elements $\sigma_1, \sigma_2, \sigma_3, \sigma_4$ in \mathbb{G} , algorithm \mathcal{B} recovers the corresponding polynomials F_1, F_2, F_3, F_4 , tests if $F_1 F_2 = F_3 F_4$ and sends “yes” or “no” to \mathcal{A} accordingly.

In the final stage, \mathcal{A} outputs a bit b' . At this point, \mathcal{B} chooses uniform values $\mathbf{x} = (x_1, \dots, x_{d-1})$, $\mathbf{y} = (y_0, \dots, y_{d-1})$, $\mathbf{z} = (z_0, \dots, z_{d-1})$ in \mathbb{F}_{q^d} and a bit b ; then, it sets $X_1 = x_1, \dots, X_{d-1} = x_{d-1}$,

$Y_0 = y_0, \dots, Y_{d-1} = y_{d-1}$. Finally, if $b = 1$ it sets $Z_0 = z_0, \dots, Z_{d-1} = z_{d-1}$, otherwise it sets $Z_0 = (\mathbf{xy})_0, \dots, Z_{d-1} = (\mathbf{xy})_{d-1}$.

Trivially, in the simulation, $b = b'$ with probability $1/2$. We can, however, also argue that what the adversary sees in a real attack where $b = 0$ is statistically indistinguishable from the simulation, and that this is also the case when $b = 1$. This will clearly imply the theorem.

If $b = 1$, it is clear that the only way in which the simulation could be inconsistent with a real attack is if, after we choose value for $\mathbf{x}, \mathbf{y}, \mathbf{z}$, either two different polynomials in L , F_i, F_j , happen to produce the same value or some query to the DDH oracle F_1, F_2, F_3, F_4 satisfies that $F_1F_2 - F_3F_4$ is not the 0 polynomial, but produces 0 after assigning values. In the $b = 1$ case, all values for coordinates of $\mathbf{x}, \mathbf{y}, \mathbf{z}$ are chosen independently, so Theorem 3 applies to show that for a single oracle query $F_1F_2 - F_3F_4$ or a single difference $F_i - F_j$, the probability of having 0 after assigning values is negligible because q is exponentially large and all polynomials involved have degree at most 2. Further, by the union bound, since we only have a polynomial number of polynomials to consider, the overall probability of having 0 after assigning values is also negligible.

For the $b = 0$ case, some notation: let $X = (X_0, \dots, X_{d-1})$, $Y = (Y_0, \dots, Y_{d-1})$, $Z = (Z_0, \dots, Z_{d-1})$, and $XY = (g_0(X, Y), \dots, g_{d-1}(X, Y))$, where $g_i(X, Y)$ is a degree-2 polynomial describing how the i 'th coordinate of the product of two field elements is computed from the input coordinates.

Now, if $b = 0$, there are two initial possibilities for inconsistency between simulation and real attack. The first is if some query F_1, F_2, F_3, F_4 to the DDH oracle satisfies that $(F_1F_2 - F_3F_4)(X, Y, Z) \neq 0$, but $(F_1F_2 - F_3F_4)(X, Y, XY)$ is the 0-polynomial. This is ruled out by Theorem 4, since the F_i 's have degree at most 1 and can therefore be thought of as affine functions. The second potential inconsistency is if two distinct polynomials F_i, F_j in L satisfy that $(F_i - F_j)(X, Y, XY)$ is the 0 polynomial. To see that this cannot happen, note that since each F_i has degree at most 1, it can be decomposed uniquely as $F_i(X, Y, Z) = F_i^x(X) + F_i^y(Y) + F_i^z(Z) + c_i$ for a constant c_i and polynomials $F_i^x(X), F_i^y(Y), F_i^z(Z)$ of degree at most 1 and constant term 0. A collision as described here can only happen if $(F_i^z - F_j^z)(Z) \neq 0$, but $(F_i^z - F_j^z)(XY) = 0$. This leads to a contradiction: we can assign values $Y_0 = 1, Y_1 = 0, \dots, Y_{d-1} = 0$, corresponding to the 1-element in \mathbb{F}_{q^d} . With this assignment, we get that $(F_i^z - F_j^z)(X) = 0$, contradicting that $(F_i^z - F_j^z)(Z) \neq 0$. Having ruled out the two initial possibilities for inconsistency, the only remaining possibility is that an unfortunate choice of values for the variables lead to collisions, as in the $b = 1$ case. Again by Theorem 3, this happens with negligible probability since the involved polynomials have degree at most 4. \square

We now look at what happens to d -DDH in a bilinear group. In such a group it is well-known that DDH is easy, and we show that this is also the case for d -DDH. The EDDH assumption presented in [HYZX08] is equivalent to d -DDH for $d = 2$. It was claimed that EDDH is hard also in generic bilinear groups, which is however refuted by the following result:

Theorem 6. *d -DDH over any bilinear group can be solved in polynomial time.*

Proof. We assume that the extension field \mathbb{F}_{q^d} has been constructed using some fixed irreducible polynomial f . Consider any two elements $\mathbf{x}, \mathbf{y} \in \mathbb{F}_{q^d}$ as vectors $\mathbf{x} = (x_0, \dots, x_{d-1}), \mathbf{y} = (y_0, \dots, y_{d-1})$ and write the product as $\mathbf{xy} = (z_0, \dots, z_{d-1})$. Now, multiplication of \mathbf{x} and \mathbf{y} takes place by multiplying the polynomials $x_0 + \dots + x_{d-1}X^{d-1}$ and $y_0 + \dots + y_{d-1}X^{d-1}$ and reducing modulo $f(X)$. From this it follows that we can write

$$z_k = \sum \alpha_{ij}^k x_i y_j$$

for coefficients $\alpha_{ij}^k \in \mathbb{F}_q$ that depend only on $f(x)$. Now, if we are given d -tuples $h(\mathbf{x}), h(\mathbf{y})$, it follows from the above that we can efficiently compute a representation in the target group G_T of \mathbf{xy} . Namely, for every k , we have

$$e(h, h)^{z_k} = \prod_{ij} (e(h, h)^{x_i y_j})^{\alpha_{ij}^k} = \prod_{ij} e(h^{x_i}, h^{y_j})^{\alpha_{ij}^k}$$

and h^{x_i}, h^{y_j} can be taken directly from $h(\mathbf{x}), h(\mathbf{y})$. So if we define

$$e(h, h)(\mathbf{xy}) = (e(h, h)^{z_0}, \dots, e(h, h)^{z_{d-1}})$$

what we have shown is that we can compute $e(h, h)(\mathbf{xy})$ efficiently from $h(\mathbf{x}), h(\mathbf{y})$.

Now, consider an input instance of d -DDH, in the form $h(\mathbf{w}), h(\mathbf{wa}), h(\mathbf{wb}), h(\mathbf{wc})$. Observe that we have $\mathbf{c} = \mathbf{ab}$ if and only if $\mathbf{wa wb} = \mathbf{w wc} = \mathbf{w}^2 \mathbf{ab}$. It now follows immediately from the above that we can decide if $\mathbf{ab} = \mathbf{c}$ by computing $e(h, h)(\mathbf{wa wb})$ and $e(h, h)(\mathbf{w wc})$ and comparing the two.

Although of course not all groups are bilinear, this result nevertheless motivates looking for alternative assumptions with similar properties that can be assumed to be hard in bilinear groups. We do this in Section 6.

5 Applications of d -DDH

In this section we present a number of applications for the d -DDH assumption. We will see that its similarity to the DDH problem allows us to use it in virtually any context where DDH can be used. We thus show that we can achieve a variety of constructions with the same amortized efficiency but based on a weaker assumption.

5.1 Pseudorandom Functions

We construct pseudorandom functions (PRF) from d -DDH by taking the construction from [NR97] and showing that the natural modification where we work in the extension field also gives a PRF.

Definition 3. Let $F = \{F_k\}$ be a family of keyed functions where $F_k : A_k \rightarrow B_k$, for every k in the key space \mathcal{K} . We say that F is a family of pseudorandom functions if for all PPT algorithms \mathcal{D} , any polynomial p and large enough λ ,

$$|\Pr[\mathcal{D}^{F_k(\cdot)}(1^\lambda) = 1] - \Pr[\mathcal{D}^{f(\cdot)}(1^\lambda) = 1]| < 1/p(\lambda),$$

where k is chosen uniformly in \mathcal{K} and f is chosen uniformly from the set of functions mapping A_k to B_k .

PRF Construction We construct a function family $F = \{f_k\}$ as follows. The index k specifies a tuple $(q, \mathbb{G}^d, \mathbf{g}, \mathbf{a}_0, \dots, \mathbf{a}_n)$ where q is a prime number, \mathbb{G} is a group of order q , \mathbf{g} is an element of \mathbb{G}^d and $\mathbf{a}_0, \dots, \mathbf{a}_n$ are random in \mathbb{F}_{q^d} . Finally, we define $f_k : \{0, 1\}^n \rightarrow \mathbb{G}^d$, $f_k(x_1, \dots, x_n) = \mathbf{g}^{\mathbf{a}_0 \prod_{i=1}^n \mathbf{a}_i}$.

Theorem 7. Under the d -DDH assumption, the family $F = \{f_k\}$ defined above is a family of pseudorandom functions.

The proof of the theorem follows the exact same line as in [NR97]. Essentially the proof is done by a hybrid argument in which we define a sequence of functions $\{h_i\}$ where h_0 is f_k and h_n is a uniformly random function. An adversary that distinguishes between h_0 and h_n will also distinguish between h_i and h_{i+1} , for some i , which reduces to the d -DDH problem.

5.2 Public Key Encryption

We now apply d -DDH to public key encryption. If we modify in the natural way the Elgamal [Gam84] scheme, we obtain CPA secure encryption based on d -DDH.

- $\text{Gen}(1^\lambda)$: Let $\mathbb{G} \leftarrow \mathcal{G}(1^\lambda)$. Choose a random element $\mathbf{g} \leftarrow \mathbb{G}^d$ and random $\mathbf{x} \leftarrow \mathbb{F}^{q^d}$. Compute $\mathbf{h} = \mathbf{g}^{\mathbf{x}}$. The secret key is then $sk = \mathbf{x}$ and the public key is $pk = (\mathbf{h})$, where \mathbb{G} can be considered public parameters.
- $\text{Enc}(pk, M)$: Let the message be $M \in \mathbb{G}^d$. Choose randomly $\mathbf{r} \leftarrow \mathbb{F}^{q^d}$. Output the ciphertext $CT = (\mathbf{g}^{\mathbf{r}}, \mathbf{h}^{\mathbf{r}} \cdot M)$.
- $\text{Dec}(sk, CT)$: Write the ciphertext as $CT = (\mathbf{e}, \mathbf{c})$. Output $M' = \mathbf{c} \cdot (\mathbf{e}^{\mathbf{x}})^{-1}$

The proof of correctness and security follows immediately as for standard Elgamal.

5.3 Applications in general

Having seen the two examples above, it should not be surprising that all DDH-based cryptographic schemes we are aware of can be based on d -DDH instead. This is basically because all involved algorithms (such as key generation, encryption, and security reduction) will work given only black-box access to a group \mathcal{G} and a finite field K . We just need that for $g \in \mathcal{G}$ and $x \in K$, $g^x \in \mathcal{G}$ is well-defined and standard “axioms” such as $g^{x+y} = g^x g^y$ and $(g^x)^y = g^{xy}$ hold. The exact same scheme and security proof can be run, based on $(\mathcal{G}, K) = (\mathbb{G}, \mathbb{F}_q)$ or based on $(\mathcal{G}, K) = (\mathbb{G}^d, \mathbb{F}_{q^d})$. The only difference is that we need the d -DDH assumption in the latter case. Thus, for instance, CCA secure encryption [CS98] and circular secure or auxiliary input secure encryption [BHHO08] follow immediately from d -DDH.

5.4 Efficiency

For all constructions mentioned here, we can define a notion of amortized complexity. For a PRF, this is the computation time needed to produce a single pseudorandom group element; for an encryption scheme it is the computation time needed to encrypt a group element.

An important point is that in all applications we are aware of, the amortized complexity is essentially the same for constructions based on DDH and on d -DDH. This is because for $\mathbf{g} \in \mathbb{G}^d$ and $\mathbf{a} \in \mathbb{F}_{q^d}$, $\mathbf{g}^{\mathbf{a}}$ corresponds to a tuple of length d where each entry is an expression of the form $\prod g_i^{\alpha_i}$. By a well-known algorithm (see [Pip76]) such a value can be computed in time roughly what you need for a single exponentiation in \mathbb{G} .

As a concrete example, computing the PRF defined above requires essentially a single exponentiation: $\mathbf{g}^{(\mathbf{a}_0 \prod_{x_i=1} \mathbf{a}_i)}$. This produces d pseudorandom elements at amortized cost roughly 1 exponentiation in \mathbb{G} , which is the same cost as the DDH based version.

Various optimizations are known that save computation in the constructions we consider here. However, all the optimizations we are aware of can be applied to both variants based on DDH and d -DDH, and therefore do not affect our conclusion on the amortized complexities.

6 The Vector DDH Problem

The main observation in this section is that we can construct a problem that is generically harder than DDH by revealing only the last entry of the final vector in an f -DDH instance. In the following, we study in detail what happens if we choose f to be x^d . It turns out that there is a simple way of expressing products in $R_d = \mathbb{F}_q[X]/(x^d)$. If we take $\mathbf{x} = (x_0, \dots, x_{d-1})$ and $\mathbf{y} = (y_0, \dots, y_{d-1})$ in R_d , we have:

$$\mathbf{xy} = \left(x_0 y_0, \dots, \sum_{k=0}^{i-1} x_k y_{i-1-k}, \dots, \sum_{k=0}^{d-1} x_k y_{d-1-k} \right). \quad (2)$$

We define the d -VDDH problem just like d -DDH, except that the problem instance is now of the form $(h(\mathbf{w}), h(\mathbf{wa}), h(\mathbf{wb}), h(\mathbf{wc}))[d]$, where we recall that $\mathbf{x}[d]$ is the d th entry of the vector \mathbf{x} , that is x_{d-1} if we start numbering from 0.

Definition 4 (The d -VDDH Problem). *Let d be an integer. Let \mathcal{G} be a PPT algorithm, which given the security parameter λ , outputs the description of a group \mathbb{G} of order $q = q(1^\lambda)$. Let \mathcal{A} be a probabilistic algorithm that takes as input (a description of) \mathbb{G} and a 3-tuple in \mathbb{G}^d plus an element in \mathbb{G} , and outputs 0 or 1.*

We say that \mathcal{A} solves the d -VDDH problem with advantage $\varepsilon_{\mathcal{A}}(\lambda)$, where

$$\varepsilon_{\mathcal{A}}(\lambda) = |\Pr[\mathcal{A}(\mathbb{G}, (\mathbf{g}, \mathbf{g}^{\mathbf{a}}, \mathbf{g}^{\mathbf{b}}, \mathbf{g}^{\mathbf{c}}[d])) = 1] - \Pr[\mathcal{A}(\mathbb{G}, (\mathbf{g}, \mathbf{g}^{\mathbf{a}}, \mathbf{g}^{\mathbf{b}}, \mathbf{g}^{\mathbf{ab}}[d])) = 1]|$$

where $\mathbf{g} \leftarrow \mathbb{G}^d$ and $\mathbf{a} \leftarrow R_d, \mathbf{b} \leftarrow R_d, \mathbf{c} \leftarrow R_d$.

Definition 5 (The d -VDDH Assumption). *For any probabilistic polynomial time algorithm \mathcal{A} as in Definition 4, it holds that $\varepsilon_{\mathcal{A}}(\lambda)$ is negligible as a function of λ .*

Recall the notation introduced in Section 3: $\mathbf{g} = (g_0, \dots, g_{d-1}) = (h^{w_0}, \dots, h^{w_{d-1}})$. Note that we WLOG can choose $w_0 = 1$, so $h(\mathbf{w}) = (g_0, g_0^{w_1}, \dots, g_0^{w_{d-1}})$. To prove that d -VDDH is generically hard, even in d -linear groups, it is useful to do the following parameter substitution: set $\mathbf{x} = \mathbf{wa}$, $\mathbf{y} = \mathbf{wb}$. The d -VDDH problem now becomes deciding whether the last element is the d th coordinate of \mathbf{xyw}^{-1} or is random.

Now, set $\mathbf{w}^{-1} = (z_0, z_1, \dots, z_{d-1})$ and consider the z_i as unknowns. Since $\mathbf{ww}^{-1} = \mathbf{1} = (1, 0, \dots, 0)$ we get $d - 1$ equations involving the z_i 's, using the product introduced in (2):

$$\left\{ \begin{array}{l} z_0 = 1 \\ z_1 = -w_1 \\ \vdots \\ z_i = -w_i - \sum_{j+l=i} z_l w_j \\ \vdots \\ z_{d-1} = -w_{d-1} - \sum_{j+l=d-1} z_l w_j \end{array} \right.$$

In particular, $z_i = -w_1 z_{i-1} - \dots - w_{i-1} z_1 - w_i$. Hence, it can be proved by simple induction that z_i has degree i as a function of the w_j 's. Now, let $p_i(\mathbf{w}, \mathbf{x}, \mathbf{y})$ be the i th entry of $\mathbf{w}^{-1} \mathbf{xy}$. Then $p_d(\mathbf{w}, \mathbf{x}, \mathbf{y})$ has degree $d + 1$ in \mathbf{w}, \mathbf{x} , and \mathbf{y} .

We are now ready to prove the generic hardness of d -VDDH.

Theorem 8. *Even given a d -linear mapping, the d -VDDH holds in the generic group model.*

The proof can be found in Appendix B.

Later, we will need a lemma stating that a generalization of the d -VDDH which considers several generators is equivalent to the original assumption.

Lemma 1. *If d -VDDH is hard for \mathcal{G} , then for any positive integer m*

$$\left\{ (\mathbf{g}_1, \dots, \mathbf{g}_m, \mathbf{g}_1^r[d], \dots, \mathbf{g}_m^r[d]) \mid \mathbf{g}_i \leftarrow \mathbb{G}^d, \mathbf{r} \leftarrow R_d \right\} \stackrel{c}{\approx} \quad (3)$$

$$\left\{ (\mathbf{g}_1, \dots, \mathbf{g}_m, \mathbf{g}_1^{r_1}[d], \dots, \mathbf{g}_m^{r_m}[d]) \mid \mathbf{g}_i \leftarrow \mathbb{G}^d, \mathbf{r}_i \leftarrow R_d \right\}. \quad (4)$$

The proof can be found in Appendix C.

7 Applications of d -VDDH

In this section we discuss a number of natural application of our d -VDDH assumption. Throughout this section we will use the ring $R_d = \mathbb{F}_q[X]/(f)$ for $f = X^d$.

7.1 Public Key Encryption

It is immediate how to construct a CPA-secure encryption schemes from the d -VDDH assumption family. We now show how to extend them to chosen-ciphertext (CCA) secure schemes. Let us first recall the definition of chosen-ciphertext security for encryption schemes.

Definition 6. *A scheme PKE is CCA secure if for any PPT adversary $\mathcal{A} = (\mathcal{A}_1, \mathcal{A}_2)$, any polynomial p and large enough λ ,*

$$\text{Adv}_{\mathcal{A},h} := \left| \Pr[\text{CCA}_0(\text{PKE}, \mathcal{A}, 1^\lambda)] - \Pr[\text{CCA}_1(\text{PKE}, \mathcal{A}, 1^\lambda)] \right| < 1/p(\lambda),$$

where $\text{CCA}_i(\text{PKE}, \mathcal{A}, 1^\lambda)$ is output from the following experiment:

$$\begin{aligned} (pk, sk) &\leftarrow \mathcal{G}(1^\lambda) \\ (m_0, m_1, \text{state}) &\leftarrow \mathcal{A}_1^{\text{Dec}(sk, \cdot)}(1^\lambda, pk) \text{ with } |m_0| = |m_1| \\ CT^* &\leftarrow \text{Enc}_{pk}(m_b), \\ \text{Output } b' &\leftarrow \mathcal{A}_2^{\text{Dec}(sk, \cdot)}(1^\lambda, \text{state}, CT^*) \end{aligned}$$

In the second phase the decryption oracle $\text{Dec}(sk, \cdot)$ returns \perp when queried on the challenge ciphertext CT^* .

We now give the construction of our CCA secure encryption scheme. Let (E, D) be a symmetric encryption scheme with key-space $K \in \mathbb{G}$. Let $\mathbb{T} : \mathbb{G}^d \rightarrow \mathbb{F}_q$ be a target collision resistant hash function (see [HK07] for a definition) and define $\hat{\mathbb{T}}(\mathbf{x}) := (\mathbb{T}(x), 0, \dots, 0) \in R_d$. (Note that for two elements $\mathbf{x} \neq \mathbf{y}$ we have that $\hat{\mathbb{T}}(\mathbf{x}) - \hat{\mathbb{T}}(\mathbf{y})$ is invertible in R_d unless $\mathbb{T}(\mathbf{x}) = \mathbb{T}(\mathbf{y})$).

- **Gen**(1^λ): Let $\mathbb{G} \leftarrow \mathcal{G}(1^\lambda)$. Choose a random generator $\mathbf{g}_1 \leftarrow \mathbb{G}^d$ and random indices $\mathbf{w}, \mathbf{x}, \mathbf{y} \leftarrow R_d$. Compute $\mathbf{g}_2 = \mathbf{g}^{\mathbf{w}}, \mathbf{u} = \mathbf{g}^{\mathbf{x}}, \mathbf{v} = \mathbf{g}^{\mathbf{y}}$. The secret key is then $sk = \mathbf{w}, \mathbf{x}, \mathbf{y}$ and the public key is $pk = (\mathbb{G}, \mathbf{g}_1, \mathbf{g}_2, \mathbf{u}, \mathbf{v})$.

- $\text{Enc}(pk, M)$: Choose randomly $\mathbf{r} \leftarrow R_d$. Compute $\mathbf{c}_1 = \mathbf{g}^{\mathbf{r}}$ and $\mathbf{c}_2 = (\mathbf{u}^{\mathbf{t}}\mathbf{v})^{\mathbf{r}}$, where $\mathbf{t} = \hat{\mathbf{T}}(\mathbf{c}_1) \in R_d$. Compute the symmetric part as $C = \mathbf{E}_K(m)$, where $K = \mathbf{g}_2[d]$. Output the ciphertext $CT = (\mathbf{c}_1, \mathbf{c}_2, C)$.
- $\text{Dec}(sk, CT)$: Write the ciphertext as $CT = (\mathbf{c}_1, \mathbf{c}_2, C)$. If $\mathbf{c}_1^{\mathbf{x}^{\mathbf{t}+\mathbf{y}}} \neq \mathbf{c}_2$ then return \perp . Otherwise return $\mathbf{D}_K(C)$, where $K = \mathbf{c}_1^{\mathbf{w}}[d]$.

It is easy to see that correctness follows by the definition of the public/secret key and by the correctness of the symmetric scheme. To prove the theorem we need that symmetric scheme is secure in the sense of authenticated encryption. That is, it acts as a one-time pad plus any decryption query (with respect to a uniform random key) is rejected. We refer again to [HK07] for a formal definition.

Theorem 9. *If (\mathbf{E}, \mathbf{D}) is a symmetric encryption scheme secure in the sense of authenticated encryption, \mathbf{T} is a target collision resistant hash function and the d -VDDH holds in \mathbb{G} , then the encryption scheme is IND-CCA secure.*

The proof is exactly the same as Theorem 2 in [HK07] where an encryption scheme is proved CCA secure from the DDH assumption. We give some intuition about the proof.

The difficulty in the security reduction is that an adversary against the d -VDDH assumption has to answer the decryption queries and hence has to distinguish between consistent ciphertexts (i.e., ciphertexts for that $\mathbf{c}_1^{\mathbf{x}^{\mathbf{t}+\mathbf{y}}} = \mathbf{c}_2$ holds) and inconsistent ones, without knowing $\mathbf{w} = \log_{\mathbf{g}_1} \mathbf{g}_2$. The simulator inputs $(\mathbf{g}_1, \mathbf{g}_2, \mathbf{c}_1^* = \mathbf{g}_1^{\mathbf{r}}, K^*)$ and wants to distinguish $K^* = \mathbf{g}_2^{\mathbf{r}}[d]$ from a uniform element in \mathbb{G} . In the simulation the values \mathbf{u}, \mathbf{v} from the public-key are set-up such that the tuple $\mathbf{c}_1^*, \mathbf{c}_2^*$ can be used as the challenge ciphertext for some efficiently computable \mathbf{c}_2^* and the value K^* as the symmetric key. More precisely, we define $\mathbf{u} = \mathbf{g}_1^{\mathbf{x}_1} \mathbf{g}_2^{\mathbf{x}_2}$, $\mathbf{v} = \mathbf{g}_1^{\mathbf{y}_1} \mathbf{g}_2^{-\mathbf{t} \cdot \mathbf{x}_2}$ for uniform $\mathbf{x}_1, \mathbf{y}_1 \in R_d$, $\mathbf{x}_2 \in R_d^*$ and $\mathbf{t}^* = \hat{\mathbf{T}}(\mathbf{c}_1^*)$. By construction, the corresponding real session key is $\mathbf{g}_2^{\mathbf{r}}[d]$ so breaking the indistinguishability of the scheme is equivalent to solving the d -VDDH problem. It leaves to deal with the decryption queries for $CT = (\mathbf{c}_1, \mathbf{c}_2, C)$. The simulator is not able to distinguish consistent from inconsistent ciphertexts. However, for ciphertexts with $\mathbf{t} = \hat{\mathbf{T}}(\mathbf{c}_1) \neq \mathbf{t}^*$ (these are the interesting cases) the simulator implements an alternative decryption algorithm by computing the symmetric key as $K = (\mathbf{c}_1 \mathbf{c}_2^{-\mathbf{x}_1 \mathbf{t} + \mathbf{y}_1})^{(\mathbf{x}_2 (\mathbf{t} - \mathbf{t}^*))^{-1}}[d]$. (Note that by the properties of $\hat{\mathbf{T}}$, $\mathbf{x}_2 (\mathbf{t} - \mathbf{t}^*) \in R^*$ so its inverse is well-defined.) This has the following consequences.

It is easy to verify that if the queried ciphertext is consistent then the alternative decryption algorithm yields the correct symmetric key $K = \mathbf{c}_1^{\mathbf{w}}$. If the queried ciphertext is inconsistent then the alternative decapsulation algorithm yields one single symmetric key K that is uniformly distributed over \mathbb{G} . (The probability space is taken over all possible $\mathbf{x}_1, \mathbf{x}_2, \mathbf{y}_1$ that yield \mathbf{u}, \mathbf{v} from the public-key given to the adversary.) Returning this key K to the adversary would completely determine the simulator's secret key and hence also the virtual symmetric key K' for the next decapsulation query. However, this key K is used to decrypt the symmetric part C of the decryption query and by the authenticity property of the latter this will always lead to a rejection. Hence the decryption query is answered correctly and no information about the secret key is leaked which makes it possible to apply the same argument again.

7.2 Generalized BHHO Encryption

In this section we define a public-key encryption scheme which is heavily inspired by the scheme in [BHHO08]. Here, however, the cryptosystem is based on d -VDDH, instead of DDH.

Let λ be the security parameter and $m = m(\lambda)$ be a parameter of the scheme. The encryption scheme is $\text{PKE} = (\text{Gen}, \text{Enc}, \text{Dec})$.

- $\text{Gen}(1^\lambda)$: Let $\mathbb{G} \leftarrow \mathcal{G}(1^\lambda)$. Choose a vector of random generators $\mathbf{g} = (\mathbf{g}_1, \dots, \mathbf{g}_m)$, $\mathbf{g}_i \leftarrow \mathbb{G}^d$ and random bit string $\mathbf{s} = (s_1, \dots, s_m) \leftarrow \{0, 1\}^m$. Compute $\mathbf{y} = \prod_{i=1}^m \mathbf{g}_i^{(s_i, 0, \dots, 0)}$, where $(s_1, 0, \dots, 0)$ is viewed as an element in R_d . The secret key is then $sk = \mathbf{s}$ and the public key is $pk = (\mathbb{G}, \mathbf{g}, \mathbf{y})$, where \mathbb{G} and \mathbf{g} can be considered public parameters.
- $\text{Enc}(pk, M)$: Let the message be $M \in \mathbb{G}$. Choose randomly $\mathbf{r} \leftarrow R_d$. Compute $f_i = \mathbf{g}_i^{\mathbf{r}}[d]$ and output the ciphertext $CT = (f_1, \dots, f_m, \mathbf{y}^{\mathbf{r}}[d] \cdot M)$.
- $\text{Dec}(sk, CT)$: Write the ciphertext as $CT = (f_1, \dots, f_m, c)$. Output $M' = c \cdot (\prod_{i=1}^m f_i^{s_i})^{-1}$

Correctness of decryption follows since

$$\begin{aligned} \prod_{i=1}^m f_i^{s_i} &= \prod_{i=1}^m (\mathbf{g}_i^{\mathbf{r}}[d])^{s_i} = \prod_{i=1}^m (g_{i1}^{r_d} \cdot g_{i2}^{r_{d-1}} \cdots g_{id}^{r_1})^{s_i} = \prod_{i=1}^m (g_{i1}^{r_d s_i} \cdot g_{i2}^{r_{d-1} s_i} \cdots g_{id}^{r_1 s_i}) \\ &= \prod_{i=1}^m (g_{i1}^{s_i}, \dots, g_{id}^{s_i})^{(r_1, \dots, r_d)} [d] = \prod_{i=1}^m (g_{i1}, \dots, g_{id})^{(s_i, 0, \dots, 0)(r_1, \dots, r_d)} [d] = \prod_{i=1}^m \mathbf{g}_i^{(s_i, 0, \dots, 0)\mathbf{r}} [d] = \mathbf{y}^{\mathbf{r}} [d] \end{aligned}$$

CPA security in the usual sense follows immediately from Lemma 1. We will, however, argue that the scheme is also leakage resilient in the auxiliary input model.

Auxiliary Input Security The definition of security w.r.t auxiliary inputs is exactly as in [DGK⁺10].

Definition 7. A scheme PKE is CPA secure w.r.t. auxiliary inputs from a function class \mathcal{H} if for any function $h \in \mathcal{H}$, any PPT adversary $\mathcal{A} = (\mathcal{A}_1, \mathcal{A}_2)$, any polynomial p and large enough λ ,

$$\text{Adv}_{\mathcal{A}, h} := \left| \Pr[\text{CPA}_0(\text{PKE}, \mathcal{A}, 1^\lambda, h)] - \Pr[\text{CPA}_1(\text{PKE}, \mathcal{A}, 1^\lambda, h)] \right| < 1/p(\lambda),$$

where $\text{CPA}_b(\text{PKE}, \mathcal{A}, 1^\lambda, h)$ is output from the following experiment:

$$\begin{aligned} (pk, sk) &\leftarrow \text{Gen}(1^\lambda) \\ (m_0, m_1, \text{state}) &\leftarrow \mathcal{A}_1(1^\lambda, pk, h(sk, pk)) \text{ with } |m_0| = |m_1| \\ CT^* &\leftarrow \text{Enc}_{pk}(m_b), \\ \text{Output } b' &\leftarrow \mathcal{A}_2(1^\lambda, \text{state}, CT^*) \end{aligned}$$

The functions we will consider are those where the secret key is hard to compute even given the leakage. More precisely, $\mathcal{H}_{ow}(f(k))$ consists of all PT functions $h : \{0, 1\}^{|sk|+|pk|} \rightarrow \{0, 1\}^*$ s.t. given $h(sk, pk)$ (for $(sk, pk) \leftarrow \text{Gen}(1^\lambda)$), no PPT algorithm can find sk with probability greater than $f(k)$. A scheme secure w.r.t auxiliary inputs from $\mathcal{H}_{ow}(f(k))$ is called $f(k)$ -AI-CPA secure.

We are now ready to state the theorem about the security of our scheme.

Theorem 10. Let $m = (4 \log q^d)^{1/\epsilon}$, for some $\epsilon > 0$. Assuming that d -VDDH is hard for \mathcal{G} , the scheme above is (2^{-m^ϵ}) -AI-CPA secure.

The complete details of the proof of Theorem 10 are given in the appendix of the full version. It follows the exact same lines as in the proof in [DGK⁺10]. Essentially, the security follows by first using Lemma 1 to argue that the first m elements in the ciphertext are indistinguishable from random. Finally we argue that if we can distinguish the last element from a random element, then we can invert h , which by assumption is hard. This reduction is done by using a Goldreich-Levin Theorem for large fields, which is proved in [DGK⁺10]. The theorem basically says that if we can distinguish the inner product $\langle \mathbf{r}, \mathbf{s} \rangle$ from a uniformly random element given $h(\mathbf{s})$, then we can invert h .

There is a trade-off between the ciphertext size and the hardness of the leakage functions that we can protect against. Obtaining security against functions that are 2^{-m^ϵ} -hard to invert, requires that $m = (4 \log q^d)^{1/\epsilon}$ instead of $m = (4 \log q)^{1/\epsilon}$, which is a polynomial overhead in the ciphertext size.

7.3 Pseudorandom Functions

In this section we present a construction for pseudorandom functions (see Definition 3) based on the d -VDDH assumption. This construction is a modification of the DDH-based one in [NR97].

PRF Construction We construct a function family $F = \{f_k\}$ as follows. The index k specifies a tuple $(q, \mathbb{G}, g_1, g_2, e, \mathbf{a}_0, \dots, \mathbf{a}_n)$ where q is a prime number, \mathbb{G} is a group of order q , g_1, g_2 are two generators of \mathbb{G} , $e : \mathbb{G}^2 \rightarrow \mathbb{G}_T$ is a bilinear map and $\mathbf{a}_0, \dots, \mathbf{a}_n$ are random in R_2 . For any such index k we denote $t_1 = e(g_1, g_1), t_2 = e(g_2, g_1)$ and $\mathbf{t} = (t_1, t_2)$. Finally, we define $f_k : \{0, 1\}^n \rightarrow \mathbb{G}_T$, $f_k(x_1, \dots, x_n) = \mathbf{t}^{\mathbf{a}_0 \prod_{x_i=1} \mathbf{a}_i} [2]$.

Theorem 11. *Under the 2-VDH assumption, the family $F = \{f_k\}$ defined above is a family of pseudorandom functions.*

Proof. Let $k = (q, \mathbf{g}, \mathbf{a}_0, \dots, \mathbf{a}_n)$ be some random index. We have to prove that f_k cannot be distinguished from a uniformly random map from $\{0, 1\}^n$ to \mathbb{G}_T . To prove this, we define for $j \in \{0, \dots, n\}$ the map $h_j : \{0, 1\}^n \rightarrow \mathbb{G}_T$. In order to compute h_j , we choose $\mathbf{a}_{j+1}, \dots, \mathbf{a}_n$ in R_2 and we parse every input $x \in \{0, 1\}^n$ as $x = yx_{j+1} \dots x_n$, where $y \in \{0, 1\}^j$. For every such y we choose \mathbf{t}_y uniformly in \mathbb{G}_T . Finally, we define $h_j(y, x_{j+1}, \dots, x_n) = \mathbf{t}_y^{\prod_{x_i=1} \mathbf{a}_i} [2]$. Notice that $h_0 = f_k$ while h_n is a uniformly random function. If there exists a PPT adversary \mathcal{A} that can distinguish h_0 from h_n with some non-negligible advantage ϵ , then there must be a j for which \mathcal{A} can distinguish with advantage at least ϵ/n between h_j and h_{j+1} . We will show that such an adversary can be used to distinguish tuples as in Lemma 1 where the elements of the tuples are in \mathbb{G} .

We define a distinguisher \mathcal{D} as follows. On input $(\mathbf{g}, \mathbf{g}^{\mathbf{a}}, \mathbf{g}^{\mathbf{b}_1}, \dots, \mathbf{g}^{\mathbf{b}_m}, \mathbf{g}^{\mathbf{c}_1} [2], \dots, \mathbf{g}^{\mathbf{c}_m} [2])$, \mathcal{D} invokes \mathcal{A} with input 1^λ and chooses $\mathbf{a}_{j+2}, \dots, \mathbf{a}_n \leftarrow R_2$. When \mathcal{A} sends a query $x \in \{0, 1\}^n$, distinguisher \mathcal{D} parses x as $yx_{j+1} \dots x_n$. If y was not queried before, \mathcal{D} considers the next fresh $\mathbf{g}^{\mathbf{b}_l}$ otherwise it takes the element that was used before. Finally, \mathcal{D} computes $h(x)$ as $\mathbf{t}^{\mathbf{b}_l \prod_{x_i=1} \mathbf{a}_i} [2]$ if $x_{j+1} = 0$. Note that $\mathbf{t}^{\mathbf{b}_l}$ can be computed using e as follows: $(e(\mathbf{g}^{\mathbf{b}_l} [1], g_1), e(\mathbf{g}^{\mathbf{b}_l} [2], g_1)) = (e(g_1^{\mathbf{b}_{l,1}}, g_1), e(g_1^{\mathbf{b}_{l,2}} g_2^{\mathbf{b}_{l,1}}, g_1)) = (t_1^{\mathbf{b}_{l,1}}, t_2^{\mathbf{b}_{l,1}} t_1^{\mathbf{b}_{l,2}}) = \mathbf{t}^{\mathbf{b}_l}$. If $x_{j+1} = 1$, distinguisher \mathcal{D} first computes $\mathbf{t}^{\mathbf{a}_l} [1]$ as $e(\mathbf{g}^{\mathbf{a}_l} [1], \mathbf{g}^{\mathbf{b}_l} [1]) = (g_1^{\mathbf{a}_l}, g_1^{\mathbf{b}_l}) = t_1^{\mathbf{a}_l \mathbf{b}_l}$, then it sets $h(x) = (\mathbf{t}^{\mathbf{a}_l \mathbf{b}_l} [1], e(\mathbf{g}^{\mathbf{c}_l} [2], g_1))^{\prod_{x_i=1} \mathbf{a}_i} [2]$. We want to argue that if $\mathbf{c}_l = \mathbf{a}_l \mathbf{b}_l$ then $h(x) = h_j(x)$, while if \mathbf{c}_l is random, we have $h(x) = h_{j+1}(x)$. Notice that this is clearly true whenever $x_{j+1} = 0$, so let's assume that $x_{j+1} = 1$. If $\mathbf{c}_l = \mathbf{a}_l \mathbf{b}_l$ then

$(t_1^{a_1 b_{l,1}}, e(g_1^{a_1 b_{l,2} + a_2 b_{l,1}} g_2^{a_1 b_{l,1}}, g_1)) = (t_1^{a_1 b_{l,1}}, t_2^{a_1 b_{l,1}} t_1^{a_1 b_{l,2} + a_2 b_{l,1}}) = \mathbf{t}^{\mathbf{a}b_l}$, thus $h(x) = \mathbf{t}^{\mathbf{a}b_l} \prod_{x_i=1}^{\mathbf{a}_i} [2] = h_j(x)$, where $\mathbf{a}_{j+1} = \mathbf{a}$. When instead \mathbf{c}_i is a random element, we notice that even with $\mathbf{t}^{\mathbf{a}b_l} [1]$ in the first component, the output of h will still be of the form $\mathbf{g}^{\mathbf{r}} \prod_{x_i=1}^{\mathbf{a}_i} [2]$ for a random \mathbf{r} .

References

- [BB04] Dan Boneh and Xavier Boyen. Efficient selective-id secure identity-based encryption without random oracles. In Christian Cachin and Jan Camenisch, editors, *EUROCRYPT*, volume 3027 of *Lecture Notes in Computer Science*, pages 223–238. Springer, 2004.
- [BBS04] Dan Boneh, Xavier Boyen, and Hovav Shacham. Short group signatures. In Matthew K. Franklin, editor, *CRYPTO*, volume 3152 of *Lecture Notes in Computer Science*, pages 41–55. Springer, 2004.
- [BHHO08] Dan Boneh, Shai Halevi, Michael Hamburg, and Rafail Ostrovsky. Circular-secure encryption from decision diffie-hellman. In David Wagner, editor, *CRYPTO*, volume 5157 of *Lecture Notes in Computer Science*, pages 108–125. Springer, 2008.
- [BMR10] Dan Boneh, Hart William Montgomery, and Ananth Raghunathan. Algebraic pseudorandom functions with improved efficiency from the augmented cascade. In Ehab Al-Shaer, Angelos D. Keromytis, and Vitaly Shmatikov, editors, *ACM Conference on Computer and Communications Security*, pages 131–140. ACM, 2010.
- [CS98] Ronald Cramer and Victor Shoup. A practical public key cryptosystem provably secure against adaptive chosen ciphertext attack. In Hugo Krawczyk, editor, *CRYPTO*, volume 1462 of *Lecture Notes in Computer Science*, pages 13–25. Springer, 1998.
- [DGK⁺10] Yevgeniy Dodis, Shafi Goldwasser, Yael Tauman Kalai, Chris Peikert, and Vinod Vaikuntanathan. Public-key encryption schemes with auxiliary inputs. In Daniele Micciancio, editor, *TCC*, volume 5978 of *Lecture Notes in Computer Science*, pages 361–381. Springer, 2010.
- [DH76] Whitfield Diffie and Martin E. Hellman. New Directions in Cryptography. *IEEE Transactions on Information Theory*, IT-22(6):644–654, 1976.
- [DY05] Yevgeniy Dodis and Aleksandr Yampolskiy. A verifiable random function with short proofs and keys. In Serge Vaudenay, editor, *Public Key Cryptography*, volume 3386 of *Lecture Notes in Computer Science*, pages 416–431. Springer, 2005.
- [Gam84] Taher El Gamal. A public key cryptosystem and a signature scheme based on discrete logarithms. In *CRYPTO*, pages 10–18, 1984.
- [HK07] Dennis Hofheinz and Eike Kiltz. Secure hybrid encryption from weakened key encapsulation. In *CRYPTO*, pages 553–571, 2007.
- [HYZX08] Huawei Huang, Bo Yang, Shenglin Zhu, and Guozhen Xiao. Generalized elgamal public key cryptosystem based on a new diffie-hellman problem. In Joonsang Baek, Feng Bao, Kefei Chen, and Xuejia Lai, editors, *ProvSec*, volume 5324 of *Lecture Notes in Computer Science*, pages 1–21. Springer, 2008.
- [Kil07] Eike Kiltz. Chosen-ciphertext secure key-encapsulation based on gap hashed diffie-hellman. In Tatsuaki Okamoto and Xiaoyun Wang, editors, *Public Key Cryptography*, volume 4450 of *Lecture Notes in Computer Science*, pages 282–297. Springer, 2007.
- [NR97] Moni Naor and Omer Reingold. Number-theoretic constructions of efficient pseudo-random functions. In *FOCS*, pages 458–467, 1997.
- [Pip76] Nicholas Pippenger. On the evaluation of powers and related problems (preliminary version). In *FOCS*, pages 258–263, 1976.
- [Sch80] J. T. Schwartz. Fast probabilistic algorithms for verification of polynomial identities. *J. ACM*, 27:701–717, October 1980.
- [Sha07] Hovav Shacham. A Cramer-Shoup encryption scheme from the Linear Assumption and from progressively weaker Linear variants. Cryptology ePrint Archive, Report 2007/074, February 2007. <http://eprint.iacr.org/>.
- [Sta96] Markus Stadler. Publicly verifiable secret sharing. In *EUROCRYPT*, pages 190–199, 1996.
- [Zip79] Richard Zippel. Probabilistic algorithms for sparse polynomials. In *EUROSAM*, pages 216–226, 1979.

A Proof of Theorem 4

We restate the Theorem here with slightly different notation that will be convenient later:

Theorem 12. *Let $f_1, f_2, f_3, f_4 : \mathbb{F}_{q^d}^3 \rightarrow \mathbb{F}_q$ be affine functions with $q > 2$ and $d > 2$. Suppose that the function $(f_1 f_2 - f_3 f_4)(X, Y, XY)$ vanishes on $\mathbb{F}_{q^d}^2$. Then the function $(f_1 f_2 - f_3 f_4)(X, Y, Z)$ vanishes on $\mathbb{F}_{q^d}^3$.*

Proof. An affine function $f : \mathbb{F}_{q^d}^3 \rightarrow \mathbb{F}_q$ is of the form

$$f(X, Y, Z) = e + \phi(X, Y, Z),$$

where $e \in \mathbb{F}_q$ is a constant and where $\phi : \mathbb{F}_{q^d}^3 \rightarrow \mathbb{F}_q$ is an \mathbb{F}_q -linear function. Clearly, ϕ is of the form

$$\phi(X, Y, Z) = \phi'(X) + \phi''(Y) + \phi'''(Z),$$

where $\phi', \phi'', \phi''' : \mathbb{F}_{q^d} \rightarrow \mathbb{F}_q$ are \mathbb{F}_q -linear functions.

It is an elementary fact from the theory of finite fields that any \mathbb{F}_q -linear function $g(X)$ mapping \mathbb{F}_{q^d} to \mathbb{F}_q is of the form $g(X) = \text{Tr}(\mathbf{a}X)$ for some $\mathbf{a} \in \mathbb{F}_{q^d}$. Here, $\text{Tr} : \mathbb{F}_{q^d} \rightarrow \mathbb{F}_q$ denotes the trace-function, i.e., $\text{Tr}(\mathbf{x}) = \mathbf{x} + \mathbf{x}^q + \dots + \mathbf{x}^{q^{d-1}}$ for all $\mathbf{x} \in \mathbb{F}_{q^d}$.

Hence, there exist $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{F}_{q^d}$ such that

$$\phi'(X) = \text{Tr}(\mathbf{a}X), \quad \phi''(Y) = \text{Tr}(\mathbf{b}Y), \quad \phi'''(Z) = \text{Tr}(\mathbf{c}Z).$$

Putting it all together, it follows that an affine function $f : \mathbb{F}_{q^d}^3 \rightarrow \mathbb{F}_q$ is of the form

$$f(X, Y, Z) = e + \text{Tr}(\mathbf{a}X) + \text{Tr}(\mathbf{b}Y) + \text{Tr}(\mathbf{c}Z).$$

Let $e_i \in \mathbb{F}_q, \mathbf{a}_i, \mathbf{b}_i, \mathbf{c}_i \in \mathbb{F}_{q^d}$ ($i = 1, 2, 3, 4$) be such that

$$f_i(X, Y, Z) = e_i + \text{Tr}(\mathbf{a}_i X) + \text{Tr}(\mathbf{b}_i Y) + \text{Tr}(\mathbf{c}_i Z) \quad (i = 1, 2, 3, 4).$$

Substituting the (polynomial) expression for the trace-function, it follows there is a constant $e \in \mathbb{F}_q$ and there are polynomials

$$h'(X) \in \mathbb{F}_q[X], \quad h''(Y) \in \mathbb{F}_q[Y], \quad h'''(Z) \in \mathbb{F}_q[Z],$$

$$h_1(X, Y) \in \mathbb{F}_q[X, Y], \quad h_2(X, Z) \in \mathbb{F}_q[X, Z], \quad h_3(Y, Z) \in \mathbb{F}_q[Y, Z]$$

such that

$$(f_1 f_2 - f_3 f_4)(X, Y, Z) = F(X, Y, Z),$$

as functions on $\mathbb{F}_{q^d}^3$, where

$$F(X, Y, Z) = e + h'(X) + h''(Y) + h'''(Z) + h_1(X, Y) + h_2(X, Z) + h_3(Y, Z) \in \mathbb{F}_q[X, Y, Z].$$

Note that the total degree of $F(X, Y, Z)$ is at most $2q^{d-1}$. Substituting $XY = Z$, the condition of the theorem implies that

$$F(X, Y, XY) \equiv 0,$$

the zero-polynomial. This implies, in particular, that the polynomial $F(X, Y, Z)$ has many zeroes. Unfortunately, however, this number is not large enough compared to its total degree for Theorem 3 (Schwartz-Zippel) to imply that F must be the zero-polynomial.

Exploiting the fact that the bi-degrees of the monomials are of a special form (resulting from the expression for the trace-function), we first argue that several terms must vanish individually. This will simplify matters significantly.

First, observe that (disregarding their coefficients),

- We can write $h'(X) = g'_1(X) + g'_2(X)$, where the monomials of $g'_1(X)$ are all of the form X^{q^i} , and the monomials of $g'_2(X)$ are all of the form $X^{q^i+q^j}$.
- We can write $h''(Y) = g''_1(Y) + g''_2(Y)$, where the monomials of $g''_1(Y)$ are all of the form Y^{q^i} , and the monomials of $g''_2(Y)$ are all of the form $Y^{q^i+q^j}$.
- We can write $h'''(Z) = g'''_1(Z) + g'''_2(Z)$, where the monomials of $g'''_1(Z)$ are all of the form Z^{q^i} , and the monomials of $g'''_2(Z)$ are all of the form $Z^{q^i+q^j}$.
- The monomials of $h_1(X, Y)$ are all of the form $X^{q^i}Y^{q^j}$.
- The monomials of $h_2(X, Z)$ are all of the form $X^{q^i}Z^{q^j}$.
- The monomials of $h_3(Y, Z)$ are all of the form $Y^{q^i}Z^{q^j}$.

Second, after substituting $XY = Z$, observe that (disregarding their coefficients),

- The monomials of $g'_1(X)$ are of the form X^{q^i} , while monomials of $g'_2(X)$ are of the form $X^{q^i+q^j}$.
- The monomials of $g''_1(Y)$ are of the form Y^{q^i} , while monomials of $g''_2(Y)$ are of the form $Y^{q^i+q^j}$.
- The monomials of $g'''_1(XY)$ are of the form $X^{q^i}Y^{q^i}$, while monomials of $g'''_2(XY)$ are of the form $X^{q^i+q^j}Y^{q^i+q^j}$.
- The monomials of $h_1(X, Y)$ are all of the form $X^{q^i}Y^{q^j}$.
- The monomials of $h_2(X, XY)$ are all of the form $X^{q^i+q^j}Y^{q^j}$.
- The monomials of $h_3(Y, XY)$ are all of the form $X^{q^i}Y^{q^i+q^j}$.

Since $F(X, Y, XY) \equiv 0$, the first (trivial) consequence is that $e = 0$, since the constant terms of the polynomials $h', h'', h''', h_1, h_2, h_3$ are all equal to 0. Next,

$$g'_1(X) \equiv 0, g'_2(X) \equiv 0, g''_1(Y) \equiv 0, g''_2(Y) \equiv 0,$$

since, for each of these polynomials, it clearly holds that its monomials cannot be “canceled” by any monomials from any of the other polynomials. The same holds for $h_2(X, XY)$ and $h_3(X, XY)$, this can be seen using the uniqueness of q -ary representation of the integers. So we have

$$h_2(X, XY) \equiv 0, h_3(Y, XY) \equiv 0.$$

We then consider what this means for $h_2(X, Z)$. The total degree of the polynomial is at most $2q^{d-1}$. Yet, if we set X to be any non-zero value, then if we let Y run through all values in \mathbb{F}_{q^d} XY runs through all values as well. By the above observation that $h_2(X, XY) = 0$, this gives us $(q^d - 1)q^d$ choices of values for (X, Z) where $h_2(X, Z) = 0$. So the number of zeros for $h_2(X, Z)$ is at least (counting also $(X, Z) = (0, 0)$):

$$1 + (q^d - 1)q^d = q^{2d} - q^d + 1 > \frac{2q^{d-1}}{q^d} \cdot q^{2d},$$

where for the sharpness of the inequality it is used that $q > 2$. This means, by Theorem 3 (Schwartz-Zippel), that $h_2(X, Z)$ has too many zeroes to be anything else than the zero-polynomial. The same holds for $h_3(Y, Z)$ by the same argument. Thus,

$$h_2(X, Z) \equiv 0, \quad h_3(Y, Z) \equiv 0.$$

Finally we conclude, again by uniqueness of q -ary representation, that $g_2'''(XY) \equiv 0$, whence $g_2'''(Z) \equiv 0$. Substituting all this back, it follows that

$$F(X, Y, Z) = g_1'''(Z) + h_1(X, Y)$$

as polynomials.

Since $F(X, Y, XY) \equiv 0$,

$$g_1'''(XY) + h_1(X, Y) \equiv 0, \tag{5}$$

and since $h_1(X, Y) = \text{Tr}(\mathbf{a}_1 X)\text{Tr}(\mathbf{b}_2 Y) + \text{Tr}(\mathbf{a}_2 X)\text{Tr}(\mathbf{b}_1 Y) - \text{Tr}(\mathbf{a}_3 X)\text{Tr}(\mathbf{b}_4 Y) - \text{Tr}(\mathbf{a}_4 X)\text{Tr}(\mathbf{b}_3 Y)$, we get

$$-g_1'''(XY) = \text{Tr}(\mathbf{a}_1 X)\text{Tr}(\mathbf{b}_2 Y) + \text{Tr}(\mathbf{a}_2 X)\text{Tr}(\mathbf{b}_1 Y) - \text{Tr}(\mathbf{a}_3 X)\text{Tr}(\mathbf{b}_4 Y) - \text{Tr}(\mathbf{a}_4 X)\text{Tr}(\mathbf{b}_3 Y) \tag{6}$$

as polynomials.

Proof for $d > 4$. Recall that if $\mathbf{a}_i \neq 0$ then $\text{Tr}(\mathbf{a}_i X)$ is a (non-trivial) \mathbb{F}_q -linear function from \mathbb{F}_{q^d} to \mathbb{F}_q , whence its kernel has dimension $d-1$. Therefore, if $d > 4$, the intersection of the 4 kernels of the $\text{Tr}(\mathbf{a}_i X)$'s is non-trivial: there exists $x_0 \neq 0$ which is in all 4 kernels. This implies that $h'''(x_0 \mathbf{y}) = 0$ for all $\mathbf{y} \in \mathbb{F}_{q^d}$. Hence, $g_1'''(Z) \equiv 0$ as a polynomial as well.

But then

$$F(X, Y, Z) = g_1'''(Z) + h_1(X, Y) = h_1(X, Y)$$

which means that in fact $F(X, Y, Z) = F(X, Y, XY)$ as polynomials, and so $F(X, Y, Z)$ must vanish on $\mathbb{F}_{q^d}^3$, as claimed.

Proof for $d = 3, 4$. To get the result in this case, we exploit some more of the information we get from the fact that $F(X, Y, XY)$ vanishes: Since $g_2'(X) \equiv 0$ we get that

$$0 \equiv g_2'(X) = \text{Tr}(\mathbf{a}_1 X)\text{Tr}(\mathbf{a}_2 X) - \text{Tr}(\mathbf{a}_3 X)\text{Tr}(\mathbf{a}_4 X)$$

as polynomials. Note that if either $\text{Tr}(\mathbf{a}_1 X) \equiv 0$ or $\text{Tr}(\mathbf{a}_2 X) \equiv 0$, then $\text{Tr}(\mathbf{a}_3 X)\text{Tr}(\mathbf{a}_4 X) \equiv 0$. Hence, since the polynomial ring $\mathbb{F}_q[X]$ is a domain, either $\text{Tr}(\mathbf{a}_3 X) \equiv 0$ or $\text{Tr}(\mathbf{a}_4 X) \equiv 0$

Two of the 4 polynomials being zero is sufficient to complete the argument later on, so assume for the moment that all 4 polynomials are non-zero, i.e. $\mathbf{a}_i \neq 0$ ($i = 1, 2, 3, 4$). Then the polynomial $\text{Tr}(\mathbf{a}_1 X)$ must divide the polynomial $\text{Tr}(\mathbf{a}_3 X)\text{Tr}(\mathbf{a}_4 X)$. But in fact, $\text{Tr}(\mathbf{a}_1 X)$ must divide either $\text{Tr}(\mathbf{a}_3 X)$ or $\text{Tr}(\mathbf{a}_4 X)$.

To see this, note that, since $\text{Tr}(\mathbf{a}_i X)$ is a (non-trivial) \mathbb{F}_q -linear function from \mathbb{F}_{q^d} to \mathbb{F}_q , its zeros form an \mathbb{F}_q -linear subspace of dimension $d-1$ in \mathbb{F}_{q^d} . Two such *distinct* subspaces can intersect in a subspace of dimension at most $d-2$. Hence, if the zeros of $\text{Tr}(\mathbf{a}_1 X)$ did not overlap completely with the zeros of $\text{Tr}(\mathbf{a}_3 X)$ nor completely with the zeros of $\text{Tr}(\mathbf{a}_4 X)$, it could have at most $2q^{d-2} < q^{d-1}$ zeros, which is a contradiction. So without loss of generality, $\text{Tr}(\mathbf{a}_1 X) = \alpha \text{Tr}(\mathbf{a}_3 X)$ for some non-zero constant $\alpha \in \mathbb{F}_{q^d}$, whence also $\text{Tr}(\mathbf{a}_2 X) = \beta \text{Tr}(\mathbf{a}_4 X)$ for some $\beta \in \mathbb{F}_{q^d}$. But since the functions $\text{Tr}(\mathbf{a}_i X)$ are surjective (onto \mathbb{F}_q), it must hold that $\alpha, \beta \in \mathbb{F}_q$.

Now, consider again equation (6) proved above. If $\mathbf{a}_i = 0$ for some $i \in \{1, 2, 3, 4\}$, then, as noted above, $\mathbf{a}_j = 0$ for some $j \in \{1, 2, 3, 4\}$ with $j \neq i$. Hence, two of the 4 terms in (6) vanish. If $\mathbf{a}_i \neq 0$ for all $i \in \{1, 2, 3, 4\}$, the substitutions $\text{Tr}(\mathbf{a}_1 X) = \alpha \text{Tr}(\mathbf{a}_3 X)$ and $\text{Tr}(\mathbf{a}_2 X) = \beta \text{Tr}(\mathbf{a}_4 X)$ are made. In both cases, it follows that

$$g_1'''(XY) = u_1(X)v_1(Y) + u_2(X)v_2(Y)$$

for \mathbb{F}_q -linear functions u_1, u_2, v_1, v_2 . Now, the argument is completed as we did before for $d > 4$, since the kernels of u_1, u_2 must intersect non-trivially if $d > 2$.

Proof for $d = 2$. To show the result in this case we do a more elaborate case analysis. To make notation more compact in the following, we let $X_i = \text{Tr}(\mathbf{a}_i X)$, $Y_i = \text{Tr}(\mathbf{b}_i Y)$ and $Z_i = \text{Tr}(\mathbf{c}_i Z)$ for $i = 1, \dots, 4$, so X_i, Y_i, Z_i are polynomials and equalities in the following involving these are understood to as equalities of polynomials.

Note that to show the result we want it is enough to show that $g_1'''(XY) = u(X)v(Y)$ for linear functions u, v , since then it follows that $g_1'''(Z) \equiv 0$ in the same way as for $d > 2$. It is also enough to show that all Z_i are 0, since then we would have $g_1'''(Z) = d_1 Z_2 + d_2 Z_1 - d_3 Z_4 - d_4 Z_3 = 0$

To identify the cases we have to consider, note that the argument exploiting $g_2'(X) \equiv 0$ we used to show the theorem for $d = 3, 4$ can also be based on $g_2''(Y) \equiv 0$. In fact, that argument implies that one of the three following cases must occur:

1. One of Y_1, Y_2 is 0 and one of Y_3, Y_4 is 0,
2. All Y_i are non-zero and $Y_1 = \alpha Y_3, Y_2 = \beta Y_4$, with $\alpha = \beta^{-1}$
3. All Y_i are non-zero and $Y_1 = \alpha Y_4, Y_2 = \beta Y_3$, with $\alpha = \beta^{-1}$,

and the same 3 cases apply to the X_i 's:

1. One of X_1, X_2 is 0 and one of X_3, X_4 is 0,
2. All X_i are non-zero and $X_1 = \gamma X_3, X_2 = \delta X_4$, with $\gamma = \delta^{-1}$
3. All X_i are non-zero and $X_1 = \gamma X_4, X_2 = \delta X_3$, with $\gamma = \delta^{-1}$,

So we now consider all combined possibilities, where fortunately some can be handled together due to symmetries.

Case 2 or 3 applies to both the X_i 's and the Y_i 's. Without loss of generality, assume case 2 applies to the Y_i 's. Plugging into (6), we get

$$-g_1'''(XY) = (\beta X_1 - X_3)Y_4 + (\alpha X_2 - X_4)Y_3 \tag{7}$$

If $Y_3 = \lambda Y_4$ for some constant λ we are clearly done, so we may assume this is not the case, i.e., $\ker(Y_3) \neq \ker(Y_4)$. From our earlier arguments, we also have that $h_3(Y, Z) = Z_1 Y_2 + Z_2 Y_1 - Z_3 Y_4 - Z_4 Y_3 = 0$. Since case 2 applies to the Y_i 's, we get

$$(\beta Z_1 - Z_3)Y_4 + (\alpha Z_2 - Z_4)Y_3 = 0$$

This, and that Y_3, Y_4 are non-zero with $\ker(Y_3) \neq \ker(Y_4)$ implies that

$$\beta Z_1 = Z_3, \quad \alpha Z_2 = Z_4, \tag{8}$$

because we can choose a \mathbf{y} such that $\mathbf{y} \in \ker(Y_3)$ but $\mathbf{y} \notin \ker(Y_4)$, and vice versa.

Now, if case 2 also applies to the X_i 's, the argument we just applied to $h_3(Y, Z)$ can also be applied to $h_2(X, Z)$, and will give us that $\delta Z_1 = Z_3$, $\gamma Z_2 = Z_4$. If all $Z_i = 0$, we are already done, so assume without loss of generality that $Z_1 \neq 0$. Then we have $\delta = \beta$ and therefore $X_2 = \beta X_4$. Plugging this into (7) and using $\alpha\beta = 1$ we get $-g_1'''(XY) = (\beta X_1 - X_3)Y_4$ and we are done.

We then assume instead that case 3 applies to the X_i 's. Plugging this into (7), we get

$$-g_1'''(XY) = (\beta\gamma X_4 - X_3)Y_4 + (\alpha\delta X_3 - X_4)Y_3$$

From $\alpha\beta = 1$, $\gamma\delta = 1$ follows that $(\beta\gamma)^{-1} = \alpha\delta$, and hence $(\beta\gamma X_4 - X_3)$ and $(\alpha\delta X_3 - X_4)$ seen as linear mappings have the same kernel. We can therefore choose $\mathbf{x} \neq 0$ in the kernel of both mappings and conclude that $-g_1'''(\mathbf{x}Y)$ is identically 0, hence $g_1'''(Z)$ is 0 and we are done.

Case 1 applies to the X_i 's or the Y_i 's. Assume without loss of generality that case 1 applies to the Y_i 's and that in fact $Y_1 = 0$, $Y_3 = 0$. Plugging this into (6), we get

$$-g_1'''(XY) = X_1Y_2 - X_3Y_4 \tag{9}$$

As a first observation, note that if one of Y_2, Y_4 is 0, or if $Y_2 = \kappa Y_4$ for a constant κ , we are clearly done. So we may assume this is not the case, in other words Y_2, Y_4 are non-zero and have distinct kernels. Exactly the same can be assumed about X_1, X_3 .

Consider now that $h_3(Y, Z) = Z_1Y_2 + Z_2Y_1 - Z_3Y_4 - Z_4Y_3 = 0$ in this case means that $Z_1Y_2 - Z_3Y_4 = 0$. By our assumption on Y_2, Y_4 this implies $Z_1 = Z_3 = 0$. We also know that $h_2(X, Z) = X_1Z_2 + X_2Z_1 - X_3Z_4 - X_4Z_3 = 0$. Plugging in $Z_1 = Z_3 = 0$, we get

$$X_1Z_2 - X_3Z_4 = 0$$

By our assumption on X_1, X_3 , we get $Z_2 = Z_4 = 0$ and we are done.

B Proof of Theorem 8

Proof. Let \mathbb{G} be a cyclic group of order q . Let g be a generator, and let $e : \mathbb{G}^d \rightarrow \mathbb{G}_T$ be a d -linear map. We consider an algorithm \mathcal{B} playing the following game with \mathcal{A} . Algorithm \mathcal{B} chooses $3d+2$ bit strings $\sigma_0, \dots, \sigma_{3d+1}$ uniformly in $\{0, 1\}^m$, for a sufficiently large m . These strings represent the encoded elements which algorithm \mathcal{A} will work with. Internally, \mathcal{B} keeps track of the encoded elements using polynomials in the ring $\mathbb{F}_q[W_1, \dots, W_{d-1}, X_0, \dots, Y_{d-1}, X_0, \dots, Y_{d-1}, T_0, T_1]$. Externally, the elements that \mathcal{B} gives to \mathcal{A} are just bit strings in $\{0, 1\}^m$. To maintain consistency, \mathcal{B} creates two lists L_1 and L_2 of pairs (F, σ) where F is a polynomial in the ring specified above and σ is a bit string. List L_1 represents elements in \mathbb{G} while L_2 represents elements in \mathbb{G}_T . Initially, L_2 is empty and L_1 is set to $\{(1, \sigma_0), (W_1, \sigma_1), \dots, (W_{d-1}, \sigma_{d-1}), (X_0, \sigma_d), \dots, (X_{d-1}, \sigma_{2d-1}), (Y_0, \sigma_{2d}), \dots, (Y_{d-1}, \sigma_{3d-1}), (T_0, \sigma_{3d}), (T_1, \sigma_{3d+1})\}$. Algorithm \mathcal{B} starts the game providing \mathcal{A} with $\sigma_0, \dots, \sigma_{3d+1}$. The simulation of the oracles goes as follows:

Group action: Given two elements σ_i, σ_j in \mathbb{G} , \mathcal{B} recovers the corresponding polynomials F_i and F_j and computes $F_i + F_j$. If $F_i + F_j$ is already in L_1 , \mathcal{B} returns to \mathcal{A} the corresponding bit string; otherwise it returns a uniform element σ in $\{0, 1\}^m$ and stores $(F_i + F_j, \sigma)$ in L_1 . Multiplication in \mathbb{G}_T is handled similarly.

Inversion: Given an element σ in \mathbb{G} , \mathcal{B} recovers its internal representation F and computes $-F$. If the polynomial $-F$ is already in L_1 , \mathcal{B} returns the corresponding bit string; otherwise it returns a uniform string σ and stores $(-F, \sigma)$ in L_1 . Inversion in \mathbb{G}_T is handled analogously.

d -linear Map: Given d elements $\sigma_{i_1}, \dots, \sigma_{i_d}$ in \mathbb{G} , adversary \mathcal{B} recovers the corresponding polynomials F_{i_1}, \dots, F_{i_d} and computes $F = \prod_j F_{i_j}$. If F is already in L_2 , \mathcal{B} returns to \mathcal{A} the corresponding bit string; otherwise it returns a uniform element σ in $\{0, 1\}^m$ and stores (F, σ) in L_2 .

Note that all polynomials in L_1 have degree at most 1, while all polynomials in L_2 have degree at most d . Finally \mathcal{A} outputs a bit b' . At this point, \mathcal{B} chooses uniform values $w_1, \dots, w_{d-1}, x_0, \dots, x_{d-1}, y_0, \dots, y_{d-1}, s$ in \mathbb{F}_q and a bit b ; then, it sets $W_1 = w_1, \dots, W_{d-1} = w_{d-1}, X_0 = x_0, \dots, X_{d-1} = x_{d-1}, Y_0 = y_0, \dots, Y_{d-1} = y_{d-1}, T_b = p_d(\mathbf{w}, \mathbf{x}, \mathbf{y}), T_{1-b} = s$.

The simulation provided by \mathcal{B} is consistent and reveals nothing about b unless two distinct polynomials F_1 and F_2 in L_1 or L_2 take on the same value after the substitution.

First, we argue that \mathcal{A} is not able to intentionally cause a collision. The values that we substitute in the variables W, X, Y and T are all independent except that of T_b . This means that the only collision that \mathcal{A} can engineer consists in producing $p_d(\mathbf{w}, \mathbf{x}, \mathbf{y})$ using a combination of the other polynomials in the list. In fact, for any collision $F_i = F_j$, T_b has to appear as one of the variables, since T_b is the only variable that depends on the value of the other assignments. Then, the equation $F_i = F_j$ reduces to a relation of the kind

$$T_b^d = T_b^{d-1} p_1 + \dots + p_d \quad (10)$$

for polynomials p_i of degree at most i . Equation (10), can be reduced to $T_b = \tilde{p}(W, X, Y)$. However, as we noticed before, the polynomials in the lists have degree at most d , while $p_d(\mathbf{w}, \mathbf{x}, \mathbf{y})$ has degree $d + 1$. Therefore, \mathcal{A} is not able to cause a collision.

Now, it only remains to bound the probability that an unlucky choice of values causes a collision. We know that two polynomials F_i, F_j in L_1 have degree at most 1, so, by theorem ..., the probability that they have the same value is at most $1/q$. Similarly, the probability that two polynomials in L_2 take on the same value after the substitution is at most d/q , since every polynomial in L_2 has degree at most d . Assuming that \mathcal{A} makes Q_g queries to the group operation oracle and Q_b queries to the bilinear oracle, the probability ε of a collision can be bounded as follows:

$$\varepsilon_{\mathcal{A}} \leq \binom{Q_g + 3d + 2}{2} \frac{1}{q} + \binom{Q_b}{2} \frac{d}{q} \leq \left(\binom{Q_g + 3d + 2}{2} + \binom{Q_b}{2} \right) \frac{d}{q} \leq \frac{d((Q_g + 3d + 2)^2 + Q_b^2)}{2q}.$$

Since the probability of \mathcal{A} outputting the correct b' is at most $1/2 + \varepsilon_{\mathcal{A}}$, the above shows that the advantage of \mathcal{A} is negligible in the security parameter, concluding the proof.

C Proof of Lemma 1

Proof. We prove the lemma above by the following hybrid argument.

Assume we have a PPT distinguisher \mathcal{D} that can distinguish with significant advantage $\varepsilon = \varepsilon(\lambda)$ between the two tuples (3) and (4). Then there exists an i s.t. \mathcal{D} can distinguish with advantage at least ε/m between

$$\{(\mathbf{g}_1, \dots, \mathbf{g}_m, \mathbf{g}_1^{\mathbf{r}}[d], \dots, \mathbf{g}_i^{\mathbf{r}}[d], \mathbf{g}_{i+1}^{\mathbf{r}^{i+1}}[d], \dots, \mathbf{g}_m^{\mathbf{r}^m}[d])\} \text{ and} \quad (11)$$

$$\{(\mathbf{g}_1, \dots, \mathbf{g}_m, \mathbf{g}_1^{\mathbf{r}}[d], \dots, \mathbf{g}_{i-1}^{\mathbf{r}}[d], \mathbf{g}_i^{\mathbf{r}^i}[d], \dots, \mathbf{g}_m^{\mathbf{r}^m}[d])\}. \quad (12)$$

This can be used to solve a given d -VDDH instance $(\mathbf{g}, \mathbf{g}^{\mathbf{a}}, \mathbf{g}^{\mathbf{b}}, \mathbf{h}[d])$, where $\mathbf{h}[d]$ is either $\mathbf{g}^{ab}[d]$ or some random value in \mathbb{G} . We construct and give to \mathcal{D} the following tuple

$$\left\{ \left(\mathbf{g}'_1, \dots, \mathbf{g}'_m, \mathbf{g}'_1{}^{\mathbf{r}_1}[d], \dots, \mathbf{g}'_m{}^{\mathbf{r}_m}[d] \right) \right\}.$$

We set $\mathbf{g}'_1 = \mathbf{g}$, $\mathbf{g}'_i = \mathbf{g}^{\mathbf{a}}$. For $j = 2, \dots, i-1$, we choose random \mathbf{x}_j and set $\mathbf{g}'_j = \mathbf{g}^{\mathbf{x}_j}$. Next we set $\mathbf{g}'_1{}^{\mathbf{r}_1}[d] = \mathbf{g}^{\mathbf{b}}[d]$, $\mathbf{g}'_i{}^{\mathbf{r}_i}[d] = \mathbf{h}[d]$. For $j = 2, \dots, i-1$, we let $\mathbf{g}'_j{}^{\mathbf{r}_j}[d] = (\mathbf{g}^{\mathbf{b}})^{\mathbf{x}_j}[d] = (\mathbf{g}^{\mathbf{x}_j})^{\mathbf{b}}[d]$. The rest of the entries in the tuple will just be random values in the appropriate groups, that is, for $j > i$ we let $\mathbf{g}'_j = \mathbf{g}^{\mathbf{w}_j}$ and $\mathbf{g}'_j{}^{\mathbf{r}_j}[d] = g^{z_j}$, for random $\mathbf{w}_j \leftarrow \mathbb{F}_q^d, z_j \leftarrow \mathbb{F}_q$, and generator $g \leftarrow \mathbb{G}$.

It is now clear that if $\mathbf{h}[d]$ is $\mathbf{g}^{ab}[d]$, then the constructed tuple is as in (11), whereas if $\mathbf{h}[d]$ is a random value in \mathbb{G} the tuple is like (12). Therefore, giving this to D enables us to distinguish the d -VDDH instance with same significant advantage ε/m , contradicting that d -VDDH is hard.

D Proof of Theorem 2

Proof. We prove here that 1-DDH is not harder than k -DDH for any k , and then we argue that this can be generalized to the statement above.

Assume we have some PPT algorithm \mathcal{A} that solves the k -DDH problem with some non-negligible advantage ε . Then we can use this to solve a 1-DDH instance. So let a 1-DDH instance be given: (g, g^a, g^b, g^c) , where c is either ab or random in \mathbb{F}_q . We construct with this a k -DDH instance: $((g, 1, \dots, 1), (g^a, 1, \dots, 1), (g^b, 1, \dots, 1), (g^c, 1, \dots, 1))$, where each of the four tuples are elements in \mathbb{G}^k . It can easily be seen, that for a fixed generator h of \mathbb{G} with $g = h^{w_0}$, the above is a valid instance $(h(\mathbf{w}), h(\mathbf{w}\mathbf{a}), h(\mathbf{w}\mathbf{b}), h(\mathbf{w}\mathbf{c}))$, for k -dimensional vectors $\mathbf{w} = (w_0, 0, \dots, 0)$, $\mathbf{a} = (a, 0, \dots, 0)$, $\mathbf{b} = (b, 0, \dots, 0)$, $\mathbf{c} = (c, 0, \dots, 0)$.

The only problem left is that the values $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are not sampled as specified in the definition. Therefore, we need to randomize the instance so we get the right distribution. We can do this by using the technique of random self-reducibility of DDH introduced by [Sta96, NR97]. First we choose three elements $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3 \leftarrow \mathbb{F}_{q^k}$. Then we compute

$$\begin{aligned} \mathbf{g}^{\mathbf{a}'} &= (\mathbf{g}^{\mathbf{a}})^{\mathbf{r}_1} \mathbf{g}^{\mathbf{r}_2}, \\ \mathbf{g}^{\mathbf{b}'} &= \mathbf{g}^{\mathbf{b}} \mathbf{g}^{\mathbf{r}_3}, \\ \mathbf{g}^{\mathbf{c}'} &= (\mathbf{g}^{\mathbf{c}})^{\mathbf{r}_1} (\mathbf{g}^{\mathbf{a}})^{\mathbf{r}_1 \mathbf{r}_3} (\mathbf{g}^{\mathbf{b}})^{\mathbf{r}_1} \mathbf{g}^{\mathbf{r}_2 \mathbf{r}_3}. \end{aligned}$$

Writing $\mathbf{c} = \mathbf{a}\mathbf{b} + \mathbf{e}$ we create from the above the instance $(h(\mathbf{w}), h(\mathbf{w}\mathbf{a}'), h(\mathbf{w}\mathbf{b}'), h(\mathbf{w}\mathbf{c}'))$, where $\mathbf{a}' = \mathbf{r}_1 \mathbf{a} + \mathbf{r}_2$, $\mathbf{b}' = \mathbf{b} + \mathbf{r}_3$, and $\mathbf{c}' = \mathbf{a}'\mathbf{b}' + \mathbf{e}\mathbf{r}_1$. If $\mathbf{c} = \mathbf{a}\mathbf{b}$, such that $\mathbf{e} = \mathbf{0}$, then \mathbf{a}', \mathbf{b}' are random elements in \mathbb{F}_{q^k} and $\mathbf{c}' = \mathbf{a}'\mathbf{b}'$. Otherwise, if $\mathbf{e} \neq \mathbf{0}$, then $\mathbf{a}', \mathbf{b}', \mathbf{c}'$ are all uniformly distributed in \mathbb{F}_{q^k} . Now, we can give the instance to \mathcal{A} , which will decide with probability ε whether $\mathbf{c}' = \mathbf{a}'\mathbf{b}' (\Leftrightarrow \mathbf{c} = \mathbf{a}\mathbf{b} \Leftrightarrow c = ab)$. Thus, we can solve the original instance with the same advantage as \mathcal{A} .

Finally, note that the argument does not depend on the basis field being \mathbb{F}_q . We might as well have started from some other $\mathbb{F}_{q^{d'}}$ and extended to \mathbb{F}_{q^d} , for some $d = kd'$. Therefore, this completes the proof.