
CENTRAL CHARACTERS FOR SMOOTH IRREDUCIBLE MODULAR REPRESENTATIONS OF $GL_2(\mathbf{Q}_p)$

by

Laurent Berger

Abstract. — We prove that every smooth irreducible $\overline{\mathbf{F}}_p$ -linear representation of $GL_2(\mathbf{Q}_p)$ admits a central character.

Introduction

Let Π be a representation of $GL_2(\mathbf{Q}_p)$. We say that Π is smooth, if the stabilizer of any $v \in \Pi$ is an open subgroup of $GL_2(\mathbf{Q}_p)$. We say that Π admits a central character, if every $z \in Z(GL_2(\mathbf{Q}_p))$ acts on Π by a scalar. The smooth irreducible representations of $GL_2(\mathbf{Q}_p)$ over an algebraically closed field of characteristic p , admitting a central character, have been studied by Barthel-Livné in [BL94, BL95] and by Breuil in [Bre03]. The purpose of this note is to prove the following theorem.

Theorem A. — *If Π is a smooth irreducible $\overline{\mathbf{F}}_p$ -linear representation of $GL_2(\mathbf{Q}_p)$, then Π admits a central character.*

The idea of the proof of theorem A is as follows. If Π does not admit a central character, and if $f = \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix}$, then for any nonzero polynomial $Q(X) \in \overline{\mathbf{F}}_p[X]$, the map $Q(f) : \Pi \rightarrow \Pi$ is bijective, so that Π has the structure of a $\overline{\mathbf{F}}_p(X)$ -vector space. The representation Π is therefore a smooth irreducible $\overline{\mathbf{F}}_p(X)$ -linear representation of $GL_2(\mathbf{Q}_p)$, which now admits a central character, since f acts by multiplication by X . It remains to apply Barthel-Livné and Breuil’s classification, which gives the structure of the components of Π after extending scalars to a finite extension K of $\overline{\mathbf{F}}_p(X)$. A corollary of this classification is that these components are all “defined” over a subring R of K , where R is a finitely generated $\overline{\mathbf{F}}_p$ -algebra. This can be used to show that Π is not of finite length, a contradiction.

2000 Mathematics Subject Classification. — 22E50.

Key words and phrases. — Smooth representation; admissible representation; parabolic induction; supersingular representation; central character; Schur’s lemma.

Acknowledgments. I am grateful to C. Breuil, G. Chenevier, P. Colmez, G. Henniart, M. Schein and M.-F. Vignéras for helpful comments.

1. Barthel-Livné and Breuil's classification

Let E be a field of characteristic p . In this section, we recall the explicit classification of smooth irreducible E -linear representations of $\mathrm{GL}_2(\mathbf{Q}_p)$, admitting a central character.

We denote the center of $\mathrm{GL}_2(\mathbf{Q}_p)$ by Z . If $r \geq 0$, then $\mathrm{Sym}^r E^2$ is a representation of $\mathrm{GL}_2(\mathbf{F}_p)$ which gives rise, by inflation, to a representation of $\mathrm{GL}_2(\mathbf{Z}_p)$. We extend it to $\mathrm{GL}_2(\mathbf{Z}_p)Z$ by letting $\begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix}$ act trivially. Consider the representation

$$\mathrm{ind}_{\mathrm{GL}_2(\mathbf{Z}_p)Z}^{\mathrm{GL}_2(\mathbf{Q}_p)} \mathrm{Sym}^r E^2.$$

The Hecke algebra

$$\mathrm{End}_{E[\mathrm{GL}_2(\mathbf{Q}_p)]} \left(\mathrm{ind}_{\mathrm{GL}_2(\mathbf{Z}_p)Z}^{\mathrm{GL}_2(\mathbf{Q}_p)} \mathrm{Sym}^r E^2 \right)$$

is isomorphic to $E[T]$ where T is a Hecke operator, which corresponds to the double class $\mathrm{GL}_2(\mathbf{Z}_p)Z \cdot \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \cdot \mathrm{GL}_2(\mathbf{Z}_p)$. If $\chi : \mathbf{Q}_p^\times \rightarrow E^\times$ is a smooth character, and if $\lambda \in E$, then let

$$\pi(r, \lambda, \chi) = \frac{\mathrm{ind}_{\mathrm{GL}_2(\mathbf{Z}_p)Z}^{\mathrm{GL}_2(\mathbf{Q}_p)} \mathrm{Sym}^r E^2}{T - \lambda} \otimes (\chi \circ \det).$$

This is a smooth representation of $\mathrm{GL}_2(\mathbf{Q}_p)$, with central character $\omega^r \chi^2$ (where $\omega : \mathbf{Z}_p^\times \rightarrow \mathbf{F}_p^\times$ is the “reduction mod p ” map). Let $\mu_\lambda : \mathbf{Q}_p^\times \rightarrow E^\times$ be given by $\mu_\lambda|_{\mathbf{Z}_p^\times} = 1$, and $\mu_\lambda(p) = \lambda$. If $\lambda = \pm 1$, then we have two exact sequences:

$$\begin{aligned} 0 &\rightarrow \mathrm{Sp} \otimes (\chi \mu_\lambda \circ \det) \rightarrow \pi(0, \lambda, \chi) \rightarrow \chi \mu_\lambda \circ \det \rightarrow 0, \\ 0 &\rightarrow \chi \mu_\lambda \circ \det \rightarrow \pi(p-1, \lambda, \chi) \rightarrow \mathrm{Sp} \otimes (\chi \mu_\lambda \circ \det) \rightarrow 0, \end{aligned}$$

where the representation Sp is the “special” representation.

Theorem 1.1. — *If E is algebraically closed, then the smooth irreducible E -linear representations of $\mathrm{GL}_2(\mathbf{Q}_p)$, admitting a central character, are as follows:*

1. $\chi \circ \det$;
2. $\mathrm{Sp} \otimes (\chi \circ \det)$;
3. $\pi(r, \lambda, \chi)$, where $r \in \{0, \dots, p-1\}$ and $(r, \lambda) \notin \{(0, \pm 1), (p-1, \pm 1)\}$.

This theorem is proved in [BL95] and [BL94], which treat the case $\lambda \neq 0$, and in [Bre03], which treats the case $\lambda = 0$.

We now explain what happens if E is not algebraically closed.

Proposition 1.2. — *If Π is a smooth irreducible E -linear representation of $\mathrm{GL}_2(\mathbf{Q}_p)$, admitting a central character, then there exists a finite extension K/E such that $(\Pi \otimes_E K)^\beta$ is a direct sum of K -linear representations of the type described in theorem 1.1.*

Proof. — Barthel and Livné’s methods show (as is observed in §5.3 of [Paš10]) that Π is a quotient of

$$\Sigma = \frac{\mathrm{ind}_{\mathrm{GL}_2(\mathbf{Z}_p)\mathbf{Z}}^{\mathrm{GL}_2(\mathbf{Q}_p)} \mathrm{Sym}^r E^2}{P(T)} \otimes (\chi \circ \det),$$

for some integer $r \in \{0, \dots, p-1\}$, character $\chi : \mathbf{Q}_p^\times \rightarrow E^\times$, and polynomial $P(Y) \in E[Y]$. Let K be a splitting field of $P(Y)$, write $P(Y) = (Y - \lambda_1) \cdots (Y - \lambda_d)$, and let $P_i(Y) = (Y - \lambda_1) \cdots (Y - \lambda_i)$ for $i = 0, \dots, d$. The representations $P_{i-1}(T)\Sigma/P_i(T)\Sigma$ are then isomorphic to $\pi(r, \lambda_i, \chi)$, for $i = 1, \dots, d$. \square

We finish this section by recalling that if $\lambda \neq 0$, then the representations $\pi(r, \lambda, \chi)$ are parabolic inductions (when $\lambda = 0$, they are called supersingular). Let χ_1 and $\chi_2 : \mathbf{Q}_p^\times \rightarrow E^\times$ be two smooth characters, and consider the parabolic induction $\mathrm{ind}_{\mathrm{B}_2(\mathbf{Q}_p)}^{\mathrm{GL}_2(\mathbf{Q}_p)} (\chi_1 \otimes \chi_2)$. The following result is proved in [BL94] and [BL95].

Theorem 1.3. — *If $\lambda \in K \setminus \{0, \pm 1\}$, and if $r \in \{0, \dots, p-1\}$, then $\pi(r, \lambda, \chi)$ is isomorphic to $\mathrm{ind}_{\mathrm{B}_2(\mathbf{Q}_p)}^{\mathrm{GL}_2(\mathbf{Q}_p)} (\chi \mu_{1/\lambda}, \chi \omega^r \mu_\lambda)$.*

2. Proof of the theorem

We now give the proof of theorem A. Let Π be a smooth irreducible $\overline{\mathbf{F}}_p$ -linear representation of $\mathrm{GL}_2(\mathbf{Q}_p)$. We have $\Pi^{(1+p\mathbf{Z}_p)\mathrm{Id}} \neq 0$ (since a p -group acting on a \mathbf{F}_p -vector space always has nontrivial fixed points), so that if Π is irreducible, then $(1 + p\mathbf{Z}_p) \cdot \mathrm{Id}$ acts trivially on Π . If $g \in \mathbf{Z}_p^\times \cdot \mathrm{Id}$, then $g^{p-1} = \mathrm{Id}$ on Π , so that $\Pi = \bigoplus_{\omega \in \mathbf{F}_p} \Pi^{g=\omega \cdot \mathrm{Id}}$. Since Π is irreducible, this implies that the elements of $\mathbf{Z}_p^\times \cdot \mathrm{Id}$ act by scalars.

If $f = \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix}$, then for any nonzero polynomial $Q(X) \in \overline{\mathbf{F}}_p[X]$, the kernel and image of the map $Q(f) : \Pi \rightarrow \Pi$ are subrepresentations of Π . If $Q(f) = 0$ on a nontrivial subspace of Π , then f admits an eigenvector for an eigenvalue $\lambda \in \overline{\mathbf{F}}_p$. This implies that $\Pi = \Pi^{f=\lambda \cdot \mathrm{Id}}$, so that Π does admit a central character. If this is not the case, then $Q(f)$ is bijective for every nonzero polynomial $Q(X) \in \overline{\mathbf{F}}_p[X]$, so that Π has the structure of a $\overline{\mathbf{F}}_p(X)$ -vector space, and is a $\overline{\mathbf{F}}_p(X)$ -linear smooth irreducible representation of $\mathrm{GL}_2(\mathbf{Q}_p)$, admitting a central character.

Let $E = \overline{\mathbf{F}}_p(X)$. Proposition 1.2 gives us a finite extension K of E , such that $(\Pi \otimes_E K)^\beta$ is a direct sum of K -linear representations of the type described in theorem 1.1. The $\overline{\mathbf{F}}_p$ -linear representation underlying $(\Pi \otimes_E K)^\beta$ is isomorphic to $\Pi^{[K:E]}$, and hence of length

$[K : E]$. We now prove that none of the K -linear representations of the type described in theorem 1.1 are of finite length, when viewed as $\overline{\mathbf{F}}_p$ -linear representations.

Let Σ be one such representation, and let $\lambda \in K$ be the corresponding Hecke eigenvalue.

Proposition 2.1. — *There exists a subring R of K , which is a finitely generated $\overline{\mathbf{F}}_p$ -algebra, such that $\Sigma = \Sigma_R \otimes_R K$, where Σ_R is an R -linear representation of $\mathrm{GL}_2(\mathbf{Q}_p)$.*

Proof. — If $\lambda \in \overline{\mathbf{F}}_p$, then theorem 1.1 shows that

$$\Sigma = \frac{\mathrm{ind}_{\mathrm{GL}_2(\mathbf{Z}_p)\mathbf{Z}}^{\mathrm{GL}_2(\mathbf{Q}_p)} \mathrm{Sym}^r \overline{\mathbf{F}}_p^2}{T - \lambda} \otimes_{\overline{\mathbf{F}}_p} K(\chi \circ \det), \text{ or } \mathrm{Sp} \otimes_{\overline{\mathbf{F}}_p} K(\chi \circ \det), \text{ or } K(\chi \circ \det).$$

We can then take $R = \overline{\mathbf{F}}_p[\chi(p)^{\pm 1}]$, and $\Sigma_R = (\mathrm{ind}_{\mathrm{GL}_2(\mathbf{Z}_p)\mathbf{Z}}^{\mathrm{GL}_2(\mathbf{Q}_p)} \mathrm{Sym}^r \overline{\mathbf{F}}_p^2 / (T - \lambda)) \otimes_{\overline{\mathbf{F}}_p} R(\chi \circ \det)$, or $\mathrm{Sp} \otimes_{\overline{\mathbf{F}}_p} R(\chi \circ \det)$, or $R(\chi \circ \det)$, respectively.

If $\lambda \notin \overline{\mathbf{F}}_p$, then by theorem 1.3, we have

$$\Sigma = \mathrm{ind}_{\mathrm{B}_2(\mathbf{Q}_p)}^{\mathrm{GL}_2(\mathbf{Q}_p)} (\chi \mu_{1/\lambda}, \chi \omega^r \mu_\lambda).$$

We can take $R = \overline{\mathbf{F}}_p[\lambda^{\pm 1}, \chi(p)^{\pm 1}]$, and let Σ_R be the set of functions $f \in \Sigma$ with values in R . □

Let $\beta \in \overline{\mathbf{F}}_p$ be such that $(X - \beta) \notin R^\times$, so that $(X - \beta)^j \Sigma_R \neq (X - \beta)^{j+1} \Sigma_R$ for all $j \in \mathbf{Z}$. The representation Σ contains $\cup_{j \in \mathbf{Z}} (X - \beta)^j \Sigma_R$, so that the underlying $\overline{\mathbf{F}}_p$ -linear representation is not of finite length, which is a contradiction. This finishes the proof of theorem A.

References

- [BL94] L. BARTHEL & R. LIVNÉ — “Irreducible modular representations of GL_2 of a local field”, *Duke Math. J.* **75** (1994), no. 2, p. 261–292.
- [BL95] ———, “Modular representations of GL_2 of a local field: the ordinary, unramified case”, *J. Number Theory* **55** (1995), no. 1, p. 1–27.
- [Bre03] C. BREUIL — “Sur quelques représentations modulaires et p -adiques de $\mathrm{GL}_2(\mathbf{Q}_p)$: I”, *Compositio Math.* **138** (2003), no. 2, p. 165–188.
- [Paš10] V. PAŠKŪNAS — “The image of Colmez’s Montréal functor”, preprint, 2010.

July 2011

LAURENT BERGER, UMPA, ENS de Lyon, UMR 5669 du CNRS, Université de Lyon

E-mail : laurent.berger@ens-lyon.fr • *Url* : perso.ens-lyon.fr/laurent.berger/