

**ON THE DETERMINANT FORMULA IN THE INVERSE
SCATTERING PROCEDURE WITH A PARTIALLY KNOWN
STEPLIKE POTENTIAL**

ODILE BASTILLE, ALEXEI RYBKIN

ABSTRACT. We are concerned with the inverse scattering problem for the full line Schrödinger operator $-\partial_x^2 + q(x)$ with a steplike potential q a priori known on $\mathbb{R}_+ = (0, \infty)$. Assuming $q|_{\mathbb{R}_+}$ is known and short range, we show that the unknown part $q|_{\mathbb{R}_-}$ of q can be recovered by

$$q|_{\mathbb{R}_-}(x) = -2\partial_x^2 \log \det (1 + (1 + \mathbb{M}_x^+)^{-1} \mathbb{G}_x),$$

where \mathbb{M}_x^+ is the classical Marchenko operator associated to $q|_{\mathbb{R}_+}$ and \mathbb{G}_x is a trace class integral Hankel operator. The kernel of \mathbb{G}_x is explicitly constructed in term of the difference of two suitably defined reflection coefficients. Since $q|_{\mathbb{R}_-}$ is not assumed to have any pattern of behavior at $-\infty$, defining and analyzing scattering quantities becomes a serious issue. Our analysis is based upon some subtle properties of the Titchmarsh-Weyl m -function associated with \mathbb{R}_- .

CONTENTS

| | |
|--|----|
| 1. Introduction | 1 |
| 2. Notation | 4 |
| 3. Weyl and scattering solutions of the Schrödinger equation | 5 |
| 4. The Titchmarsh-Weyl m -function | 7 |
| 5. The reflection and transmission coefficients | 10 |
| 6. A trace class operator | 16 |
| 7. Classical Marchenko inverse scattering | 17 |
| 8. Main result | 18 |
| References | 20 |

1. INTRODUCTION

As the title suggests, we are concerned with the recovery of an unknown potential of the full line Schrödinger operator $-\partial_x^2 + q(x)$ from a certain set of scattering data. In its classical formulation, the scattering data consist of the reflection coefficient R , bound states $\{-\varkappa_n^2\}_{n=1}^N$ and their norming constants $\{c_n\}_{n=1}^N$.

Date: July, 2011.

1991 Mathematics Subject Classification. 34L25, 34B20, 47B35.

Key words and phrases. Marchenko inverse scattering, Schrödinger operator, Titchmarsh-Weyl m -function.

Based on research supported in part by the NSF under grant DMS 1009673.

It is well-known (see e.g. [9]) that the inverse problem

$$S := \{R(k), -\varkappa_n^2, c_n\}_{k \in \mathbb{R}, 1 \leq n \leq N} \longrightarrow q(x) \quad (1.1)$$

for all x is uniquely solvable through the famous Faddeev-Marchenko inverse scattering procedure. In fact, there is an explicit formula, referred to sometimes as Bargmann or Dyson, (see e.g. [10])

$$q(x) = -2\partial_x^2 \log \det(1 + \mathbb{M}_x), \quad (1.2)$$

where \mathbb{M}_x is the so-called Marchenko operator, a Hankel integral operator whose kernel is constructed in terms of S .

In practice, however, norming constants $\{c_n\}$ are not available. Moreover the authors are unaware of their physical meaning either. Notice that the inverse problem

$$\{R(k)\}_{k \in \mathbb{R}} \longrightarrow q(x) \quad (1.3)$$

is solved uniquely only if the operator $-\partial_x^2 + q(x)$ has no (negative) bound states. The classical example of the so-called one-soliton potential

$$q(x) = -2\varkappa^2 \operatorname{sech}^2 \left(\varkappa x + \log \frac{\sqrt{2\varkappa}}{c} \right)$$

produces the scattering data

$$\{0, -\varkappa^2, c\}$$

with $R(k) = 0$ for all k , suggesting that the inverse problem (1.3) is ill-posed in general.

The inverse problem (1.1) was originally posed and solved for q 's decaying sufficiently fast at infinity. The complete treatment of this problem is done in [9] under the Faddeev condition¹

$$\int_{\mathbb{R}} (1 + |x|) |q(x)| dx < \infty. \quad (1.4)$$

While the Faddeev condition (1.4) is typically satisfied in nuclear physics, many interesting inverse problems (e.g. in geophysics) deal with potentials which do not decay at one of the infinities but still decay at the other infinity. Such potentials are commonly called steplike. A suitable analog of inverse problem (1.1) is well-posed as well (see e.g. [1] and the literature cited therein). New circumstances arise, however, due to a richer negative spectrum of $-\partial_x^2 + q(x)$. But the classical Marchenko theory can be successfully adjusted to such setting too.

In [7] the inverse problem was solved for $q(x) \rightarrow C$, a nonzero constant, as $x \rightarrow -\infty$ and $q(x) \rightarrow 0$ as $x \rightarrow \infty$ sufficiently fast (with some gaps fixed in [8]). The case of q 's periodic on the, say, left hand side and decaying on the other have been treated first by Hruslov [13] (see also [1]).

The more general case of q 's with no certain pattern of behavior at $-\infty$ and identically zero on the right hand side has been recently treated by one of the authors in [16]. To the best of our knowledge, in the context of steplike potentials, the determinant formula (1.2) is not available in the literature.

¹Certain aspects of the theory developed in [9] actually require the stronger condition

$$\int_{\mathbb{R}} (1 + |x|^2) |q(x)| dx < \infty.$$

The situation with steplike potentials is similar to the case (1.4) in that the reflection coefficient alone does not determine the potential uniquely. It is natural to ask what can make up for the unavailable data in (1.1) related to the negative spectrum? This problem has generated considerable interest (see e.g. [3, 4, 6, 12, 15]). In the context of Faddeev potentials, the inverse problem

$$\{R(k), q(x)\}_{k \in \mathbb{R}, x \geq 0} \longrightarrow q(x), \quad x < 0 \tag{1.5}$$

is shown to be well-posed. The problem (1.5) is referred to as the inverse scattering problem with partial information on the potential. In [11], the inverse problem (1.5) is solved for essentially arbitrary potentials as long as the reflection coefficient can be suitably defined. The actual procedure of solving (1.5) in [11] is not scattering but rather spectral through solving the Gelfand-Levitan integral equation.

In [16], one of the authors found a way to adapt the classical Marchenko inverse scattering algorithm to solve (1.5). No analog of the determinant formula (1.2) appears to be found for inverse problems with partial information on the potential (1.1) in the context of steplike potentials. The present paper intends to deal with this issue. More specifically, we show that if q is locally square integrable,

$$\sup_{x \leq 0} \int_{x-1}^x |q|^2 < \infty, \tag{1.6}$$

and the known part $q_+ = q|_{(0, \infty)}$ is subject to (1.4) then

$$q(x) = -2\partial_x^2 \log \det \left(1 + (1 + \mathbb{M}_x^+)^{-1} \mathbb{G}_x \right), \quad x < 0. \tag{1.7}$$

Here \mathbb{M}_x^+ is the Marchenko operator constructed in terms of the scattering data for q_+ and \mathbb{G}_x is the integral Hankel operator with the kernel expressed in terms of the difference $R - R_+$ of the (right) reflection coefficients R for the whole potential q and R_+ corresponding to q_+ .

We emphasize that the main issue here is the existence of the determinant in (1.7) in the classical Fredholm sense. We prove that the operator $(1 + \mathbb{M}_x^+)^{-1} \mathbb{G}_x$ in (1.7) is trace class for every x . Our arguments are based upon the simple but important observation that the difference $R - R_+$ is an analytic function (even though R and R_+ are not) and certain limiting procedures (which, as frequently happens in similar situations, are quite involved).

From the geophysical point of view our situation is related to reflection seismology where one is concerned with recovering certain properties q of deeper layers of the Earth given already known properties q_+ of shallower layers and measured reflections R . Of course if the media do not tend to be homogeneous as the depth increases, a larger range of frequencies is required to investigate deeper layers.

Another example of the inverse problem with partially known steplike potential is related to neutron reflectometry (see e.g. [2]) where properties of an unknown material are studied by scattering neutrons and measuring their reflection. The role of q_+ is played by a layer of known material, called a coating.

The paper is organized as follows. Section 2 lists our notation. Section 3 introduces the scattering quantities in our setting. Section 4 details relevant properties of the Titchmarsh-Weyl m -function which are then applied in our context in Section 5. Section 6 and 7 respectively deal with trace class and Marchenko operators and Section 8 contains the main result.

2. NOTATION

We adhere to standard terminology from analysis, namely $\mathbb{R}_\pm := [0, \pm\infty)$, \mathbb{C} is the complex plane,

$$\mathbb{C}^+ = \{z \in \mathbb{C} : \text{Im } z > 0\} \quad , \quad i\mathbb{R}_+ = \{z \in \mathbb{C} : z = iy, y \in \mathbb{R}_+\},$$

in the upper half plane,

$$\mathbb{R} + ih = \{z \in \mathbb{C} : z = x + ih, x \in \mathbb{R}\}$$

is the real line shifted h units up. $\|\cdot\|_X$ stands for a norm in a Banach (Hilbert) space X . We use $(S \subseteq \mathbb{R}$ and S will typically be \mathbb{R} or \mathbb{R}_\pm):

- the usual Lebesgue spaces

$$L^p(S) := \left\{ f : \|f\|_{L^p(S)} := \left(\int_S |f(x)|^p dx \right)^{1/p} < \infty \right\} \quad , \quad 1 \leq p < \infty$$

$$L^\infty(S) := \left\{ f : \|f\|_{L^\infty(S)} := \text{esssup}_{x \in S} |f(x)| < \infty \right\},$$

$$L^p_{loc}(S) := \{ \cap L^p(\Delta) : \Delta \subset S, \Delta \text{ compact} \}.$$

- the short range or Faddeev class (important in scattering theory)

$$L^1_1(S) = \left\{ f : \|f\|_{L^1_1(S)} := \int_S (1 + |x|) |f(x)| dx < \infty \right\}.$$

- the Birman-Solomyak spaces ($1 \leq p < \infty$)

$$\ell^\infty(L^p(\mathbb{R}_\pm)) := \left\{ f : \|f\|_{\ell^\infty(L^p(\mathbb{R}_\pm))} := \sup_{x \in \mathbb{R}_\pm} \left(\int_x^{x \pm 1} |f(x)|^p dx \right)^{1/p} < \infty \right\}.$$

Next, \mathfrak{S}_2 denotes the Hilbert-Schmidt class of linear operators A :

$$\mathfrak{S}_2 = \left\{ A : \|A\|_{\mathfrak{S}_2} := [\text{tr}(A^*A)]^{1/2} < \infty \right\}$$

while \mathfrak{S}_1 is the trace class

$$\mathfrak{S}_1 = \left\{ A : \|A\|_{\mathfrak{S}_1} := \text{tr}[(A^*A)^{1/2}] < \infty \right\}.$$

$\text{Spec}(A)$ stands for the spectrum of an operator A and if it is selfadjoint, $\text{Spec}_{ac}(A)$, $\text{Spec}_d(A)$ denote, respectively, the absolutely continuous and discrete components of $\text{Spec}(A)$.

The following portion of notation will be used extensively in reference to the potential q and quantities associated with it. If $\chi_S(x)$ is the characteristic function of a set S , i.e. $\chi_S(x) = 1, x \in S$ and 0 otherwise, then we define:

$$q_+(x) := q(x)\chi_{\mathbb{R}_+}, \quad q_-(x) = q(x)\chi_{\mathbb{R}_-}, \quad \tilde{q}(x) = q(x)\chi_{[-a, a]} \text{ for some } a > 0.$$

We also denote

$$\delta q := q - \tilde{q}$$

and when the cutoff approximation is taken to infinity, i.e. $a \rightarrow \infty$, we write $\delta q \rightarrow 0$.

Any quantity X of arbitrary nature (functions, operators, etc.) related to \tilde{q} will be denoted \tilde{X} and

$$\delta X := X - \tilde{X}.$$

3. WEYL AND SCATTERING SOLUTIONS OF THE SCHRÖDINGER EQUATION

Throughout this section we assume the following.

Hypothesis 3.1. *The potential q is real, locally integrable and such that²*

- (1) q is limit point case at $-\infty$
- (2) q_+ is Faddeev class (short range) and hence has the Jost solution at $+\infty$.

Condition (1) means that the equation

$$-\partial_x^2 u + qu = k^2 u \quad (3.1)$$

has a unique, up to a multiplicative constant, solution Ψ_- , called Weyl, such that $\Psi_-(x, k) \in L^2(\mathbb{R}_-)$ for any $k^2 \in \mathbb{C}^+$. Condition (2) implies that q is limit point case at $+\infty$ and the Weyl solution $\Psi_+(x, k)$ can be taken to have the asymptotic behavior:

$$\Psi_+(x, k) = e^{ikx} + o(1) \quad , \quad x \rightarrow \infty \quad (3.2)$$

for all real³ k .

In particular, we have as in classical scattering theory $\text{Spec}_{ac}(-\partial_x^2 + q_+) = \mathbb{R}_+$. Furthermore, $\Psi_+, \overline{\Psi}_+$ are both solutions to (3.1) for a.e. real k and linearly independent with constant Wronskian

$$W := W(\overline{\Psi}_+(x, k), \Psi_+(x, k)) = 2ik. \quad (3.3)$$

So they form a basis of solutions for (3.1) for a.e. real k and in particular:

$$C(k)\Psi_-(x, k) = \overline{\Psi}_+(x, k) + R(k)\Psi_+(x, k) \quad (3.4)$$

with some $C(k), R(k)$. We call R the reflection coefficient from the right incident. Under our hypothesis, neither C nor R can be analytically continued into the upper half plane. Figure 1 illustrates a potential from our hypothesis and the asymptotic behavior of $C\Psi_-$ at $+\infty$.




FIGURE 1. Scattering channels for $q = q_- + q_+$

We now consider separately scattering solutions corresponding to q_{\pm} . I.e., first by our hypothesis at $-\infty$, there is a solution $\varphi_-(x, k)$ to

$$-\partial_x^2 u + q_- u = k^2 u$$

²For terminology used without explanation, see e.g. [9, 19].

³the Weyl solution Ψ_+ coincides in this case with the Jost solution.

of the form: ($k \in \mathbb{R}$)

$$\varphi_-(x, k) = \begin{cases} D(k)\Psi_-(x, k) & , \quad x < 0 \\ e^{-ikx} + R_-(k)e^{ikx} & , \quad x \geq 0 \end{cases}$$

with some $D(k), R_-(k)$ (see Figure 2).




FIGURE 2. Scattering channels for q_-

For q_+ , there exist particular Jost solutions $\varphi_{\ell,+}$ and $\varphi_{r,+}$ to


$$-\partial_x^2 u + q_+ u = k^2 u$$

such that: ($k \in \mathbb{R}$)

$$T_+(k)\varphi_{\ell,+}(x, k) = \begin{cases} e^{ikx} + L_+(k)e^{-ikx} & , \quad x < 0 \\ T_+(k)\Psi_+(x, k) & , \quad x \geq 0 \end{cases}$$

$$T_+(k)\varphi_{r,+}(x, k) = \begin{cases} T_+(k)e^{-ikx} & , \quad x < 0 \\ \overline{\Psi_+(x, k)} + R_+(k)\Psi_+(x, k) & , \quad x \geq 0 \end{cases}$$

for k real and where T_+ is called the transmission coefficient and L_+, R_+ the reflection coefficients from the left (right) incident. Because q_+ is short range, T_+ can be analytically continued in the upper half plane and has only a finite number of poles $\{i\kappa_n^+\}_{n=1}^N$. Since, in addition, q_+ is supported on the right half line, L_+ can also be analytically extended to the upper half plane and shares the same poles as T_+ . However, in general, R_+ can not be extended off the real axis. The asymptotic behavior of these solutions is illustrated in Figure 3.


 FIGURE 3. Scattering channels for q_+

Note that all right reflection and transmission coefficients can be expressed in terms of Wronskians. Of particular interest, we have for a.e. real k :

$$\begin{aligned}
 W &= 2ik = W(\Psi_+(x, k), C(k)\Psi_-(x, k)) \\
 R(k) &= \frac{W(\overline{\Psi_+(x, k)}, C(k)\Psi_-(x, k))}{W} \\
 R_-(k) &= \frac{W(e^{-ikx}, \varphi_-(x, k))}{W} \Big|_{x \geq 0} \\
 R_+(k) &= \frac{W(\overline{\Psi_+(x, k)}, T_+(k)\varphi_{r,+}(x, k))}{W} \Big|_{x \geq 0} \\
 \frac{1}{T_+(k)} &= \frac{W(\varphi_{r,+}(x, k), \varphi_{\ell,+}(x, k))}{W}
 \end{aligned}$$

Note that any truncation $\tilde{q} = q|_{[-a, a]}$ is compactly supported which implies that \tilde{R}, \tilde{R}_+ can be analytically continued into \mathbb{C}^+ [9] except at a finite number of poles. These poles are located on $i\mathbb{R}_+$ such that their squares correspond respectively to the discrete spectrum of $-\partial_x^2 + \tilde{q}$ and $-\partial_x^2 + \tilde{q}_+$.

4. THE TITCHMARSH-WEYL m -FUNCTION

In this section, we review properties of the Titchmarsh-Weyl m -function which is the logarithmic derivative of the Weyl solutions $\Psi_{\pm}(x, k)$ as $x \rightarrow \pm 0$. It will be a central object in redefining scattering quantities in the next section. We will have to impose some additional conditions on the potential q . Most of the material already appeared in [16] but are repeated here for the reader's convenience.

Definition 4.1. *The Titchmarsh-Weyl m -function is defined by:*

$$m_{\pm}(k^2) = \pm \frac{\partial_x \Psi_{\pm}(x, k)}{\Psi_{\pm}(x, k)} \Big|_{x=\pm 0}$$

Some of the important properties of the m -function are (see e.g. [14, 16]):

- m_{\pm} is analytic for all $k^2 \in \mathbb{C}^+$ and has the Herglotz property, i.e. $m_{\pm} : \mathbb{C}^+ \rightarrow \mathbb{C}^+$
- symmetry $m_{\pm}(\bar{z}) = \overline{m_{\pm}(z)}$

- the singularities of m_{\pm} correspond to the spectrum of the half line Dirichlet Schrödinger operator, i.e. $-\partial_x^2 + q$ on \mathbb{R}_{\pm} with $u(\pm 0) = 0$.
- the Borg-Marchenko uniqueness theorem: $m_1 = m_2 \Rightarrow q_1 = q_2$.

The following representation of m_{\pm} will be useful.

Proposition 4.2. *Let q be a real function on \mathbb{R} such that $q \in \ell^\infty(L^2(\mathbb{R}_-)) \cap L^1(\mathbb{R}_+)$. Let $\gamma = \max(\gamma_-, \gamma_+)$ where*

$$\begin{aligned}\gamma_- &= \max\left(\sqrt{2\|q_-\|_{\ell^\infty(L^2(\mathbb{R}_-))}}, e\|q_-\|_{\ell^\infty(L^2(\mathbb{R}_-))}\right), \\ \gamma_+ &= \frac{\|q_+\|_{L^1(\mathbb{R}_+)}}{2}.\end{aligned}$$

Then for $k = \alpha + ih$, $h > \gamma$,

$$m_{\pm}(k^2) = ik \mp \int_0^{\pm\infty} e^{\pm 2ikx} A_{\pm}(x) dx \quad (4.1)$$

with some real function $A_{\pm}(x)$, called the A -amplitude. The integral in (4.1) is absolutely convergent and the A -amplitude has the following properties.

- (1) $A_{\pm} - q_{\pm}$ is continuous on \mathbb{R}_{\pm} and for $\pm x > 0$:

$$|A_{\pm}(x) - q_{\pm}(x)| \leq \left(\pm \int_0^x |q_{\pm}(s)| ds\right)^2 e^{\pm 2\gamma x} \quad (4.2)$$

- (2) If $q_1, q_2 \in \ell^\infty(L^2(\mathbb{R}_-)) \cap L^1(\mathbb{R}_+)$ then

$$q_1(x) = q_2(x) \text{ on } [0, \pm a] \Rightarrow A_1(x) = A_2(x) \text{ on } [0, \pm a]. \quad (4.3)$$

- (3) For any $h > \gamma$,

$$\|e^{\mp 2hx} A_{\pm}(x)\|_{L^1(\mathbb{R}_{\pm})} \leq C(h, q_{\pm}) < \infty$$

and $C(h, q_{\pm})$ is a nonincreasing function of h .

- (4) For any $h > \gamma$,

$$e^{2hx} A_-(x) \in L^2(\mathbb{R}_-).$$

Proof. The representation (4.1) appeared in [14] for short range q 's and in [18] for more general q 's. Properties (4.2)-(4.3) were derived for $q_+ \in L^1(\mathbb{R}_+)$ in [18] then for $q_+ \in \ell^\infty(L^1(\mathbb{R}_+))$ in [5] but since

$$m_-(q_-(x), k^2) = m_+(q_-(-x), k^2)$$

and $\ell^\infty(L^2(\mathbb{R}_-)) \subset \ell^\infty(L^1(\mathbb{R}_-))$, we have adjusted the results accordingly. So only (3)-(4) require a proof. We will consider A_- and $p = 1, 2$. Using Minkowski's inequality, one needs only to show $e^{2hx} q_-(x)$ and $e^{2hx}(A_-(x) - q_-(x))$ are in $L^p(\mathbb{R}_-)$.

Dropping the subscripts, we have

$$\begin{aligned}
 \int_{-\infty}^0 |e^{2hx} q(x)|^p dx &= \int_{-\infty}^0 e^{2hpx} |q(x)|^p dx \\
 &= \sum_{n=0}^{\infty} \int_{-n-1}^{-n} e^{2hpx} |q(x)|^p dx \\
 &\leq \sum_{n=0}^{\infty} e^{-2hpn} \int_{-n-1}^{-n} |q(x)|^p dx \\
 &\leq \sum_{n=0}^{\infty} e^{-2hpn} \|q(x)\|_{\ell^\infty(L^p(\mathbb{R}_-))}^p \\
 &= \frac{1}{1 - e^{-2hp}} \|q(x)\|_{\ell^\infty(L^p(\mathbb{R}_-))}^p.
 \end{aligned}$$

For the next term, we will make use of the following: ($x \leq 0$)

$$\begin{aligned}
 \int_x^0 |q(s)| ds &\leq \sum_{n=1}^{-\lfloor x \rfloor} \int_{-n}^{-n+1} |q(s)| ds \leq \sum_{n=1}^{-\lfloor x \rfloor} \|q\|_{\ell^\infty(L^1(\mathbb{R}_-))} \\
 &\leq (1-x) \|q\|_{\ell^\infty(L^1(\mathbb{R}_-))} \leq (1-x) \|q\|_{\ell^\infty(L^p(\mathbb{R}_-))}.
 \end{aligned}$$

where the last inequality is a direct consequence of Hölder's inequality. So,

$$\begin{aligned}
 \int_{-\infty}^0 |e^{2hx} (A(x) - q(x))|^p dx &= \int_{-\infty}^0 e^{2hpx} |A(x) - q(x)|^p dx \\
 &\leq \int_{-\infty}^0 e^{2p(h-\gamma)x} \left(\int_x^0 |q(s)| ds \right)^{2p} dx \\
 &\leq \int_{-\infty}^0 e^{2p(h-\gamma)x} (1-x)^{2p} \|q\|_{\ell^\infty(L^p(\mathbb{R}_-))}^{2p} dx.
 \end{aligned}$$

One readily verifies that for any $m = 0, 1, 2, \dots$ and $b > 0$

$$\int_{-\infty}^0 (1-x)^m e^{bx} dx = \frac{m!}{b^{m+1}} \sum_{k=0}^m \frac{b^k}{k!}.$$

Therefore, we have

$$\int_{-\infty}^0 |e^{2hx} (A(x) - q(x))|^p dx \leq \frac{2p!}{[2p(h-\gamma)]^{2p+1}} \sum_{j=0}^{2p} \frac{[2p(h-\gamma)]^j}{j!} \|q\|_{\ell^\infty(L^p(\mathbb{R}_-))}^{2p}.$$

So (3)-(4) are verified with

$$C(h, q_-) = \frac{1}{1 - e^{-2h}} \|q_-(x)\|_{\ell^\infty(L^1(\mathbb{R}_-))} + \frac{1}{4(h-\gamma)^3} \sum_{j=0}^2 \frac{[2(h-\gamma)]^j}{j!} \|q_-\|_{\ell^\infty(L^1(\mathbb{R}_-))}^2.$$

Similarly for A_+ , we can take in (3)

$$C(h, q_+) = \|q_+\|_{L^1(\mathbb{R}_+)} + \frac{1}{h-\gamma} \|q_+\|_{L^1(\mathbb{R}_+)}^2. \quad \square$$

Remark 4.3. In the case of a truncated potential \tilde{q} , since $\tilde{\gamma}_\pm \leq \gamma_\pm$ and $C(h, \tilde{q}_\pm) \leq C(h, q_\pm)$, all above results remain true for the same h . If, in addition, $q_+ \in$

$L^2_{loc}(\mathbb{R}_+)$ then $\tilde{q}_+ \in \ell^\infty(L^2(\mathbb{R}_+))$ and thus $e^{-2hx}\tilde{A}_+(x) \in L^2(\mathbb{R}_+)$ for h large enough.

Corollary 4.4. *Let $q \in \ell^\infty(L^2(\mathbb{R}_-))$, and let $h \geq \gamma_-$ where γ_- is defined as in Proposition 4.2. Then*

- (1) $ik - m_-(k^2) \in L^2(\mathbb{R} + ih)$,
- (2) if $\tilde{q}_- = q_-|_{[-a,0]}$ for some $a > 0$, then

$$\delta m_-(k^2) := m_-(k^2) - \tilde{m}_-(k^2) \rightarrow 0 \text{ in } L^2(\mathbb{R} + ih) \quad , \quad a \rightarrow \infty.$$

Proof. Note that for $k = \alpha + ih$ where $\alpha \in \mathbb{R}, h > \gamma_-$,

$$ik - m_-(k^2) = \int_{-\infty}^0 e^{-2i\alpha x} e^{2hx} A_-(x) dx$$

where by Proposition 4.2, $e^{2hx} A_-(x) \in L^2(\mathbb{R} + ih)$. The Plancherel formula in our case takes the form:

$$\left\| \int_{\mathbb{R}_-} e^{-2i\alpha x} f(x) dx \right\|_{L^2(\mathbb{R})} = \sqrt{\pi} \|f(x)\|_{L^2(\mathbb{R}_-)}$$

and hence

$$\|ik - m_-(k^2)\|_{L^2(\mathbb{R} + ih)} = \sqrt{\pi} \|e^{2hx} A_-(x)\|_{L^2(\mathbb{R}_-)} < \infty$$

In the case $\tilde{q}_- = q_-|_{[-a,0]}$, we have $\tilde{\gamma}_- \leq \gamma_-$ and by Proposition 4.2 (2),

$$\delta m_-(k^2) = \int_{-\infty}^{-a} e^{-2ikx} \delta A(x) dx$$

and therefore for any $h > \gamma_-$,

$$\|\delta m_-(k^2)\|_{L^2(\mathbb{R} + ih)} = \sqrt{\pi} \|e^{2hx} \delta A(x)\|_{L^2((-\infty, -a])} \rightarrow 0 \quad , \quad a \rightarrow \infty. \quad \square$$

5. THE REFLECTION AND TRANSMISSION COEFFICIENTS

In this section we establish some properties of one of our main objects:

$$G(k) := \Delta R(k) = R(k) - R_+(k).$$

As mentioned when first introduced, neither R nor R_+ can be analytically extended to the upper half plane for a potential q under the conditions of Hypothesis 3.1 (or those in Proposition 4.2). But by rewriting G exclusively in terms of R_- , T_+ , and L_+ , we will see that G can be analytically extended to the upper half plane. We also derive key properties of R_- , T_+ , L_+ and G in \mathbb{C}^+ which will be used later to recover $q_-(x)$ – assuming R, R_+ are known.

First we rewrite the reflection and transmission coefficients in terms of the m -function. Setting

$$m_\pm = m_\pm(k^2 + i0) := \lim_{\epsilon \rightarrow 0^+} m_\pm(k^2 + i\epsilon),$$

we have:

$$R(k) = -\frac{m_- + \overline{m}_+ \overline{\Psi_+(0, k)}}{m_- + m_+ \Psi_+(0, k)}, \quad (5.1)$$

$$R_+(k) = -\frac{\overline{m}_+ + ik \overline{\Psi_+(0, k)}}{m_+ + ik \Psi_+(0, k)}, \quad (5.2)$$

$$R_-(k) = \frac{ik - m_-}{ik + m_-}, \quad (5.3)$$

$$L_+(k) = \frac{ik - m_+}{ik + m_+}, \quad (5.4)$$

$$T_+(k) = \frac{2ik}{(ik + m_+) \Psi_+(0, k)}. \quad (5.5)$$

The above are obtained using continuity of the various solutions and their derivatives in x at the point $x = 0$ or, alternately, the Wronskians. Recall that the above are defined for a.e. real k .

From (5.3)-(5.4) and properties of the Titchmarsh-Weyl m -function, one notes that L_+, R_- have a meromorphic extension to the upper half plane. By Proposition 4.2 (3), L_+, R_- are smooth on $\mathbb{R} + ih$ for any $h > \gamma$. Furthermore, by the Borg-Marchenko uniqueness theorem L_+, R_- determine uniquely respectively $q_\pm(x)$ [14].

Now, using (3.3) and (5.1)-(5.5), one readily verifies that:

$$G(k) = \frac{T_+^2(k) R_-(k)}{1 - L_+(k) R_-(k)} \quad (5.6)$$

and thus G can be analytically extended to the upper half plane (recall that T_+ also has an analytic extension since q_+ is short range [9]).

Proposition 5.1. *Let q be as in Proposition 4.2 and let*

$$h_\pm = \inf \left\{ h : h > \gamma_\pm \text{ and } C(h, q_\pm) < \frac{h}{2} \right\}$$

Then for all $h > \max(h_+, h_-)$

- (1) $R_-, L_+ \in (L^\infty \cap L^2)(\mathbb{R} + ih)$ with their L^∞ norm no greater than $1/3$.
- (2) $\delta L_+ \rightarrow 0$ in $L^\infty(\mathbb{R} + ih)$ when $a \rightarrow \infty$.
- (3) $R_- \in L^1(\mathbb{R} + ih)$.
- (4) $\delta R_- \rightarrow 0$ in $L^1(\mathbb{R} + ih)$.

Proof. The above results are direct consequences of Proposition 4.2. For all $k \in \mathbb{R} + ih$, such that $h > \max(h_+, h_-)$,

$$\begin{aligned} |ik - m_\pm(k^2)| &= \left| \int_0^{\pm\infty} e^{\pm 2ikx} A_\pm(x) dx \right| \\ &\leq \|e^{\mp 2hx} A_\pm(x)\|_{L^1(\mathbb{R}_\pm)} \leq \frac{h}{2} \\ |ik + m_\pm(k^2)| &= \left| 2ik \mp \int_0^{\pm\infty} e^{2ikx} A_\pm(x) dx \right| \\ &\geq 2|k| \cdot \left| 1 - \frac{1}{2|k|} \int_0^{\pm\infty} e^{\pm 2ikx} A_\pm(x) dx \right| \\ &\geq \frac{3|k|}{2} \end{aligned}$$

where we have used $|k| \geq h$. Thus $\frac{1}{ik+m_{\pm}(k^2)} \in L^2(\mathbb{R} + ih)$ and it follows immediately from (5.3)-(5.4) that $\|R_-(k)\|_{L^\infty(\mathbb{R}+ih)}, \|L_+(k)\|_{L^\infty(\mathbb{R}+ih)} \leq \frac{1}{3}$ and $L_+, R_- \in L^2(\mathbb{R} + ih)$.

We further obtain $R_- \in L^1(\mathbb{R} + ih)$ by the Cauchy-Schwartz inequality for $h > h_-$:

$$\|R_-(k)\|_{L^1(\mathbb{R}+ih)} \leq \left\| \frac{1}{ik+m_-(k^2)} \right\|_{L^2(\mathbb{R}+ih)} \cdot \|ik - m_-(k^2)\|_{L^2(\mathbb{R}+ih)}.$$

By Remark 4.3, the above is also true for \tilde{R}_-, \tilde{L}_+ .

Now for all $k \in \mathbb{R} + ih$ with $h > h_+$,

$$\begin{aligned} |\delta L_+(k)| &= \left| \frac{-2ik\delta m_+(k^2)}{(ik+m_+(k^2))(ik+\tilde{m}_+(k^2))} \right| \leq \frac{8}{9h} |\delta m_+(k^2)| \\ &\leq \frac{8}{9h} \|e^{-2hx}\delta A_+(x)\|_{L^1([a,\infty))} \rightarrow 0 \quad , \quad a \rightarrow \infty. \end{aligned}$$

We also have

$$\delta R_-(k) = \frac{-2ik\delta m_-(k^2)}{(ik+m_-(k^2))(ik+\tilde{m}_-(k^2))}.$$

Using L^∞ norms and the Cauchy-Schwartz inequality:

$$\begin{aligned} \|\delta R_-\|_{L^1(\mathbb{R}+ih)} &\leq \left\| \frac{-2ik}{ik+\tilde{m}_-(k^2)} \right\|_{L^\infty(\mathbb{R}+ih)} \cdot \left\| \frac{\delta m_-(k^2)}{ik+m_-(k^2)} \right\|_{L^1(\mathbb{R}+ih)} \\ &\leq \frac{4}{3} \left\| \frac{1}{ik+m_-(k^2)} \right\|_{L^2(\mathbb{R}+ih)} \cdot \|\delta m_-(k^2)\|_{L^2(\mathbb{R}+ih)} \end{aligned}$$

and the right hand side, by Corollary 4.4, goes to zero when $a \rightarrow \infty$. \square

Remark 5.2. *Property (3) will play a crucial role. Note that if $q(x) = c\delta(x)$, where δ is Dirac's δ -function, then*

$$R(k) = \frac{c}{2ik - c}$$

which is not in $L^1(\mathbb{R} + ih)$. This suggests that the condition $\ell^\infty(L^2(\mathbb{R}))$ may not be relaxed to read $\ell^\infty(L^1(\mathbb{R}))$.

Corollary 5.3. *For any finite z , and q_-, h under the conditions of Proposition 5.1, $e^{2ikz}R_-(k) \in L^1(\mathbb{R} + ih)$.*

Proof. Immediately follows from $\|e^{2ikz}R_-(k)\|_{L^1(\mathbb{R}+ih)} = e^{-2hz}\|R_-(k)\|_{L^1(\mathbb{R}+ih)}$. \square

While trivial, the above corollary plays an important part in our arguments. The reflection coefficient for the shifted potential $q(x+z)$ is $R(k)e^{2ikz}$ where $R(k)$ is the reflection coefficient corresponding to $q(x)$.

Lemma 5.4. *Let $q_+ \in L^1_1(\mathbb{R}_+)$ and let $h > \beta$ where*

$$\beta = 2 \max \left\{ \|q_+\|_{L^1_1(\mathbb{R}_+)}, 1 \right\}.$$

Then

$$\|T_+(k)\|_{L^\infty(\mathbb{R}+ih)}, \left\| \tilde{T}_+(k) \right\|_{L^\infty(\mathbb{R}+ih)} \leq 2^\beta$$

and $\delta T_+(k) \rightarrow 0$ in $L^\infty(\mathbb{R} + ih)$ as $a \rightarrow \infty$.

Proof. The following are well-known facts (e.g. [9]) for q_+ short range and supported on \mathbb{R}_+ :

- (1) $T_+(k)$ is analytic in \mathbb{C}^+ except at a finite number of simple poles $\{i\kappa_n^+\}_{n=1}^N$ where

$$N \leq 1 + \int_{\mathbb{R}_+} |x| |q_+(x)| dx. \quad (5.7)$$

- (2) $|T_+(k)|^2 + |L_+(k)|^2 = 1$ for a.e. real k and $T_+(-k) = \overline{T_+(k)}$.
 (3) $T_+(k)$ admits the following representation for any $k \in \mathbb{C}^+$:

$$T_+(k) = \prod_{n=1}^N \frac{k + i\kappa_n^+}{k - i\kappa_n^+} \exp\left(\frac{i}{\pi} \int_{\mathbb{R}} \frac{\log |T_+(\omega)|^{-1}}{\omega - k} d\omega\right).$$

We also have the Lieb-Thirring inequality [20]

$$\sum_{n=1}^N \kappa_n^+ \leq L_{1/2,1} \int_{\mathbb{R}_+} |q_+(x)| dx \quad (5.8)$$

where $1/2 \leq L_{1/2,1} \leq 1.005$. Thus, for any $k \in \mathbb{R} + ih$, $h > \beta$

$$\begin{aligned} |T_+(k)| &= \left| \prod_{n=1}^N \frac{k + i\kappa_n^+}{k - i\kappa_n^+} \right| \exp\left(\frac{1}{\pi} \int_{\mathbb{R}} \operatorname{Re} \frac{i}{\omega - k} \log |T_+(\omega)|^{-1} d\omega\right) \\ &= \prod_{n=1}^N \sqrt{1 + \frac{4h\kappa_n^+}{(h - \kappa_n^+)^2}} \exp\left(\frac{-h}{\pi} \int_{\mathbb{R}} \frac{\log |T_+(\omega)|^{-1}}{h^2 + (\omega - \alpha)^2} d\omega\right). \end{aligned}$$

Since $|T_+(\omega)| \leq 1$ for a.e. real ω , the above becomes

$$|T_+(k)| \leq \prod_{n=1}^N \sqrt{1 + \frac{4h\kappa_n^+}{(h - \kappa_n^+)^2}}.$$

But by (5.8), we also have for each n :

$$\kappa_n^+ \leq L_{1/2,1} \|q_+\|_{L^1(\mathbb{R}_+)} \leq \frac{2}{3}\beta < \frac{2}{3}h \quad (5.9)$$

and hence for all $k \in \mathbb{R} + ih$ where $h > \beta$

$$|T_+(k)| \leq \left(\frac{11}{3}\right)^{\beta/2} \leq 2^\beta \quad (5.10)$$

where we have used $N \leq \beta$ from (5.7). Inequalities (5.7)-(5.10) are also valid for $\tilde{T}_+(k)$ and thus T_+, \tilde{T}_+ are uniformly bounded on $\mathbb{R} + ih$.

From [9], we now use the following results

$$\frac{1}{T_+(k)} = 1 - \frac{1}{2ik} \int_{\mathbb{R}} q_+(x) y_+(x, k) dx$$

where $y_+(x, k) := e^{-ikx} \varphi_{\ell,+}(x, k)$ satisfies for $\operatorname{Im} k \geq 0$,

$$y_+(x, k) = 1 + \int_x^\infty D_k(t-x) q_+(t) y_+(t, k) dt, \quad D_k(y) = \frac{e^{2iky} - 1}{2ik}, \quad (5.11)$$

$$|y_+(x, k)| \leq K(\beta)(1 + |x|)$$

where K is a constant depending only on β . Note that

$$\delta T_+(k) = \frac{T_+(k)\tilde{T}_+(k)}{2ik} \left[\int_{\mathbb{R}} q_+(x)\delta y_+(x, k)dx + \int_{\mathbb{R}} \tilde{y}_+(x, k)\delta q_+(x)dx \right]. \quad (5.12)$$

From (5.11),

$$\delta y_+(x, k) = \int_x^\infty D_k(t-x)\tilde{y}_+(t, k)\delta q_+(t)dt + \int_x^\infty D_k(t-x)q_+(t)\delta y_+(t, k)dt \quad (5.13)$$

and since for $k \neq 0$, $|D_k(y)| \leq \frac{1}{|k|}$ for all $y \geq 0$, $\text{Im } k \geq 0$,

$$\begin{aligned} |\delta y_+(x, k)| &\leq \frac{K(\beta)}{|k|} \|\delta q_+\|_{L^1_+(\mathbb{R}_+)} + \int_x^\infty \frac{|q_+(t)|}{|k|} |\delta y_+(t, k)| dt \\ &\leq \frac{K(\beta)}{|k|} e^{\frac{\beta}{2|k|}} \|\delta q_+\|_{L^1_+(\mathbb{R}_+)} \quad , \quad k \neq 0 \quad , \quad \text{Im } k \geq 0 \end{aligned}$$

by iteration on the Volterra integral equation for $\frac{\delta y_+(x, k)k}{K(\beta)\|\delta q_+\|_{L^1_+(\mathbb{R}_+)}}$ derived from (5.13).

Hence for (5.12), we have

$$\|\delta T_+(k)\|_{L^\infty(\mathbb{R}+ih)} \leq 2^{2\beta} \frac{K(\beta)}{\beta} \|\delta q_+\|_{L^1_+(\mathbb{R}_+)}$$

where the right hand side goes to zero for $a \rightarrow \infty$ by the dominated convergence theorem. \square

Proposition 5.5. *Under Hypothesis 3.1,*

$$\Delta R(k) := R(k) - R_+(k)$$

is analytic in \mathbb{C}^+ except on a set

$$\mathcal{S} = \{i\mathcal{z}_n^+\} \cup \sigma \subset i\mathbb{R}_+$$

where

$$\left\{ -(\mathcal{z}_n^+)^2 \right\} = \text{Spec}_d(-\partial_x^2 + q_+) \quad , \quad \sigma = \{ \lambda : \lambda^2 \in \text{Spec}(-\partial_x^2 + q) \cap \mathbb{R}_- \}.$$

Proof. By direct computation, we have

$$G(k) = T_+(k) \frac{ik - m_-(k^2)}{m_+(k^2) + m_-(k^2)} g(k) \quad (5.14)$$

where

$$g(k) = \frac{T_+(k)}{1 + L_+(k)} = \frac{1}{\Psi_+(0, k)}.$$

We gather the following facts:

- (1) it is well-known that $T_+(k), L_+(k)$ are analytic in $\mathbb{C}^+ \setminus \{i\mathcal{z}_n^+\}_{n=1}^N$ where $\left\{ -(\mathcal{z}_n^+)^2 \right\}$ is the negative simple discrete spectrum of $-\partial_x^2 + q_+$. So we also have that $g(k)$ is meromorphic in \mathbb{C}^+ but with poles different from those of T_+, L_+ . Poles of $g(k)$ correspond to the poles of m_+ , i.e. κ 's such that $\Psi_+(0, \kappa) = 0$.
- (2) recall that m_\pm is analytic in \mathbb{C}^+ except for some singularities⁴ $\kappa \in i\mathbb{R}_+$, hence so is $\frac{ik - m_-(k^2)}{(m_+ + m_+)(k^2)}$. Note that $m_+(k^2) + m_-(k^2) = 0$ corresponds

⁴Singularities of m_+ are a finite number poles since q_+ is short range [14] whereas the set of singularities of m_- , while bounded by γ_- , need not be made of poles – it could be continuous.

to $W(\psi_-, \psi_+) = 0$. But then if these two solutions to $\partial_x^2 + q$ are linearly dependent, then $\psi_{\pm}(x, k) \in L^2(\mathbb{R})$ and so $\lambda \in \sigma$.

- (3) we now consider singularities of m_{\pm} . Those cases correspond exactly to $\psi_{\pm}(0, \kappa) = 0$ with $\psi'_{\pm}(0, \kappa) \neq 0$. But if $\psi_-(0, \kappa) = 0$, we can assume $\psi_+(0, \kappa) \neq 0$ (otherwise $\kappa \in \sigma$) and so $m_+(\kappa^2)$ finite. Therefore, by (5.14),

$$G(\kappa) = -\frac{T(\kappa)}{\psi'_+(0, \kappa)}$$

is finite unless $\kappa \in \{i\mathcal{X}_n^+\}$. Now if $\psi_+(0, \kappa) = 0$, we have by (5.5) that ($\kappa \neq 0$)

$$T_+(\kappa) = \frac{2i\kappa}{\psi'_+(0, \kappa)}$$

is finite and by (5.14) since we can assume $m_-(\kappa^2)$ is finite, then

$$G(\kappa) = T_+(\kappa) \frac{i\kappa + m_-(\kappa^2)}{\psi'_+(0, \kappa)}$$

is finite too.

Thus, we find that $G(k)$ is analytic in $\mathbb{C}^+ \setminus (\{i\mathcal{X}_n^+\} \cup \sigma)$. \square

Remark 5.6. *By Proposition 4.2, we have $\gamma > \sup |\mathcal{S}|$ and thus $G(k)$ is smooth on $\mathbb{R} + ih$ for any $h > \gamma$. Note that σ need not be finite, but is bounded.*

Proposition 5.7. *Let q be a real function on \mathbb{R} such that $q \in \ell^\infty(L^2(\mathbb{R}_-)) \cap L^1_1(\mathbb{R}_+)$ and let $h > h_0$ where $h_0 = \max(h_+, h_-, \beta)$ where h_{\pm} are as in Proposition 5.1 and β as in Lemma 5.4. Then*

$$G(k) = \Delta R(k) \in L^1(\mathbb{R} + ih) \quad \text{and} \quad \delta G(k) \rightarrow 0 \text{ in } L^1(\mathbb{R} + ih).$$

Proof. Omitting the variable k for brevity, from (5.6), Proposition 5.1 and Lemma 5.4, we have:

$$\|G\|_{L^1(\mathbb{R} + ih)} \leq \frac{9}{8} \cdot 2^{2h_0} \cdot \|R_-\|_{L^1(\mathbb{R} + ih)} < \infty.$$

By direct computation,

$$\delta G = \frac{G_1 \delta L_+ + G_2 \delta T_+ + G_3 \delta R_-}{(1 - L_+ R_-)(1 - \tilde{L}_+ \tilde{R}_-)} \quad (5.15)$$

where

$$G_1 = T_+^2 R_- \tilde{R}_- \quad , \quad G_2 = R_-(1 - L_+ \tilde{R}_-)(T_+ + \tilde{T}_+) \quad , \quad G_3 = \tilde{T}_+^2.$$

By Proposition 5.1,

$$\left\| \frac{1}{(1 - L_+ R_-)(1 - \tilde{L}_+ \tilde{R}_-)} \right\|_{L^\infty(\mathbb{R} + ih)} \leq \left(\frac{9}{8} \right)^2$$

so it is enough to show that each term inside the brackets in (5.15) goes to zero in $L^1(\mathbb{R} + ih)$ as $a \rightarrow \infty$. Indeed

$$\begin{aligned} \|G_1 \delta L_+\|_{L^1(\mathbb{R} + ih)} &\leq \frac{2^{2h_0}}{3} \cdot \|R_-\|_{L^1(\mathbb{R} + ih)} \cdot \|\delta L_+\|_{L^\infty(\mathbb{R} + ih)} \rightarrow 0, \\ \|G_2 \delta T_+\|_{L^1(\mathbb{R} + ih)} &\leq \frac{10}{9} \cdot 2^{2h_0} \|R_-\|_{L^1(\mathbb{R} + ih)} \cdot \|\delta T_+\|_{L^\infty(\mathbb{R} + ih)} \rightarrow 0, \\ \|G_3 \delta R_-\|_{L^1(\mathbb{R} + ih)} &\leq 2^{2h_0} \cdot \|\delta R_-\|_{L^1(\mathbb{R} + ih)} \rightarrow 0. \end{aligned} \quad \square$$

Corollary 5.8. *Let z be a fixed real parameter and let q, h be as in Proposition 5.7. Then $G_z(k) := e^{2ikz} \Delta R(k) \in L^1(\mathbb{R} + ih)$ and $\delta G_z(k) \rightarrow 0$ in $L^1(\mathbb{R} + ih)$.*

Proof. Note that $G_z(k)$ correspond to the shifted potential $q(x + z)$. For such potential, $R_-(k)$ becomes $R_-(k)e^{2ikz}$, $T_+(k)$ remains the same and $L_+(k)$ becomes $L(k)e^{-2ikz}$. So by Corollary 5.3 and Proposition 5.7, $G_z(k) \in L^1(\mathbb{R} + ih)$ and $\delta G_z(k) \rightarrow 0$ in $L^1(\mathbb{R} + ih)$. \square

6. A TRACE CLASS OPERATOR

In this section we introduce a lemma which appeared in a more general form in [17] and will be a central argument in the main result of this paper in the next section.

Proposition 6.1. *Let A be smooth on $\mathbb{R} + ih$ for some $h > 0$, and $A \in L^1(\mathbb{R} + ih)$. Then the integral operator \mathbb{A} on $L^2(\mathbb{R}_+)$ with kernel*

$$\mathbf{A}(x, y) = \int_{\mathbb{R} + ih} e^{ik(x+y)} A(k) \frac{dk}{2\pi}, \quad x, y \geq 0$$

is trace class, and

$$\|\mathbb{A}\|_{\mathfrak{S}_1} \leq \frac{1}{4\pi h} \|A\|_{L^1(\mathbb{R} + ih)}.$$

Proof. Denote $A_h(\alpha) = A(\alpha + ih)$ and $\widehat{f}(z) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{ikz} f(k) dk$. Then rewrite \mathbb{A} as an operator on $L^2(\mathbb{R})$ by considering $x, y \in \mathbb{R}$ and:

$$\mathbf{A}(x, y) = \chi(x) e^{-h(x+y)} \widehat{A}_h(x+y) \chi(y)$$

where χ is the characteristic function on \mathbb{R}_+ . By convolution and a change of variable, we have:

$$\begin{aligned} \widehat{A}_h(x+y) &= \left(\widehat{\sqrt{A_h}} * \widehat{\sqrt{A_h}} \right) (x+y) \\ &= \int_{\mathbb{R}} \widehat{\sqrt{A_h}}(x-s) \widehat{\sqrt{A_h}}(y+s) ds. \end{aligned}$$

So $\mathbb{A} = \mathbb{A}_1 \mathbb{A}_2$ where $\mathbb{A}_1, \mathbb{A}_2$ are operators on $L^2(\mathbb{R})$ with kernels

$$\mathbf{A}_1(x, s) = \chi(x) e^{-hx} \widehat{\sqrt{A_h}}(x-s),$$

$$\mathbf{A}_2(s, y) = \chi(y) e^{-hy} \widehat{\sqrt{A_h}}(y+s).$$

One readily has

$$\begin{aligned} \|\mathbb{A}_k\|_{\mathfrak{S}_2}^2 &= \iint_{\mathbb{R}^2} |\mathbf{A}_k(\xi, \eta)|^2 d\xi d\eta \\ &= \int_{\mathbb{R}} \chi(z) e^{-2hz} dz \int_{\mathbb{R}} \left| \widehat{\sqrt{A_h}}(S) \right|^2 dS \\ &= \frac{1}{4\pi h} \left\| \widehat{\sqrt{A_h}} \right\|_{L^2(\mathbb{R})}^2 = \frac{1}{4\pi h} \|A_h\|_{L^1(\mathbb{R})} \end{aligned}$$

where we have used the Plancherel equality $\|\widehat{f}\|_2^2 = \frac{1}{2\pi} \|f\|_2^2$ and hence

$$\|\mathbb{A}\|_{\mathfrak{S}_1} \leq \|\mathbb{A}_1\|_{\mathfrak{S}_2} \|\mathbb{A}_2\|_{\mathfrak{S}_2} = \frac{1}{4\pi h} \|A_h\|_{L^1(\mathbb{R})} = \frac{1}{4\pi h} \|A\|_{L^1(\mathbb{R} + ih)}. \quad \square$$

7. CLASSICAL MARCHENKO INVERSE SCATTERING

In this section we review well-known facts about how to recover a potential q in the Faddeev class from the scattering data associated with the Schrödinger operator $-\partial_x^2 + q(x)$ via the Marchenko inverse scattering procedure (see e.g. [1, 9, 10]).

For $q \in L^1_+(\mathbb{R})$, the scattering data consisting of

- the discrete spectrum $\{-\varkappa_n^2\}_{n=1}^N$ of the Schrödinger operator $-\partial_x^2 + q(x)$ on $L^2(\mathbb{R})$,
- norming constants $\{c_n\}_{n=1}^N$ associated to the bound states of the Schrödinger operator,
- and the reflection coefficient $R(k)$, $k \in \mathbb{R}$

determine together the potential uniquely. By the inverse scattering procedure

$$q(x) = -2\partial_x k_x(0^+), \quad (7.1)$$

where $k_x \in L^2(\mathbb{R}_+)$ solves the Marchenko equation

$$k_x(y) + M_x(y) + \int_0^\infty M_x(y+z)k_x(z)dz = 0 \quad , \quad y > 0 \quad (7.2)$$

with

$$M_x(\cdot) = M(\cdot + 2x), \quad (7.3)$$

$$M(s) = \sum_{n=1}^N c_n^2 e^{-\varkappa_n s} + \frac{1}{2\pi} \int_{\mathbb{R}} e^{iks} R(k) dk. \quad (7.4)$$

If we define the Marchenko operator on $L^2(\mathbb{R}_+)$ as:

$$(\mathbb{M}_x f)(y) = \int_0^\infty M_x(y+s)f(s)ds \quad , \quad f \in L^2(\mathbb{R}_+) \quad (7.5)$$

then (7.2) becomes

$$(1 + \mathbb{M}_x)k_x(y) = -M_x(y)$$

and $1 + \mathbb{M}_x$ is boundedly invertible [9].

Assuming the Fredholm determinant in (7.5) is well-defined, one can also rewrite (7.1) as [10]

$$q(x) = -2\partial_x^2 \log \det(1 + \mathbb{M}_x), \quad (7.6)$$

known as the Bargmann, Dyson, or determinant formula (see e.g. [16]). The determinant is well-defined if \mathbb{M}_x is trace class. However, we don't know if it is the case for a generic short range potential.

We choose to detour this fact. To this end, we express the Marchenko kernel in a different form. Recall the well-known fact (see e.g. [1]) that if a short range potential is supported on \mathbb{R}_- , then $R(k)$ can be analytically continued in \mathbb{C}^+ , its poles are $\{i\varkappa_n\}_{n=1}^N$ and

$$\text{Res}(R(k), i\varkappa_n) = ic_n^2,$$

where c_n is the norming constant associated to \varkappa_n in the scattering data.

So for any $h > \max\{\varkappa_n\}$, one can deform the contour [16] in (7.4) and by the residue theorem rewrite the Marchenko kernel (7.4)

$$M(s) = \frac{1}{2\pi} \int_{\mathbb{R}+ih} e^{iks} R(k) dk. \quad (7.7)$$

Note that (7.7) can then be used for any compactly supported q 's with compact support using a shifting argument.

Remark 7.1. We will consider a potential which is locally square integrable on the line, in $\ell^\infty(L^2(\mathbb{R}_-))$ and such that q_+ is Faddeev class. The classical inverse scattering results do not apply directly since q is not short range and the negative part of the spectrum of $-\partial_x^2 + q(x)$ need not be finite. However, since $q \in L^2_{loc}(\mathbb{R}) \subset L^1_{loc}(\mathbb{R})$, we have

- $\tilde{q} \in L^1_1(\mathbb{R})$ so the classical inverse scattering procedure applies to the truncated potential,
- \tilde{q} is compactly supported so we can use (7.7),
- $\tilde{q} \in \ell^\infty(L^2(\mathbb{R}))$ which we will show implies that $\tilde{\mathbb{M}}_x$ is trace class and so (7.6) applies.

The above will be a basis for our limiting procedure.

8. MAIN RESULT

We now present our main result which gives a formula to recover a nondecaying unknown potential q_- assuming that q_+ and the reflection coefficient R are known.

Theorem 8.1. Let q be a real, locally square integrable potential on \mathbb{R} such that $(q_\pm = q|_{\mathbb{R}_\pm})$

- $\sup_{x \leq 0} \int_{x-1}^x |q_-(s)|^2 ds < \infty$ (uniformly in L^2_{loc}),
- $\int_{\mathbb{R}_+} (1+x)|q_+(x)| dx < \infty$ (short range)

and let $R(k)$, $R_+(k)$ be the right reflection coefficient corresponding to q, q_+ respectively.

Let \mathbb{M}_x^+ be the Marchenko operator associated with the scattering data

$$\{R_+(k), -(\varkappa_n^+)^2, c_n^+\}_{k \in \mathbb{R}, 1 \leq n \leq N}$$

for q_+ (given by (7.3)-(7.5)) and let \mathbb{G}_x be the Hankel integral operator associated with $R - R_+$. I.e.

$$(\mathbb{G}_x f)(y) = \int_0^\infty \mathbb{G}_x(y+s)f(s)ds \quad , \quad f \in L^2(\mathbb{R}_+), \quad (8.1)$$

$$\mathbb{G}_x(s) = \frac{1}{2\pi} \int_{\mathbb{R}+ih} e^{ik(s+2x)}(R - R_+)(k)dk \quad (8.2)$$

with some $h > 0$ sufficiently large.

Then for any $x < 0$

$$q_-(x) = -2\partial_x^2 \log \det \left(1 + (1 + \mathbb{M}_x^+)^{-1} \mathbb{G}_x \right) \quad (8.3)$$

with the determinant defined in the classical Fredholm sense.

Remark 8.2. Theorem 8.1 solves the inverse scattering problem for a steplike potential with the knowledge of its short range part. Indeed, given (short range) q_+ one solves the direct scattering problem and finds the scattering data

$$\{R_+(k), -(\varkappa_n)^2, c_n\}_{k \in \mathbb{R}, 1 \leq n \leq N}$$

for q_+ . Then, we construct \mathbb{M}_x^+ by (7.3)-(7.5) and given the (right) reflection coefficient for the whole potential q , one constructs by (8.1)-(8.2) the Hankel operator \mathbb{G}_x . The unknown (non decaying) part of q is recovered for each $x < 0$ by (8.3).

Proof. We will first prove the statement for \tilde{q} . For a fixed $a > 0$, \tilde{q} is compactly supported. Hence, \tilde{R} can be analytically continued in \mathbb{C}^+ except at a finite number of poles $\{i\tilde{z}_n\}_{n=1}^{\tilde{N}}$, and the Marchenko kernel (7.4) becomes

$$\tilde{M}(s) = \frac{1}{2\pi} \int_{\mathbb{R}+ih} e^{iks} \tilde{R}(k) dk \quad , \quad h > \max\{\tilde{z}_n\}_{n=1}^{\tilde{N}}.$$

Define $q_a(x) = \tilde{q}(x+a)$ as in Figure 4. Then q_a is supported on \mathbb{R}_- and $q_a \in \ell^\infty(L^2(\mathbb{R}_-))$.




FIGURE 4. Shifted potential $q_a(x) = \tilde{q}(x+a)$

Now its reflection coefficient $R_{a,-}(k) \in L^1(\mathbb{R}+ih)$ for $h > h_a$ where h_a is defined as in Proposition 5.1 for q_a . But $\tilde{R}(k) = R_{a,-}(k)e^{-2ika}$, so they share the same poles and by Corollary 5.3, $\tilde{R}(k), \tilde{R}(k)e^{2ikx} \in L^1(\mathbb{R}+ih)$ for $h > h_a$. We also have that $\tilde{R}(k)e^{2ikx}$ is smooth on $\mathbb{R}+ih$ for $h > \max\{\tilde{z}_n\}_{n=1}^{\tilde{N}}$ so by Proposition 6.1, \tilde{M}_x is trace class. Hence the following Bargmann formula applies:

$$\tilde{q}(x) = -2\partial_x^2 \log \det(1 + \tilde{M}_x). \quad (8.4)$$

Now write

$$\tilde{R} = \tilde{R}_+ + \tilde{G} \quad , \quad \tilde{G} = \Delta\tilde{R} = \tilde{R} - \tilde{R}_+$$

and split the Marchenko operator accordingly⁵:

$$\tilde{M}_x = \tilde{M}_x^+ + \tilde{G}_x.$$

The same h_a is enough to ensure $\tilde{M}_x^+ \in \mathfrak{S}_1$ since $\tilde{R}_+(k)e^{2ika}$ corresponds to $\tilde{q}_a(x)$ (above the shaded region in Figure 4).

In addition, for $h > h_0$ where h_0 is the same⁶ as in Proposition 5.7, $\tilde{G}, G, \delta G \in L^1(\mathbb{R}+ih)$ and by Proposition 5.5 and the subsequent remark, we also have $\tilde{G}, G, \delta G$ smooth on $\mathbb{R}+ih$. Since by Corollary 5.8, the same applies to $\tilde{G}_x, G_x, \delta G_x$, we can apply Proposition 6.1 and conclude that $\tilde{G}_x, G_x, \delta G_x \in \mathfrak{S}_1$.

Therefore, first we rewrite the Bargmann formula (8.4) as:

$$\begin{aligned} \tilde{q}(x) &= -2\partial_x^2 \log \det(1 + \tilde{M}_x^+ + \tilde{G}_x) \\ &= -2\partial_x^2 \log \det(1 + \tilde{M}_x^+) - 2\partial_x^2 \log \det(1 + (1 + \tilde{M}_x^+)^{-1}\tilde{G}_x) \end{aligned} \quad (8.5)$$

where we have used the fact from classical Marchenko theory that $1 + \tilde{M}_x^+$ is boundedly invertible. But

$$\tilde{q}_+(x) = -2\partial_x^2 \log \det(1 + \tilde{M}_x^+)$$

⁵Because \tilde{q}_+ is compactly supported, \tilde{M}_x^+ can be equivalently expressed as (7.4) or (7.7).

⁶Note that h_0 is independent of a .

and $q_+(x) = 0$ for $x < 0$. So (8.5) becomes for $x < 0$:

$$\tilde{q}_-(x) = -2\partial_x^2 \log \det(1 + (1 + \tilde{\mathbb{M}}_x^+)^{-1} \tilde{\mathbb{G}}_x). \quad (8.6)$$

Now, we now use the fact that $(1 + \mathbb{M}_x)$ remains boundedly invertible in its limit so the right hand side of (8.3) is well-defined. In addition, by Proposition 6.1, for $h > h_0$

$$\|\delta \mathbb{G}_x\|_{\mathfrak{S}_1} \leq \frac{1}{4\pi h} \|\delta G_x\|_{L^1(\mathbb{R}+ih)}$$

and the right hand side of the inequality goes to zero by Corollary 5.8 for $a \rightarrow \infty$. Therefore, we find indeed that the limit of (8.6) is (8.3). \square

Note that if $\mathbb{M}_x^+ \in \mathfrak{S}_1$ then (8.3) simplifies to

$$q(x) = -2\partial_x^2 \log \det(1 + \mathbb{M}_x), \quad x \in \mathbb{R}, \quad (8.7)$$

where $\mathbb{M}_x = \mathbb{M}_x^+ + \mathbb{G}_x$. It is, of course, well-known (see e.g. [9]) that under our condition on q_+ , $\mathbb{M}_x \in \mathfrak{S}_2$ but we couldn't prove it for \mathfrak{S}_1 . We were unable to find a rigorous proof of such a statement in the literature either. (It is typically assumed (frequently implicitly) or referred to as "too involved".) However, since $\mathbb{M}_x \in \mathfrak{S}_2$, then $\det(1 + \mathbb{M}_x)$ can, in fact, be regularized differently from (8.3) (see [17] for details).

In conclusion, we emphasize that the fact that (8.3) is understood in the classical sense is indeed quite important as it guarantees the convergence of various types of approximation of $(1 + \mathbb{M}_x^+)^{-1} \mathbb{G}_x$ in trace norm. This, in turn, means a certain stability of the inverse problem algorithm based upon (8.3).

REFERENCES

- [1] Aktosun, T.; Klaus M. Chapter 2.2.4: *Inverse theory: problem on the line*. In: E. R. Pike and P. C. Sabatier (eds.), *Scattering*, Academic Press, London, 2001, pp. 770-785.
- [2] Aktosun, T.; Sacks, P. *Inversion of reflectivity data for nondecaying potentials*. *SIAM J. Appl. Math.* 60 (2000), no. 4, 1340-56.
- [3] Aktosun, T. *Inverse Schrödinger scattering on the line with partial knowledge of the potential*. *SIAM J. Appl. Math.* 56 (1996), 219-31.
- [4] Aktosun, T.; Klaus, M.; van der Mee, C. *On the Riemann-Hilbert problem for the one dimensional Schrödinger equation*. *J. Math. Phys.* 34 (1993), 2651-90.
- [5] Avdonin, S.; Mikhaylov, V.; Rybkin, A. *The boundary control approach to the Titchmarsh-Weyl m -function. I. The response operator and the A -amplitude*. *Comm. Math. Phys.* 275 (2007), no. 3, 791-803.
- [6] Braun, M.; Sofianos, S.; Lipperheide, R. *One-dimensional Marchenko inversion in the presence of bound states*. *Inverse Problems* 11 (1995), L1-3.
- [7] Buslaev, V.; Fomin, V. *An inverse scattering problem for the one-dimensional Schrödinger equation on the entire axis*. *Vestnik Leningrad Univ.* 17 (1962), 56-64. [In Russian]
- [8] Cohen, A.; Kappeler, T. *Scattering and inverse scattering for steplike potentials in the Schrödinger equation*. *Indiana Univ. Math. J.* 34 (1985), 127-80.
- [9] Deift, P.; Trubowitz, E. *Inverse scattering on the line*. *Comm. Pure Appl. Math.* 32 (1979), no. 2, 121-251.
- [10] Faddeev, L. D. *Inverse problem of quantum scattering theory. II*. *J. Sov. Math.* 5 (1976), 334-96.
- [11] Gesztesy, F.; Simon, B. *Inverse spectral analysis with partial information on the potential. I. The case of an a.c. component in the spectrum*. Papers honouring the 60th birthday of Klaus Hepp and of Walter Hunziker, Part II (Zürich, 1995), *Helv. Phys. Acta* 70 (1997), no. 1-2, 66-71.
- [12] Grébert, B.; Weder, R. *Reconstruction of a potential on the line that is a priori known on the half line*. *SIAM J. Appl. Math.* 55 (1995), 242-54.

- [13] Hruslov, Ě. Ja. *Asymptotic behavior of the solution of the Cauchy problem for the Korteweg-de Vries equation with steplike initial data*. Mat. Sb. (N.S.) 99(141) (1976), no. 2, 261–81, 296.
- [14] Ramm, A. G. *Property C for ODE and applications to inverse problems.*. In: Operator Theory and Its Applications, A.G. Ramm, P. N. Shivakumar, A. V. Strauss (eds), Amer. Math. Soc., Fields Institute Communications V. 25, Providence, 2000, pp. 15-75.
- [15] Rundell, W.; Sacks P. *On the determination of potential without bound state data*. J. Comput. Appl. Math. 55 (1994), 325–47.
- [16] Rybkin, A. *On the Marchenko inverse scattering procedure with partial information on the potential*. Inverse Problems 25 (2009), no. 9, 095011 (34pp).
- [17] Rybkin, A. *The Hirota τ -function and well-posedness of the KdV equation with an arbitrary step like initial profile decaying on the right half line*, preprint (2010).
- [18] Simon, B. *A new approach to inverse spectral theory, I. Fundamental formalism*. Ann. of Math. 150 (1999), no. 2, 1029–57.
- [19] Titchmarsh, E.C. *Eigenfunction expansions associated with second-order differential equations*. Oxford University Press, London, 1950, 184 pp.
- [20] Weidl, T. *On the Lieb-Thirring constants $L_{\gamma,1}$ for $\gamma \geq 1/2$* . Comm. Math. Phys. 178 (1996), no. 1, 135–46.

UNIVERSITY OF ALASKA FAIRBANKS

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF ALASKA FAIRBANKS, PO BOX 756660, FAIRBANKS, AK 99775

E-mail address: `orbastille@alaska.edu`, `arybkin@alaska.edu`