

CERTIFIED METAMODELS FOR SENSITIVITY INDICES ESTIMATION

ALEXANDRE JANON¹, MAËLLE NODET¹ AND CLÉMENTINE PRIEUR¹

Abstract. Global sensitivity analysis of a numerical code, more specifically estimation of Sobol indices associated with input variables, generally requires a large number of model runs. When those demand too much computation time, it is necessary to use a reduced model (metamodel) to perform sensitivity analysis, whose outputs are numerically close to the ones of the original model, while being much faster to run. In this case, estimated indices are subject to two kind of errors: sampling error, caused by the computation of the integrals appearing in the definition of the Sobol indices by a Monte-Carlo method, and metamodel error, caused by the replacement of the original model by the metamodel. In cases where we have certified bounds for the metamodel error, we propose a method to quantify both types of error, and we compute confidence intervals for first-order Sobol indices.

Résumé. L'analyse de sensibilité globale d'un modèle numérique, plus précisément l'estimation des indices de Sobol associés aux variables d'entrée, nécessite généralement un nombre important d'exécutions du modèle à analyser. Lorsque celles-ci requièrent un temps de calcul important, il est judicieux d'effectuer l'analyse de sensibilité sur un modèle réduit (ou métamodèle), fournissant des sorties numériquement proches du modèle original mais pour un coût nettement inférieur. Les indices estimés sont alors entachés de deux sortes d'erreur: l'erreur d'échantillonnage, causée par l'estimation des intégrales définissant les indices de Sobol par une méthode de Monte-Carlo, et l'erreur de métamodèle, liée au remplacement du modèle original par le métamodèle. Lorsque nous disposons de bornes d'erreurs certifiées pour le métamodèle, nous proposons une méthode pour quantifier les deux types d'erreurs et fournir des intervalles de confiance pour les indices de Sobol du premier ordre.

1. CONTEXT

1.1. Monte-Carlo estimation of first-order Sobol indices

Let $Y = f(X_1, \dots, X_p)$ be our (scalar) output of interest, where the input variables X_1, \dots, X_p are modelised as independent random variables of known distribution. For $i = 1, \dots, p$, we recall the first-order Sobol index:

$$S_i = \frac{\text{Var}(\mathbf{E}(Y|X_i))}{\text{Var}(Y)}$$

which measures, on a scale of 0 to 1, the fraction of the total variability of the output caused by the variability in X_i alone.

As f is generally implicitly known (f can *e.g.* be a functional of a solution of a partial differential equation parametrized by functions of the input variables X_1, \dots, X_p), one has no analytical expression for S_i and has to resort to numerical estimation. The variances in the definition of S_i can be expressed as multidimensional

¹ Laboratoire Jean Kuntzman, Université Joseph Fourier (Grenoble I), INRIA/MOISE, 51 rue des Mathématiques, BP 53, 38041 Grenoble cedex 9, France

integrals over the input parameters space. We use Monte-Carlo estimates for multidimensional integrals: let $\{\mathbf{X}^k\}_{k=1,\dots,N}$ and $\{\mathbf{X}'^k\}_{k=1,\dots,N}$ be two independent, identically distributed vector samples of $\mathbf{X} = (X_1, \dots, X_p)$. For $k = 1, \dots, N$, we note:

$$y_k = f(\mathbf{X}^k) \quad \text{and} \quad y'_k = f(X_1'^k, \dots, X_{i-1}'^k, X_i^k, X_{i+1}'^k, \dots, X_p'^k)$$

We take the following statistical estimator for S_i , introduced in [2]:

$$\widehat{S}_i(\mathcal{E}) = \frac{\frac{1}{N} \sum_{k=1}^N y_k y'_k - \left(\frac{1}{N} \sum_{k=1}^N y_k \right) \left(\frac{1}{N} \sum_{k=1}^N y'_k \right)}{\frac{1}{N} \sum_{k=1}^N (y_k)^2 - \left(\frac{1}{N} \sum_{k=1}^N y_k \right)^2}$$

where $\mathcal{E} = (\{\mathbf{X}^k\}_{k=1,\dots,N}, \{\mathbf{X}'^k\}_{k=1,\dots,N})$ is our couple of samples the estimator depends on.

1.2. Reduced basis metamodels

In order to apply the reduced basis metamodelling, we further assume that the output $f(\mathbf{X})$ depends on a function $u(\mathbf{X})$ where, for every input \mathbf{X} , $u(\mathbf{X})$ satisfies a \mathbf{X} -dependent partial differential equation (PDE).

To make things clear, let us consider an example: we take $p = 2$, so that $\mathbf{X} = (X_1, X_2)$, and take for $u(t, x; X_1, X_2)$ the solution of the following (X_1, X_2) -dependent initial-boundary value problem (viscous Burgers' equation):

$$\begin{cases} \frac{\partial u}{\partial t} + \frac{1}{2} \frac{\partial}{\partial x} (u^2) - \nu \frac{\partial^2 u}{\partial x^2} = 1 \\ u(t=0, x) = u_{0m}^2 + 5 \sin(0.5x) \quad \forall x \in [0; 1] \\ u(t, x=0) = b_0 \\ u(t, x=1) = b_1 \end{cases} \quad (1)$$

where our input parameters are $(X_1, X_2) = (\nu, u_{0m})$, and b_0 and b_1 are so that we have *compatibility conditions*: $b_0 = u_{0m}^2$ and $b_1 = u_{0m}^2 + 5 \sin(0.5)$.

This problem can be analyzed by means of the Cole-Hopf substitution (see [3] for instance), which turns the equation of interest into heat equation, leading to an integral representation of u and well-posedness for $u \in C^0([0, T], H^1(\cdot, 1))$.

Note that the x symbol denotes the spatial variable $u(\mathbf{X})$ depends on, and is unrelated with the parameters X_1 and X_2 . Our output can be, for instance: $f(\mathbf{X}) = \int_0^T \int_0^1 u(t, x, \mathbf{X}) dx dt$.

For a given value of \mathbf{X} , $u(\mathbf{X})$ is generally approximated using a numerical method, such as the finite-element method. These methods work by searching for $u(\mathbf{X})$ in a linear subspace of high dimension \mathcal{N} ; this leads to a large linear system (or a succession of linear systems) to solve for the coefficients of (the approximation of) $u(\mathbf{X})$ in a fixed basis of X . This gives what we call the “full” discrete solution, that we denote again by $u(\mathbf{X})$. Even if efficient methods have been developed to solve the linear systems arising from such discretizations, the large number of unknowns that are to be found is often responsible for large computation times.

The reduced basis method is based on the fact that \mathcal{N} has to be large because the basis we expand $u(\mathbf{X})$ in does not depend on the PDE problem that is being solved; hence it is too “generic”: it can represent well a large number of functions, but allows much more degrees of freedom than wanted. We split the computation into two phases: the offline phase, where we seek a “reduced space”, whose dimension n is much smaller than \mathcal{N} , and which is suitable for effectively representing $u(\mathbf{X})$ for various values of the input parameter \mathbf{X} ; and the online phase, where, for each required value of the input parameters, we solve the “projected” PDE on the reduced space.

This method is interesting if we are to solve the PDE for a number of values of the parameter sufficiently large so that the fixed cost of the offline phase is cancelled by the gain in marginal cost offered by the online phase *vs.* the standard discretization. This is often the case with Monte-Carlo estimations.

One crucial feature of the reduced basis method, which we will rely on later, is that it provides a *certified error bound* $\epsilon_u(\mathbf{X}, t)$, which satisfies ($\|\cdot\|$ being the usual norm on $L^2([0, 1])$):

$$\|u(\mathbf{X}; t) - \tilde{u}(\mathbf{X}; t)\| \leq \epsilon_u(\mathbf{X}, t) \quad \forall \mathbf{X}, \forall t \in [0; T]$$

and, of course, ϵ_u can be *fully* and *quickly* computed (with a computation time of the same order of magnitude than the one for \tilde{u}). This error bound on u can lead to an error bound ϵ on the output:

$$\left| f(\mathbf{X}) - \tilde{f}(\mathbf{X}) \right| \leq \epsilon(\mathbf{X}) \quad \forall \mathbf{X}$$

where $\tilde{f}(\mathbf{X})$ denotes a functional of the reduced solution.

One can turn to [6] for a detailed introduction to the reduced basis method, and to [4] for the extension to the viscous Burgers equation (1).

2. CONSTRUCTION OF COMBINED CONFIDENCE INTERVALS

2.1. Metamodel error

For a couple of samples $\mathcal{E} = (\{\mathbf{X}^k\}_{k=1, \dots, N}, \{\mathbf{X}'^k\}_{k=1, \dots, N})$, we can use our metamodel output \tilde{f} and our metamodel error bound ϵ to compute, for $k = 1, \dots, N$:

$$\tilde{y}_k = \tilde{f}(\mathbf{X}^k), \quad \tilde{y}'_k = \tilde{f}(X_1'^k, \dots, X_{i-1}'^k, X_i^k, X_{i+1}'^k, \dots, X_p'^k)$$

and:

$$\epsilon_k = \epsilon(\mathbf{X}^k), \quad \epsilon'_k = \epsilon(X_1'^k, \dots, X_{i-1}'^k, X_i^k, X_{i+1}'^k, \dots, X_p'^k)$$

In [5], we show that we can compute rigorous and accurate bounds \widehat{S}_i^m and \widehat{S}_i^M depending only on the $\tilde{y}_k, \tilde{y}'_k, \epsilon_k$ and ϵ'_k (thus requiring no evaluation of the full model) so that:

$$\widehat{S}_i^m(\mathcal{E}) \leq \widehat{S}_i(\mathcal{E}) \leq \widehat{S}_i^M(\mathcal{E})$$

where $\widehat{S}_i(\mathcal{E})$ is the (unknown) value of the estimator of S_i computed on the couple of samples \mathcal{E} .

2.2. Combined confidence intervals

To take sampling error in account, we use a bootstrap procedure (see [1]) on the two bounding estimators \widehat{S}_i^m and \widehat{S}_i^M . More specifically, we draw N numbers with repetition from $\{1, 2, \dots, N\}$, so as to get a random list L . We then get two *bootstrap replications* by computing \widehat{S}_i^m and \widehat{S}_i^M using the samples couple $(\{\mathbf{X}^k\}_{k \in L}, \{\mathbf{X}'^k\}_{k \in L})$ instead of $(\{\mathbf{X}^k\}_{k=1, \dots, N}, \{\mathbf{X}'^k\}_{k=1, \dots, N})$. We repeat those computations for a fixed number B of times, so as to obtain B couples of replications $S_i^{m,1}, \dots, S_i^{m,B}$ and $S_i^{M,1}, \dots, S_i^{M,B}$. Now, for a fixed risk level $\alpha \in]0; 1[$, let S_i^{inf} and S_i^{sup} be, respectively, the $\alpha/2$ quantile of $S_i^{m,1}, \dots, S_i^{m,B}$ and S_i^{sup} be the $1 - \alpha/2$ quantile of $S_i^{M,1}, \dots, S_i^{M,B}$. We take $[S_i^{inf}; S_i^{sup}]$ as our combined confidence interval for S_i . This confidence interval accounts for both metamodel and sampling error.

3. NUMERICAL RESULTS

3.1. Target model

Our underlying model is given by the Burgers equation (1). Our output functional is taken to be: $f(\mathbf{X}) = \frac{1}{N} \sum_{i=0}^N u(t = T, x = \frac{i}{N}; \mathbf{X})$.




FIGURE 1. Convergence benchmark for sensitivity index of ν . We plotted, for a fixed sample of size $N = 300$, estimator bounds \widehat{S}^m and \widehat{S}^M , and endpoints of the 95% combined confidence interval, for different reduced basis sizes.

We set $\mathcal{N} = 60$, $\Delta t = .01$, $T = .05$, while the uncertain parameters ν and u_{0m} are assumed to be of uniform distributions, with respective ranges $[1; 20]$ and $[-0.3; 0.3]$. We also take $B = 300$ bootstrap replications and a risk level $\alpha = 0.05$.

Note that more flexible parametrizations of right-hand sides in (1) can be considered; results remain qualitatively the same. We chose this parametrization for simplicity reasons.

3.2. Convergence benchmark

Figure 1 shows the lower \widehat{S}^m and upper \widehat{S}^M bounds for different reduced basis sizes n and fixed sample of size $N = 300$, as well as the endpoints of the combined confidence intervals. This figure exhibits the fast convergence of our bounds to the true value of S_a as the reduced basis size increases. We also see that the part of the error due to sampling (gaps between confidence interval upper bound and upper bound, and between confidence interval lower bound and lower bound) remains constant, as sample size stays the same.

3.3. Comparison with estimation on the full model

To demonstrate the interest of using sensitivity analysis on the reduced model, we computed the combined intervals for the two sensitivity indices using sample size $N = 22000$ and $n = 11$ (those parameters were found using our parameter tuning procedure for a target precision $p = 0.02$). We found $[0.0674128; 0.0939712]$ for sensitivity index for ν , and $[0.914772; 0.926563]$ for sensitivity with respect to u_{0m} . These confidence intervals have mean length: $0.019 \approx 0.02$ as desired. This computation took 58.77 s of CPU time to complete.

To obtain a result of the same precision, we carry a simulation on the *full* model, for $N = 22000$ (sample size can be chosen smaller than before, as there will be no metamodel error); we get a bootstrap confidence interval with mean length of ≈ 0.0193 . This computation takes 294 s of CPU time to complete. Hence, on this example, using a reduced-basis model roughly divides overall computation time by a factor of 5, without any sacrifice on the precision and the rigorousness (as our metamodel error quantification procedure is fully proven and certified) of the confidence interval. We expect higher time savings with more complex (for example, two- or three-dimensional in space) models.

This work has been partially supported by the French National Research Agency (ANR) through COSINUS program (project COSTA-BRAVA no ANR-09-COSI-015).

REFERENCES

- [1] GEB Archer, A. Saltelli, and IM Sobol. Sensitivity measures, ANOVA-like techniques and the use of bootstrap. *Journal of Statistical Computation and Simulation*, 58(2):99–120, 1997.
- [2] T. Homma and A. Saltelli. Importance measures in global sensitivity analysis of nonlinear models. *Reliability Engineering & System Safety*, 52(1):1–17, 1996.
- [3] E. Hopf. The partial differential equation $u_t + u u_x = \mu x x$. *Communications on Pure and Applied Mathematics*, 3(3):201–230, 1950.
- [4] A. Janon, M. Nodet, and C. Prieur. Certified reduced-basis solutions of viscous Burgers equations parametrized by initial and boundary values. Preprint available at <http://hal.inria.fr/inria-00524727/en>, 2010, *submitted*.
- [5] A. Janon, M. Nodet, and C. Prieur. Confidence intervals for sensitivity indices using reduced-basis metamodels. Preprint available at <http://hal.inria.fr/inria-00567977/en>, 2011, *submitted*.
- [6] N.C. Nguyen, K. Veroy, and A.T. Patera. Certified real-time solution of parametrized partial differential equations. *Handbook of Materials Modeling*, pages 1523–1558, 2005.