

HYDROLYSIS OF AZINPHOSMETHYL INDUCED BY THE SURFACE OF SMECTITES

M. SÁNCHEZ-CAMAZANO AND M. J. SÁNCHEZ-MARTÍN

Instituto de Recursos Naturales y Agrobiología, C.S.I.C., Apdo. 257
37071 Salamanca, Spain

Abstract—The effect of layer charge and of the interlayer cations of smectites on the hydrolysis of azinphosmethyl (0,0-dimethyl S-((4-oxo-1,2,3-benzotriazine-3(4H)-yl)methyl) phosphorodithioate) in an aqueous medium was investigated. Ultraviolet spectroscopy was used for monitoring the hydrolysis process. Hydrolysis of the pesticide is catalyzed by Ca-hectorite (layer charge 0.216) but is not catalyzed by Ca-nonttronite nor by Ca-montmorillonites with a layer charge above 0.216. The Mg- and Cu-hectorites and Cu-montmorillonite with a layer charge of 0.264 also show catalytic activity. The catalytic activities of the Ca²⁺ and Cu²⁺ cations as exchange cations of the smectite and as salts are compared. In agreement with previously reported work, the results show that the hydrolysis of azinphosmethyl may involve the adsorption of the pesticide into the interlayer space of the smectites, forming a bidentate complex with the interlayer cations. This interaction must enhance the electrophilic nature of the phosphorus atom, thereby facilitating its nucleophilic attack by the OH⁻ ion and producing rupture of the P-S bond.

Key Words—Hydrolysis, Adsorption, Azinphosmethyl, Smectite, Organophosphorus pesticides.

INTRODUCTION


Clay minerals are able to inactivate adsorbed pesticides by catalyzing their decomposition. In most cases, this degradation occurs through hydrolysis (Hance, 1970; Mortland, 1970; Theng, 1974). This ability of clays to convert toxic organic compounds into their corresponding nontoxic forms may be of great importance if it is taken into account that these minerals are a common component of the colloidal fraction of soils, and also that they are used in the practical formulation of pesticides. Finely-divided clays are often used as inert diluting agents to reduce the concentration of pesticides to levels appropriate for their use in agriculture, and also for improving their handling properties.

The catalytic effect of kaolinite (Saltzman *et al.* 1976; Mingelgrin *et al.*, 1977), montmorillonite (Mortland and Raman, 1967; Sánchez-Camazano and Sánchez-Martín, 1983; Pusino *et al.*, 1988) and some soils (Konrad *et al.*, 1967, 1969; Konrad and Chester, 1969) in the hydrolysis of organophosphorus pesticides has been demonstrated.

A previous work by the authors on different aspects of the adsorption of azinphosmethyl by smectites (Sánchez-Martín and Sánchez-Camazano, 1984) included the study of the adsorption of this pesticide by different minerals from this group in an aqueous medium. In that study, occasional traces of azinphosmethyl hydrolysis were observed. Accordingly, it was thought of interest to investigate the effect of the layer charge and interlayer cations of smectite on the catalytic hydrolysis of this pesticide.

Azinphosmethyl is the common name of the organophosphorus insecticide (0,0-dimethyl S-((4-oxo-

1,2,3-benzotriazine-3(4H)-yl)methyl) phosphorodithioate). Its structural formula is:



MATERIALS AND EXPERIMENTAL METHODS

The clay samples employed in the present work are shown in Table 1. In all cases the <2- μ m fraction was used. Homoionic samples were prepared according to the method described by González-García and González-García (1953). The azinphosmethyl (98% purity), was supplied by Chem Service, West Chester, PA. It is a solid crystalline substance with a melting point of 73°C. Its solubility in water is 29 μ g/ml and it is soluble in most organic solvents (Worthing and Walker, 1987).

To study the influence of the interlayer cations and of the layer charge of the smectite on azinphosmethyl hydrolysis, 10 mg of the different samples (=0.008 meq of exchange cation) were treated with 10 ml of an aqueous solution containing 20 μ g/ml of the pesticide. The suspensions were kept for different periods of time with intermittent shaking in a chamber maintained at 30°C. The suspensions were centrifuged at 5500 rpm for 30 min. Hydrolysis was followed in the supernatant solution by UV spectrophotometry; absorption maximum of azinphosmethyl 228 nm, and absorption maximum of the hydrolysis product at 210 nm. The linear




Figure 1. Hydrolysis of azinphosmethyl by Mg-hectorite as followed by UV spectroscopy: spectra of azinphosmethyl solution free of clay (W.S.) and of azinphosmethyl solution treated with clay over increasing periods of time.

Table 2 shows the results of the hydrolysis of azinphosmethyl in the presence of smectites and salts after 10 days. It should be noted that similar results are observed when using distilled or sterilized water in the experiments, which indicates that there is no microbial degradation. The chemical is hydrolyzed by the Ca-, Mg- and Cu- hectorites. The percentage of hydrolysis for the first two samples is similar. 20% of the hydrolysis produced by the Cu sample is apparent; the true




Figure 3. Hydrolysis of azinphosmethyl by Ca²⁺ and Cu²⁺ cations in solution as followed by UV spectroscopy: spectra of azinphosmethyl solution free of salts (W.S.) and of azinphosmethyl solution treated with CaCl₂ and with CuSO₄ for 10 days.




Figure 2. Hydrolysis of azinphosmethyl by Cu-hectorite as followed by UV spectroscopy: spectra of azinphosmethyl solution free of clay (W.S.) and of azinphosmethyl solution treated with clay over increasing periods of time.

percentage is possibly higher, as will be detailed below. It may be seen that the compound is not hydrolyzed by the Ca-smectites with a charge higher than that of hectorite. Neither is it hydrolyzed by the Tidinit Mg-montmorillonite, whereas it is hydrolyzed by the Cu sample of this mineral (charge 0.264). Additionally, the catalytic activity of Cu²⁺ is greater as a salt in solution than as an interlayer cation, unlike the situation with the Ca²⁺ ion. The results show that the catalytic activity of the smectites is related to the saturating cation and possibly to its layer charge, precisely the factors that generally govern the adsorption of organic molecules in the interlayer space of smectites and, in particular, that of azinphosmethyl (Sánchez-Martín and Sánchez-Camazano, 1984). The interlayer adsorption of the pesticide may therefore be involved in the cat-

Table 2. Hydrolysis of azinphosmethyl after 10 days contact with different homoionic samples and salts (% on initial chemical added).

Sample ¹	Layer charge	Hydrolysis ²
H-Ca	0.216	42
H-Mg	0.216	46
H-Cu	0.216	20
M-Ca	0.264	0
M-Mg	0.264	0
M-Cu	0.264	20
M-Ca	0.350	0
N-Ca	0.335	0
CuSO ₄	—	100
CaCl ₂	—	0
Water solution	—	0

¹ H, hectorite; M, montmorillonite; N, nontronite.

² Observed by UV spectroscopy. The true levels are possibly higher (see text).

alytic process of hydrolysis, in addition to the adsorption on the external planar surfaces of the clays.

Table 3 shows the influence of the interlayer cation of the hectorite on hydrolysis. It is seen that after adsorption equilibrium has been reached, at 24 or 48 hr and, although the percentage of hydrolyzed compound increases, the sample does not adsorb more azinphosmethyl. This suggests that the adsorption sites may continue to be occupied by hydrolysis products, although the product absorbing under UV partly passes into the solution. The Cu-hectorite, which owing to the characteristics of the Cu cation may adsorb more azinphosmethyl and produce a stronger degree of hydrolysis, apparently only hydrolyzes 20%. In a previous work (Sánchez-Martín and Sánchez-Camazano, 1984) on the mechanism of interaction of azinphosmethyl with homoionic samples of montmorillonite, IR spectroscopy findings suggested that the P-S and C-O groups of the pesticide molecule interact simultaneously with the interlayer cations of the phyllosilicate to form a bidentate complex. The Cu sample was an exception; interaction with the cation probably occurred through the P-S group and a nitrogen atom on the triazine ring, also forming a bidentate complex. The strong affinity of the Cu²⁺ cation for electrons, together with this different mechanism of interaction, may allow a greater proportion of hydrolysis products to remain in the interlayer space. The authors have observed that 4-oxo, 3H-1,2,3-benzotriazine is adsorbed into the interlayer space of Cu-hectorite. X-ray and IR results showed that the interaction occurs, like azinphosmethyl, between the Cu²⁺ cation and a nitrogen atom of the triazine ring.

From the foregoing, it may be inferred that the catalytic hydrolysis of azinphosmethyl by smectites must be the result of the formation of the aforementioned bidentate complex. The normal hydrolysis reaction is strongly mediated by the deficiency of electrons in the

Table 3. Adsorption and hydrolysis of azinphosmethyl (% on initial chemical added) by different homoionic hectorites.

Time (h)	Ca hectorite		Mg hectorite		Cu hectorite	
	Adsorption	Hydrolysis ¹	Adsorption	Hydrolysis ¹	Adsorption	Hydrolysis ¹
2	56	0	20	0	32	0
8	58	24	25	0	52	0
18	60	27	30	0	61	0
24	62	27	42	0	71	20
48	62	31	48	41	74	20
72	62	34	48	42	74	20
96	62	37	48	43	74	20
120	62	39	48	46	74	20
192	62	42	48	46	74	20

¹ Observed by UV spectroscopy. The true levels are possibly higher (see text).

phosphorus atom. The interaction in question increases the electrophilic nature of the phosphorus atom by facilitating its nucleophilic attack by the OH⁻ ion, thus leading to rupture of the P-S bond. The strong catalytic activity of the Cu²⁺ cation as a salt in solution may be due to the same easy accessibility for the pesticide. The null catalytic activity of the Ca²⁺ cation in solution is probably due to the fact that under these conditions this cation forms a monodentate rather than a bidentate complex. The bidentate complex would only be formed by the Ca²⁺ cation on the restricted internal surface of the hectorite, governed by the configuration that the molecules necessarily have to adopt owing to the negative surface of oxygen atoms of the phyllosilicate. Many reactions catalyzed by the surface of phyllosilicates involve the selective intercalation of compounds on the restricted surface of these minerals (Theng, 1974; Setton, 1986).

CONCLUSIONS

Hydrolysis of azinphosmethyl in an aqueous medium is catalyzed by different smectites. The catalytic activity of these silicates in the hydrolysis seems to be related to their layer charge and to the nature of the saturating cations. Adsorption of azinphosmethyl on clay surfaces, especially adsorption into the interlayer space, may be involved in the hydrolysis process. The formation of a bidentate interlayer complex between azinphosmethyl and the interlayer cations, in agreement with previously reported work, may strongly enhance the normal hydrolysis reaction of the pesticide in aqueous media.

ACKNOWLEDGMENTS

This work was supported by the Dirección General de Investigación Científica y Técnica (DGICYT, Spain) under Project No. PB87-0240. The authors are grateful to U. Mingelgrin for his suggestions and for critical comments on the manuscript.

REFERENCES

- American Petroleum Institute (1950) *Research Project 49. Clay Minerals Standard: Preliminary Reports 1-8*: Columbia University, New York.
- Eto, M. (1974) *Organophosphorus Pesticides: Organic and Biological Chemistry*: CRC Press, Inc., Cleveland, Ohio, 57-121.
- González-García, F. and González-García, S. (1953) Modificaciones producidas por tratamiento térmico en las propiedades fisicoquímicas de los silicatos de la serie isomorfa montmorillonita-beidellita: *An. Edafol. Agrobiol.* **12**, 925-992.
- Gutiérrez-Ríos, E. and González-García, F. (1949) Génesis de la Montmorillonita de Marruecos español: *An. Edafol. Agrobiol.* **8**, 537-547.
- Hance, R. J. (1970) *Influence of sorption on the decomposition of pesticides. Sorption and transport processes in soils. SCI monograph 37*: Academic Press, New York, 92-107.
- Konrad, J. G. and Chester, G. (1969) Degradation in soils of ciodrin, an organophosphate insecticide: *J. Agric. Food Chem.* **17**, 226-230.
- Konrad, J. G., Armstrong, D. E., and Chester, G. (1967) Soil degradation of diazinon, a phosphorothioate insecticide: *Agron. J.* **59**, 591-594.
- Konrad, J. G., Chester, G., and Armstrong, D. E. (1969) Soil degradation of malathion, a phosphorodithioate insecticide: *Soil. Sci. Soc. Am. Proc.* **33**, 259-262.
- Melnikov, N. N. (1971) Chemistry of pesticides: *Res. Rev.* **36**, 371-372.
- Mingelgrin, U., Saltzman, S., and Yaron, B. (1977) A possible model for the surface-induced hydrolysis of organophosphorus pesticides on kaolinite clays: *Soil Sci. Soc. Am. Proc.* **41**, 519-523.
- Mortland, M. M. (1970) Clay organic-complex and interactions: *Adv. Agron.* **22**, 75-117.
- Mortland, M. M. and Raman, K. V. (1967) Catalytic hydrolysis of some organic phosphate pesticides by copper (II): *J. Agric. Food Chem.* **15**, 163-167.
- Pusino, A., Gessa, C., and Kozlowski, H. (1988) Catalytic hydrolysis of quinalphos on homoionic clays: *Pestic. Sci.* **24**, 1-8.
- Saltzman, S., Mingelgrin, U., and Yaron, B. (1976) Role of water in the hydrolysis of parathion and methylparathion on kaolinite: *J. Agric. Food Chem.* **24**, 739-743.
- Sánchez-Camazano, M. and Sánchez-Martín, M. J. (1983) Montmorillonite-catalyzed hydrolysis of phosmet: *Soil Sci.* **136**, 89-93.
- Sánchez-Martín, M. J. and Sánchez-Camazano, M. (1984) Aspects of the adsorption of azinphosmethyl by smectites: *J. Agric. Food Chem.* **32**, 720-725.
- Setton, R. (1986) *Chemical Reactions in Organic and Inorganic Constrained Systems*: Reidel Publishing Company, Netherlands. 452 pp.
- Theng, B. K. J. (1974) *The Chemistry of Clay-Organic Reactions*: Wiley, New York, 26-281.
- Worthing, C. R. and Walker, S. B. (1987) *The Pesticide Manual. A World Compendium*: The Lavenham Press Ltd., Lavenham, England, 41-42.

(Received 7 February 1991; accepted 24 July 1991; Ms. 2073)