

## ON INCOMPACTNESS FOR CHROMATIC NUMBER OF GRAPHS

SAHARON SHELAH

ABSTRACT. We deal with compactness. Assume the existence of non-reflecting stationary set of cofinality  $\kappa$ . We prove that one can define a graph  $G$  whose chromatic number is  $> \kappa$ , while the chromatic number of every subgraph  $G' \subseteq G, |G'| < |G|$  is  $\leq \kappa$ . The main case is  $\kappa = \aleph_0$ .

---

*Date:* April 5, 2012.

*2010 Mathematics Subject Classification.* Primary: 03E05; Secondary: 05C15.

*Key words and phrases.* set theory, graphs, chromatic number, compactness, non-reflecting stationary sets.

The author thanks Alice Leonhardt for the beautiful typing.

We would like to thank the Israel Science Foundation for partial support of this research. Publication 1006.

## Anotated Content

§0 Introduction

§1 From non-reflecting stationary set.

[We show that “ $S \subseteq S_\kappa^\lambda$  is stationary not reflecting” implies imcompactness for length  $\lambda$  for “chromatic number =  $\kappa$ ”.]

§2 From almost free.

[Here we weaken the assumption in §1 to “ $\mathcal{A} \subseteq {}^\kappa \text{Ord}$  is almost free”.]

## § 0. INTRODUCTION

## § 0(A). The questions and results.

During the Hajnal conference (June 2011) Magidor asked me on incompactness of “having chromatic number  $\aleph_0$ ”; that is, there is a graph  $G$  with  $\lambda$  nodes, chromatic number  $> \aleph_0$  but every subgraph with  $< \lambda$  nodes has chromatic number  $\aleph_0$  when:

- (\*)<sub>1</sub>  $\lambda$  is regular  $> \aleph_1$  with a non-reflecting stationary  $S \subseteq S_{\aleph_0}^\lambda$ , possibly though better not, assuming some version of GCH.

Subsequently also when:

- (\*)<sub>2</sub>  $\lambda = \aleph_{\omega+1}$ .

Such problems were first asked by Erdős-Hajnal, see [EH74]; we continue [Sh:347].

First answer was using BB, see [Sh:309, 3.24] so assuming

- ⊕ (a)  $\lambda = \mu^+$   
 (b)  $\mu^{\aleph_0} = \mu$   
 (c)  $S \subseteq \{\delta < \lambda : \text{cf}(\delta) = \aleph_0\}$  is stationary not reflecting

or just

- ⊕' (a)  $\lambda = \text{cf}(\lambda)$   
 (b)  $\alpha < \lambda \Rightarrow |\alpha|^{\aleph_0} < \lambda$   
 (c) as above.

However, eventually we get more: if  $\lambda = \aleph^{\aleph_0} = \text{cf}(\lambda)$  and  $S \subseteq S_{\aleph_0}^\lambda$  is stationary non-reflective then we have  $\lambda$ -incompactness for  $\aleph_0$ -chromatic. In fact, in §2 we replace  $\aleph_0$  by  $\kappa = \text{cf}(\kappa) < \lambda$  using a stronger hypothesis.

Moreover, if  $\lambda^\kappa > \lambda$  we still get  $(\lambda^\kappa, \lambda)$ -incompactness for  $\aleph_0$ -chromatic number and even  $(\kappa, \aleph_0)$ -chromatic number. In §2 we use quite free family of countable sequences.

In subsequent work we shall solve also the parallel of the second question of Magidor, i.e.

- (\*)<sub>2</sub> for regular  $\kappa > \aleph_0$  colours getting  $\aleph_{\kappa \cdot \varepsilon + 1}$  for every  $\varepsilon < \kappa$ .

We thank Menachem Magidor for asking, Peter Komjath for stimulating discussion and Paul Larson and Shimoni Garti for some comments.

## § 0(B). Preliminaries.

**Definition 0.1.** For a graph  $G$ , let  $\text{ch}(G)$ , the chromatic number of  $G$  be the minimal cardinal  $\chi$  such that there is colouring  $\mathbf{c}$  of  $G$  with  $\chi$  colours, that is  $\mathbf{c}$  is a function from the set of nodes of  $G$  into  $\chi$  or just a set of of cardinality  $\leq \chi$  such that  $\mathbf{c}(x) = \mathbf{c}(y) \Rightarrow \{x, y\} \notin \text{edge}(G)$ .

**Definition 0.2.** 1) We say “we have  $\lambda$ -incompactness for the  $(< \chi)$ -chromatic number” or  $\text{INC}_{\text{chr}}(\lambda, < \chi)$  when: there is a graph  $G$  with  $\lambda$  nodes, chromatic number  $\geq \chi$  but every subgraph with  $< \lambda$  nodes has chromatic number  $< \chi$ .

2) If  $\chi = \mu^+$  we may replace “ $< \chi$ ” by  $\mu$ ; similarly in 0.3.

We also consider

- Definition 0.3.** 1) We say “we have  $(\mu, \lambda)$ -incompactness for  $(< \chi)$ -chromatic number” or  $\text{INC}_{\text{chr}}(\mu, \lambda, < \chi)$  when there is an increasing continuous sequence  $\langle G_i : i \leq \lambda \rangle$  of graphs each with  $\leq \mu$  nodes,  $G_i$  an induced subgraph of  $G_\lambda$  with  $\text{ch}(G_\lambda) \geq \chi$  but  $i < \lambda \Rightarrow \text{ch}(G_i) < \chi$ .
- 2) Replacing (in part (1))  $\chi$  by  $\bar{\chi} = (< \chi_0, \chi_1)$  means  $\text{ch}(G_\lambda) \geq \chi_1$  and  $i < \lambda \rightarrow \text{ch}(G_i) < \chi_0$ ; similarly in 0.2 and parts 3),4) below.
- 3) We say we have incompactness for length  $\lambda$  for  $(< \chi)$ -chromatic (or  $\bar{\chi}$ -chromatic) number when we fail to have  $(\mu, \lambda)$ -compactness for  $(< \chi)$ -chromatic (or  $\bar{\chi}$ -chromatic) number for some  $\mu$ .
- 4) We say we have  $[\mu, \lambda]$ -incompactness for  $(< \chi)$ -chromatic number or  $\text{INC}_{\text{chr}}[\mu, \lambda, < \chi]$  when there is a graph  $G$  with  $\mu$  nodes,  $\text{ch}(G) \geq \chi$  but  $G^1 \subseteq G \wedge |G^1| < \lambda \Rightarrow \text{ch}(G^1) < \chi$ .
- 5) Let  $\text{INC}_{\text{chr}}^+(\mu, \lambda, < \chi)$  be as in part (1) but we add that even the  $\text{cl}(G_i)$ , the colouring number of  $G_i$  is  $< \chi$  for  $i < \lambda$ , see below.
- 6) Let  $\text{INC}_{\text{chr}}^+[\mu, \lambda, < \chi]$  be as in part (4) but we add  $G^1 \subseteq G \wedge |G^1| < \lambda \Rightarrow \text{cl}(G^1) < \chi$ .
- 7) If  $\chi = \kappa^+$  we may write  $\kappa$  instead of “ $< \chi$ ”.

- Definition 0.4.** 1) For regular  $\lambda > \kappa$  let  $S_\kappa^\lambda = \{\delta < \lambda : \text{cf}(\delta) = \kappa\}$ .
- 2) We say  $C$  is a  $(\geq \theta)$ -closed subset of a set  $B$  of ordinals where: if  $\delta = \sup(\delta \cap B) \in B$ ,  $\text{cf}(\delta) \geq \theta$  and  $\delta = \sup(C \cap \delta)$  then  $\delta \in C$ .

**Definition 0.5.** For a graph  $G$ , the colouring number  $\text{cl}(G)$  is the minimal  $\kappa$  such that there is a list  $\langle a_\alpha : \alpha < \alpha(*) \rangle$  of the nodes of  $G$  such that  $\alpha < \alpha(*) \Rightarrow \kappa > |\{\beta < \alpha : \{a_\beta, a_\alpha\} \in \text{edge}(G)\}|$ .

§ 1. FROM NON-REFLECTING STATIONARY IN COFINALITY  $\aleph_0$ 

**Claim 1.1.** *There is a graph  $G$  with  $\lambda$  nodes and chromatic number  $> \kappa$  but every subgraph with  $< \lambda$  nodes have chromatic number  $\leq \kappa$  when:*

- ⊕ (a)  $\lambda, \kappa$  are regular cardinals
- (b)  $\kappa < \lambda = \lambda^\kappa$
- (c)  $S \subseteq S_\kappa^\lambda$  is stationary, not reflecting.

*Proof.* Stage A: Let  $\bar{X} = \langle X_i : i < \lambda \rangle$  be a partition of  $\lambda$  to sets such that  $|X_i| = \lambda$  or just  $|X_i| = |i|^\kappa$  and  $\min(X_i) \geq i$  and let  $X_{<i} = \cup\{X_j : j < i\}$  and  $X_{\leq i} = X_{<(i+1)}$ . For  $\alpha < \lambda$  let  $\mathbf{i}(\alpha)$  be the unique ordinal  $i < \lambda$  such that  $\alpha \in X_i$ . We choose the set of points = nodes of  $G$  as  $Y = \{(\alpha, \beta) : \alpha < \beta < \lambda, \mathbf{i}(\beta) \in S \text{ and } \alpha < \mathbf{i}(\beta)\}$  and let  $Y_{<i} = \{(\alpha, \beta) \in Y : \mathbf{i}(\beta) < i\}$ .

Stage B: Note that if  $\lambda = \kappa^+$ , the complete graph with  $\lambda$  nodes is an example (no use of the further information in ⊕). So without loss of generality  $\lambda > \kappa^+$ .

Now choose a sequence satisfying the following properties, exists by [Sh:g, Ch.III]:

- ⊕ (a)  $\bar{C} = \langle C_\delta : \delta \in S \rangle$
- (b)  $C_\delta \subseteq \delta = \sup(C_\delta)$
- (c)  $\text{otp}(C_\delta) = \kappa$
- (d)  $\bar{C}$  guesses clubs.

Let  $\langle \alpha_{\delta, \varepsilon}^* : \varepsilon < \kappa \rangle$  list  $C_\delta$  in increasing order.

For  $\delta \in S$  let  $\Gamma_\delta$  be the set of sequence  $\bar{\beta}$  such that:

- ⊕ $_{\bar{\beta}}$  (a)  $\bar{\beta}$  has the form  $\langle \beta_\varepsilon : \varepsilon < \kappa \rangle$
- (b)  $\bar{\beta}$  is increasing with limit  $\delta$
- (c)  $\alpha_{\delta, \varepsilon}^* < \beta_{2\varepsilon+i} < \alpha_{\delta, \varepsilon+1}^*$  for  $i < 2, \varepsilon < \kappa$
- (d)  $\beta_{2\varepsilon+i} \in X_{<\alpha_{\delta, \varepsilon+1}^*} \setminus X_{\leq \alpha_{\delta, \varepsilon}^*}$  for  $i < 2, \varepsilon < \kappa$
- (e)  $(\beta_{2\varepsilon}, \beta_{2\varepsilon+1}) \in Y$  hence  $\in Y_{<\alpha_{\delta, \varepsilon+1}^*} \subseteq Y_{<\delta}$  for each  $\varepsilon < \kappa$

(can ask less).

So  $|\Gamma_\delta| \leq |\delta|^\kappa \leq |X_\delta| \leq \lambda$  hence we can choose a sequence  $\langle \bar{\beta}_\gamma : \gamma \in X'_\delta \subseteq X_\delta \rangle$  listing  $\Gamma_\delta$ .

Now we define the set of edges of  $G$ :  $\text{edge}(G) = \{(\alpha_1, \alpha_2), (\min(C_\delta), \gamma) : \delta \in S, \gamma \in X'_\delta \text{ hence the sequence } \bar{\beta}_\gamma = \langle \beta_{\gamma, \varepsilon} : \varepsilon < \kappa \rangle \text{ is well defined and we demand } (\alpha_1, \alpha_2) \in \{(\beta_{\gamma, 2\varepsilon}, \beta_{\gamma, 2\varepsilon+1}) : \varepsilon < \kappa\}\}$ .

Stage C: In every subgraph of  $G$  of cardinality  $< \lambda$  has chromatic number  $\leq \kappa$ .

For this we shall prove that:

$$\oplus_1 \text{ ch}(G|Y_{<i}) \leq \kappa \text{ for every } i < \lambda.$$

This suffice as  $\lambda$  is regular, hence every subgraph with  $< \lambda$  nodes is included in  $Y_{<i}$  for some  $i < \lambda$ .

For this we shall prove more by induction on  $j < \lambda$ :

$\oplus_{2,j}$  if  $i < j, i \notin S, \mathbf{c}_1$  a colouring of  $G \upharpoonright Y_{<i}, \text{Rang}(\mathbf{c}_1) \subseteq \kappa$  and  $u \in [\kappa]^\kappa$  then there is a colouring  $\mathbf{c}_2$  of  $G \upharpoonright Y_{<j}$  extending  $\mathbf{c}_1$  such that  $\text{Rang}(\mathbf{c}_2 \upharpoonright (Y_{<j} \setminus Y_{<i})) \subseteq u$ .

Case 1:  $j = 0$

Trivial.

Case 2:  $j$  successor,  $j - 1 \notin S$

By the induction hypothesis without loss of generality  $j = i + 1$ , but then every node from  $Y_j \setminus Y_i$  is an isolated node in  $G \upharpoonright Y_{<j}$ , because if  $\{(\alpha, \beta), (\alpha', \beta')\}$  is an edge of  $G \upharpoonright Y_j$  then  $\mathbf{i}(\beta), \mathbf{i}(\beta') \in S$  hence necessarily  $\mathbf{i}(\beta) \neq j - 1 = i, \mathbf{i}(\beta') \neq j - 1 = i$  hence both  $(\alpha, \beta), (\alpha, \beta')$  are from  $Y_i$ .

Case 3:  $j$  successor,  $j - 1 \in S$

Let  $j - 1$  be called  $\delta$  so  $\delta \in S$ . But  $i \notin S$  by the assumption in  $\oplus_{2,j}$  hence  $i < \delta$ . Let  $\varepsilon(*) < \kappa$  be such that  $\beta_{\delta, \varepsilon(*)} > i$ .

Let  $\langle u_\varepsilon : \varepsilon \leq \kappa \rangle$  be a sequence of subsets of  $u$ , a partition of  $u$  to sets each of cardinality  $\kappa$ ; actually the only disjointness used is that  $u_\kappa \cap (\bigcup_{\varepsilon < \kappa} u_\varepsilon) = \emptyset$ .

We let  $i_0 = i, i_{1+\varepsilon} = \cup\{\beta_{\delta, \varepsilon(*)+1+\zeta} + 1 : \zeta < 1 + \varepsilon\}, i_\kappa = \delta, i_{\kappa+1} = \delta + 1 = j$ .

Note that:

- $\varepsilon < \kappa \Rightarrow i_\varepsilon \notin S_j$ .

[Why? For  $\varepsilon = 0$  by the assumption on  $i$ , for  $\varepsilon$  successor  $i_\varepsilon$  is successor and for  $i$  limit clearly  $\text{cf}(i_\varepsilon) = \text{cf}(\varepsilon) < \kappa$  and  $S \subseteq S_\kappa^\lambda$ .]

We now choose  $\mathbf{c}_{2,\zeta}$  by induction on  $\zeta \leq \kappa + 1$  such that:

- $\mathbf{c}_{2,0} = \mathbf{c}_1$
- $\mathbf{c}_{2,\zeta}$  is a colouring of  $G \upharpoonright Y_{<i_\zeta}$
- $\mathbf{c}_{2,\zeta}$  is increasing with  $\zeta$
- $\text{Rang}(\mathbf{c}_{2,\zeta} \upharpoonright (Y_{<i_{\zeta+1}} \setminus Y_{<i_\zeta})) \subseteq u_\xi$  for every  $\xi < \zeta$ .

For  $\zeta = 0, \mathbf{c}_{2,0}$  is  $\mathbf{c}_1$  so is given.

For  $\zeta = \varepsilon + 1 < \kappa$ : use the induction hypothesis, possible as necessarily  $i_\varepsilon \notin S$ .

For  $\zeta \leq \kappa$  limit: take union.

For  $\zeta = \kappa + 1$ , note that each node of  $Y_{<i_\zeta} \setminus Y_{<i_\kappa}$  is not connected to any other such node and connected to some node in  $Y_{<i_\kappa}$  has the form  $(\min(C_\delta), \gamma), \gamma \in X'_\delta$ , hence  $\bar{\beta}_\gamma$  is well defined, so the node  $(\min(C_\delta), \gamma)$  is connected in  $G$ , more exactly in  $G \upharpoonright Y_{\leq \delta}$  exactly to the  $\kappa$  nodes  $\{(\beta_{\gamma, 2\varepsilon}, \beta_{\gamma, 2\varepsilon+1}) : \varepsilon < \kappa\}$ , but for every  $\varepsilon < \kappa$  large enough,  $\mathbf{c}_{2,\kappa}((\beta_{\gamma, 2\varepsilon}, \beta_{\gamma, 2\varepsilon+1})) \in u_\varepsilon$  hence  $\notin u_\kappa$  and  $|u_\kappa| = \kappa$  so we can choose a colour.

Case 4:  $j$  limit

By the assumption of the claim there is a club  $e$  of  $j$  disjoint to  $S$  and without loss of generality  $\min(e) = i$ . Now choose  $\mathbf{c}_{2,\xi}$  a colouring of  $Y_{<\xi}$  by induction on  $\xi \in e \cup \{j\}$ , increasing with  $\xi$  such that  $\text{Rang}(\mathbf{c}_{2,\xi} \upharpoonright (Y_{<\xi} \setminus Y_{<i})) \subseteq u, \mathbf{c}_{2,0} = \mathbf{c}_1$

- For  $\xi = \min(e) = i$  the colouring  $\mathbf{c}_{2,\xi} = \mathbf{c}_{2,i} = \mathbf{c}_1$  is given,
- for  $\xi$  successor in  $e$ , i.e.  $\in \text{nacc}(e) \setminus \{i\}$ , use the induction hypothesis with  $\xi, \max(e \cap \xi)$  here playing the role of  $j, i$  there recalling  $\max(e \cap \xi) \in e, e \cap S = \emptyset$
- for  $\xi = \sup(e \cap \xi)$  take union.

Lastly, for  $\xi = j$  we are done.

Stage D:  $\text{ch}(G) > \kappa$ .

Why? Toward a contradiction, assume  $\mathbf{c}$  is a colouring of  $G$  with set of colours  $\subseteq \kappa$ . For each  $\gamma < \lambda$  let  $u_\gamma = \{\mathbf{c}((\alpha, \beta)) : \gamma < \alpha < \beta < \lambda \text{ and } (\alpha, \beta) \in Y\}$ . So  $\langle u_\gamma : \gamma < \lambda \rangle$  is  $\subseteq$ -decreasing sequence of subsets of  $\kappa$  and  $\kappa < \lambda = \text{cf}(\lambda)$ , hence for some  $\gamma(*) < \lambda$  and  $u_* \subseteq \kappa$  we have  $\gamma \in (\gamma(*), \lambda) \Rightarrow u_\gamma = u_*$ .

Hence  $E = \{\delta < \lambda : \delta \text{ is a limit ordinal } > \gamma(*) \text{ and } (\forall \alpha < \delta)(\mathbf{i}(\alpha) < \delta) \text{ and for every } \gamma < \delta \text{ and } i \in u_* \text{ there are } \alpha < \beta \text{ from } (\gamma, \delta) \text{ such that } (\alpha, \beta) \in Y \text{ and } \mathbf{c}((\alpha, \beta)) = i\}$  is a club of  $\lambda$ .

Now recall that  $\bar{C}$  guesses clubs hence for some  $\delta \in S$  we have  $C_\delta \subseteq E$ , so for every  $\varepsilon < \kappa$  we can choose  $\beta_{2\varepsilon} < \beta_{2\varepsilon+1}$  from  $(\alpha_{\delta, \varepsilon}^*, \alpha_{\delta, \varepsilon+1}^*)$  such that  $(\beta_{2\varepsilon}, \beta_{2\varepsilon+1}) \in Y$  and  $\varepsilon \in u_* \Rightarrow \mathbf{c}((\beta_{2\varepsilon}, \beta_{2\varepsilon+1})) = \varepsilon$ . So  $\langle \beta_\varepsilon : \varepsilon < \kappa \rangle$  is well defined, increasing and belongs to  $\Gamma_\delta$ , hence  $\bar{\beta}_\gamma = \langle \beta_\varepsilon : \varepsilon < \kappa \rangle$  for some  $\gamma \in X_\delta$ , hence  $(\alpha_{\delta, 0}^*, \gamma)$  belongs to  $Y$  and is connected in the graph to  $(\beta_{2\varepsilon}, \beta_{2\varepsilon+1})$  for  $\varepsilon < \kappa$ . Now if  $\varepsilon \in u_*$  then  $\mathbf{c}((\beta_{2\varepsilon}, \beta_{2\varepsilon+1})) = \varepsilon$  hence  $\mathbf{c}((\alpha_{\delta, 0}^*, \gamma)) \neq \varepsilon$  for every  $\varepsilon \in u_*$ , so  $\mathbf{c}((\alpha_{\delta, 0}^*, \gamma)) \in \kappa \setminus u_*$ . But  $u_* = u_{\alpha_{\delta, 0}^*}$  and  $\mathbf{c}((\alpha_{\delta, 0}^*, \gamma)) \in \kappa \setminus u_*$ , so we get contradiction to the definition of  $u_{\alpha_{\delta, 0}^*}$ . □<sub>1.1</sub>

Similarly

**Claim 1.2.** *There is an increasing continuous sequence  $\langle G_i : i \leq \lambda \rangle$  of graphs each of cardinality  $\lambda^\kappa$  such that  $\text{ch}(G_\lambda) > \kappa$  and  $i < \lambda \Rightarrow \text{ch}(G_i) \leq \kappa$  when:*

- ⊕ (a)  $\lambda = \text{cf}(\lambda)$
- (b)  $S \subseteq \{\delta < \lambda : \text{cf}(\delta) = \kappa\}$  is stationary not reflecting.

*Proof.* Like 1.1 but the  $X_i$  are not necessarily  $\subseteq \lambda$  or use 2.2. □<sub>1.2</sub>

## § 2. FROM ALMOST FREE

**Definition 2.1.** Suppose  $\eta_\beta \in {}^\kappa \text{Ord}$  for every  $\beta < \alpha(*)$  and  $u \subseteq \alpha(*)$ , and  $\alpha < \beta < \alpha(*) \Rightarrow \eta_\alpha \neq \eta_\beta$ .

1) We say  $\{\eta_\alpha : \alpha \in u\}$  is free when there exists a function  $h : u \rightarrow \kappa$  such that  $\langle \{\eta_\alpha(\varepsilon) : \varepsilon \in [h(\alpha), \kappa)\} : \alpha \in u \rangle$  is a sequence of pairwise disjoint sets.

2) We say  $\{\eta_\alpha : \alpha \in u\}$  is weakly free when there exists a sequence  $\langle u_{\varepsilon, \zeta} : \varepsilon < \kappa \rangle$  of subsets of  $u$  with union  $u$ , such that the function  $\eta_\zeta \mapsto \eta_\zeta(\varepsilon)$  is a one-to-one function on  $u_{\varepsilon, \zeta}$ , for each  $\varepsilon < \kappa$ .

**Claim 2.2.** 1) We have  $\text{INC}_{\text{chr}}(\mu, \lambda, \kappa)$  and even  $\text{INC}_{\text{chr}}^+(\mu, \lambda, \kappa)$ , see Definition 0.3(1) when:

- ⊞ (a)  $\alpha(*) \in [\mu, \mu^+)$  and  $\lambda$  is regular  $\leq \mu$  and  $\mu = \mu^\kappa$
- (b)  $\bar{\eta} = \langle \eta_\alpha : \alpha < \alpha(*) \rangle$
- (c)  $\eta_\alpha \in {}^\kappa \mu$
- (d)  $\langle u_i : i \leq \lambda \rangle$  is a  $\subseteq$ -increasing continuous sequence of subsets of  $\alpha(*)$  with  $u_\lambda = \alpha(*)$
- (e)  $\bar{\eta} \upharpoonright u_\alpha$  is free iff  $\alpha < \lambda$  iff  $\bar{\eta} \upharpoonright u_\alpha$  is weakly free.

2) We have  $\text{INC}_{\text{chr}}[\mu, \lambda, \kappa]$  and even  $\text{INC}_{\text{chr}}^+[\mu, \lambda, \kappa]$ , see Definition 0.3(4) when:

- ⊞<sub>2</sub> (a), (b), (c) as in ⊞ from 2.2
- (d)  $\bar{\eta}$  is not free
- (e)  $\bar{\eta} \upharpoonright u$  is free when  $u \in [\alpha(*)]^{< \lambda}$ .

*Proof.* For  $\mathcal{A} \subseteq {}^\kappa \text{Ord}$ , we define  $\tau_{\mathcal{A}}$  as the vocabulary  $\{P_\eta : \eta \in \mathcal{A}\} \cup \{F_\varepsilon : \varepsilon < \kappa\}$  where  $P_\eta$  is a unary predicate,  $F_\varepsilon$  a unary function (may be partial).

For part (1) without loss of generality for each  $i < \lambda$ ,  $u_i$  is an initial segment of  $\alpha(*)$  and let  $\mathcal{A} = \{\eta_\alpha : \alpha < \alpha(*)\}$ .

For part (2) let  $\langle \eta_\alpha : \alpha < \alpha(*) \rangle$  list  $\mathcal{A}$  and in both cases let  $<_{\mathcal{A}}$  be the well ordering  $\{(\eta_\alpha, \eta_\beta) : \alpha < \beta < \alpha(*)\}$  of  $\mathcal{A}$ .

We further let  $K_{\mathcal{A}}$  be the class of structures  $M$  such that (pedantically,  $K_{\mathcal{A}}$  depend also on the sequence  $\langle \eta_\alpha : \alpha < \alpha(*) \rangle$ ):

- ⊞<sub>1</sub> (a)  $M = (|M|, F_\varepsilon^M, P_\eta^M)_{\varepsilon < \kappa, \eta \in \mathcal{A}}$
- (b)  $\langle P_\eta^M : \eta \in \mathcal{A} \rangle$  is a partition of  $|M|$ , so for  $a \in M$  let  $\eta_a = \eta_a^M$  be the unique  $\eta \in \mathcal{A}$  such that  $a \in P_\eta^M$
- (c) if  $a_\ell \in P_{\eta_\ell}^M$  for  $\ell = 1, 2$  and  $F_\varepsilon^M(a_2) = a_1$  then  $\eta_1(\varepsilon) = \eta_2(\varepsilon)$  and  $\eta_1 <_{\mathcal{A}} \eta_2$ .

Let  $K_{\mathcal{A}}^*$  be the class of  $M$  such that

- ⊞<sub>2</sub> (a)  $M \in K_{\mathcal{A}}$
- (b)  $\|M\| = \lambda^\kappa$
- (c) if  $\eta \in \mathcal{A}$ ,  $u \subseteq \kappa$  and  $\eta_\varepsilon <_{\mathcal{A}} \eta$ ,  $\eta_\varepsilon(\varepsilon) = \eta(\varepsilon)$  and  $a_\varepsilon \in P_{\eta_\varepsilon}^M$  for  $\varepsilon \in u$  then for some  $a \in P_\eta^M$  we have  $\varepsilon \in u \Rightarrow F_\varepsilon^M(a) = a_\varepsilon$  and  $\varepsilon \in \kappa \setminus u \Rightarrow F_\varepsilon^M(a)$  not defined.

Clearly

$\boxplus_3$  there is  $M \in K_{\mathcal{A}}^*$

$\boxplus_4$  for  $M \in K_{\mathcal{A}}$  let  $G_M$  be the graph with:

- set of nodes  $|M|$
- set of edges  $\{\{a, F_\varepsilon^M(a)\} : a \in |M|, \varepsilon < \kappa \text{ when } F_\varepsilon^M(a) \text{ is defined}\}$ .

Now

$\boxplus_5$  if  $u \subseteq \alpha^*$ ,  $\mathcal{B} = \{\eta_\alpha : \alpha \in u\} \subseteq \mathcal{A}$  and  $\bar{\eta} \upharpoonright u$  is free, and  $M \in K_{\mathcal{A}}$  then  $G_{M, \mathcal{B}} := G_M \upharpoonright (\cup\{P_\eta^M : \eta \in \mathcal{B}_u\})$  has chromatic number  $\leq \kappa$ .

[Why? Let  $h : u \rightarrow \kappa$  witness that  $\bar{\eta} \upharpoonright u$  is free and for  $\varepsilon < \kappa$  let  $\mathcal{B}_\varepsilon := \{\eta_\alpha : \alpha \in u \text{ and } h(\alpha) = \varepsilon\}$ , so  $u = \cup\{u_\varepsilon : \varepsilon < \kappa\}$ , hence it is enough to prove for each  $\varepsilon < \kappa$  that  $G_{\mu, \mathcal{B}_\varepsilon}$  has chromatic number  $\leq \kappa$ . To prove this by induction on  $\alpha \leq \alpha^*$  we choose  $\mathbf{c}_\alpha^\varepsilon$  such that:

- $\boxplus_{5.1}$  (a)  $\mathbf{c}_\alpha^\varepsilon$  is a function  
 (b)  $\langle \mathbf{c}_\beta : \beta \leq \alpha \rangle$  is increasing continuous  
 (c)  $\text{Dom}(\mathbf{c}_\alpha^\varepsilon) = B_\alpha^\varepsilon := \cup\{P_{\eta_\beta}^M : \beta < \alpha \text{ and } \eta_\beta \in \mathcal{B}_\varepsilon\}$   
 (d)  $\text{Rang}(\mathbf{c}_\alpha^\varepsilon) \subseteq \kappa$   
 (e) if  $a, b \in \text{Dom}(\mathbf{c}_\alpha)$  and  $\{a, b\} \in \text{edge}(G_M)$  then  $\mathbf{c}_\alpha(a) \neq \mathbf{c}_\alpha(b)$ .

Clearly this suffices. Why is this possible?

If  $\alpha = 0$  let  $\mathbf{c}_\alpha^\varepsilon$  be empty, if  $\alpha$  is a limit ordinal let  $\mathbf{c}_\alpha^\varepsilon = \cup\{\mathbf{c}_\beta^\varepsilon : \beta < \alpha\}$  and if  $\alpha = \beta + 1 \wedge \alpha(\beta) \neq G$  let  $\mathbf{c}_\alpha = \mathbf{c}_\beta$ .

Lastly, if  $\alpha = \beta + 1 \wedge h(\beta) = \varepsilon$  we define  $\mathbf{c}_\alpha^\varepsilon$  as follows for  $a \in \text{Dom}(\mathbf{c}_\alpha^\varepsilon)$ ,  $\mathbf{c}_\alpha^\varepsilon(a)$  is:

Case 1:  $a \in B_\beta^\varepsilon$ .

Then  $\mathbf{c}_\alpha^\varepsilon(a) = \mathbf{c}_\beta^\varepsilon(a)$ .

Case 2:  $a \in B_\alpha^\varepsilon \setminus B_\beta^\varepsilon$ .

Then  $\mathbf{c}_\alpha^\varepsilon(a) = \min(\kappa \setminus \{c_\beta^\varepsilon(F_\zeta^M(a)) : \zeta < \varepsilon \text{ and } F_\zeta^M(a) \in \text{Dom}(\mathbf{c}_\beta^\varepsilon)\})$ .

This is O.K. as:

- $\boxplus_{5.2}$  (a)  $B_\alpha^\varepsilon = B_\beta^\varepsilon \cup P_{\eta_\beta}^M$   
 (b) if  $a \in B_\beta^\varepsilon$  then  $\mathbf{c}_\beta^\varepsilon(a)$  is well defined (so case 1 is O.K.)  
 (c) if  $\{a, b\} \in \text{edge}(G_M)$ ,  $a \in P_{\eta_\beta}^M$  and  $b \in B_\alpha^\varepsilon$  then  $b \in B_\beta^\varepsilon$  and  $b \in \{F_\zeta^M(a) : \zeta < \varepsilon\}$   
 (d)  $\mathbf{c}_\alpha^\varepsilon(a)$  is well defined in Case 2, too  
 (e)  $\mathbf{c}_\alpha^\varepsilon$  is a function from  $B_\alpha^\varepsilon$  to  $\kappa$   
 (f)  $\mathbf{c}_\alpha^\varepsilon$  is a colouring.

[Why? Clause (a) by  $\boxplus_{5.1}(b)$ , clause (b) by the induction hypothesis and clause (c) by  $\boxplus_2(c) + \boxplus_4$ . Next, clause (d) holds as  $\{c_\beta^\varepsilon(F_\zeta^M(a)) : \zeta < \varepsilon \text{ and } F_\zeta^M(a) \in \text{Dom}(\mathbf{c}_\beta^\varepsilon)\}$  is a set of cardinality  $\leq |\varepsilon| < \kappa$ . Clause (e) holds by the choices of the  $\mathbf{c}_\alpha^\varepsilon(a)$ 's. Lastly, clause (f) holds by the induction hypothesis of  $\mathbf{c}_\beta^\varepsilon$ , clause (c) and the choice of  $\mathbf{c}_\alpha^\varepsilon$ .]

So indeed  $\boxplus_5$  holds.]

$\boxplus_6$   $\text{chr}(G_M) > \kappa$  if  $M \in K_{\mathcal{A}}^*$ .

Why? Toward contradiction assume  $\mathbf{c} : G_M \rightarrow \kappa$  is a colouring. For each  $\eta \in \mathcal{A}$  and  $\varepsilon < \kappa$  let  $\Lambda_{\eta,\varepsilon} = \{\nu : \nu \in \mathcal{A}, \nu <_{\mathcal{A}} \eta, \nu(\varepsilon) = \eta(\varepsilon) \text{ and for some } a \in P_\nu^M \text{ we have } \mathbf{c}(a) = \varepsilon\}$ .

Let  $\mathcal{B}_\varepsilon = \{\eta \in \mathcal{A} : |\Lambda_{\eta,\varepsilon}| < \kappa\}$ . Now if  $\mathcal{A} \neq \cup\{\mathcal{B}_\varepsilon : \varepsilon < \kappa\}$  then pick any  $\eta \in \mathcal{A} \setminus \cup\{\mathcal{B}_\varepsilon : \varepsilon < \kappa\}$  and by induction on  $\varepsilon < \kappa$  choose  $\nu_\varepsilon \in \Lambda_{\eta,\varepsilon} \setminus \{\nu_\zeta : \zeta < \varepsilon\}$ , possible as  $\eta \notin \mathcal{B}_\varepsilon$  by the definition of  $\mathcal{B}_\varepsilon$ . By the definition of  $\Lambda_{\eta,\varepsilon}$  there is  $a_\varepsilon \in P_{\nu_\varepsilon}^M$  such that  $\mathbf{c}(\nu_\varepsilon) = \varepsilon$ . So as  $M \in K_{\mathcal{A}}^*$  there is  $a \in P_\eta^M$  such that  $\varepsilon < \kappa \Rightarrow F_\varepsilon^M(a) = a_\varepsilon$ , but  $\{a, a_\varepsilon\} \in \text{edge}(G_M)$  hence  $\mathbf{c}(a) \neq \mathbf{c}(a_\varepsilon) = \varepsilon$  for every  $\varepsilon < \kappa$ , contradiction. So  $\mathcal{A} = \cup\{\mathcal{B}_\varepsilon : \varepsilon < \kappa\}$ .

For each  $\varepsilon < \kappa$  we choose  $\zeta_\eta < \kappa$  for  $\eta \in \mathcal{B}_\varepsilon$  by induction on  $<_{\mathcal{A}}$  such that  $\zeta_\eta \notin \{\zeta_\nu : \nu \in \Lambda_{\eta,\varepsilon} \cap \mathcal{B}_\varepsilon\}$ . Let  $\mathcal{B}_{\varepsilon,\zeta} = \{\eta \in \mathcal{B}_\varepsilon : \zeta_\eta = \zeta\}$  for  $\varepsilon, \zeta < \kappa$  so  $\mathcal{A} = \cup\{\mathcal{B}_{\varepsilon,\zeta} : \varepsilon, \zeta < \kappa\}$  and clearly  $\eta \mapsto \eta(\varepsilon)$  is a one-to-one function with domain  $\mathcal{B}_{\varepsilon,\zeta}$ , contradiction to “ $\bar{\eta} = \bar{\eta} \upharpoonright u_\lambda$  is not weakly free”.  $\square_{2.2}$

**Observation 2.3.** 1) If  $\mathcal{A} \subseteq {}^\kappa \mu$  and  $\eta \neq \nu \in \mathcal{A} \Rightarrow (\forall^\infty \varepsilon < \kappa)(\eta(\varepsilon) \neq \nu(\varepsilon))$  then  $\mathcal{A}$  is free iff  $\mathcal{A}$  is weakly free.

2) The assumptions of 2.2(2) hold when:  $\mu \geq \lambda > \kappa$  are regular,  $S \subseteq S_\kappa^\mu$  stationary,  $\bar{\eta} = \langle \eta_\delta : \delta \in S \rangle$ ,  $\eta_\delta$  an increasing sequence of ordinals of length  $\kappa$  with limit  $\delta$  such that  $u \subseteq [\lambda]^{<\lambda} \Rightarrow \langle \text{Rang}(\eta_\delta) : \eta \in u \rangle$  has a one-to-one choice function.

**Conclusion 2.4.** Assume for every graph  $G$ , if  $H \subseteq G \wedge |H| < \lambda \Rightarrow \text{chr}(H) \leq \kappa$  then  $\text{chr}(G) \leq \kappa$ .

Then:

- (A) if  $\mu > \kappa = \text{cf}(\mu)$  and  $\mu \geq \lambda$  then  $\text{pp}(\mu) = \mu^+$
- (B) if  $\mu > \text{cf}(\mu) \geq \kappa$  and  $\mu \geq \lambda$  then  $\text{pp}(\mu) = \mu^+$ , i.e. the strong hypothesis
- (C) if  $\kappa = \aleph_0$  then above  $\lambda$  the SCH holds.

*Proof.* Clause (A): By 2.2 and [Sh:g, Ch.II], [Sh:g, Ch.IX,§1].

Clause (B): Follows from (A) by [Sh:g, Ch.VIII,§1].

Clause (C): Follows from (B) by [Sh:g, Ch.IX,§1].  $\square_{2.4}$

**Discussion 2.5.** Do we have  $\text{IC}_{\text{chr}}(\lambda, \aleph_\omega, \aleph_0)$  for some  $\lambda$ ? Assume not.

- 1) If  $\mu \in \mathbf{C}_{\aleph_0}$ , then necessarily  $\text{pp}(\mu) = \mu^+$  hence  $2^\mu = \mu^+$ .
- 2) Let  $\mu = \sum_n \lambda_n$ ,  $\lambda_n = \text{cf}(\lambda_n) < \lambda_{n+1}$ ,  $\bar{f} = \langle f_\alpha : \alpha < \mu^+ \rangle$  witness  $\mu^+ = \text{tcf}(\prod \lambda_n, <_{J_{\aleph_0}^{\text{bd}}})$ .

If  $S_* = \text{bad}_{\leq \aleph_\omega}(\bar{f}) = \text{bad}(\bar{f} \cap S_{< \aleph_\omega}^{\mu^+})$  non-stationary, we are done. Otherwise for some  $\ell \in \{1, 2, 3\}$  there is  $S \subseteq S_{\aleph_\ell}^{\mu^+}$  stationary not reflecting in any  $\delta \in S_{< \aleph_\omega}^{\mu^+}$ , so we have  $\diamond_S$ .

3) Recall for every  $n$ ,  $\text{IC}_{\text{chr}}(\beth_n^+(\kappa), \beth_n^+(\kappa), \kappa)$ ; so we can assume  $\bigwedge_n \beth_n < \aleph_\omega$ , so

$\beth_\omega = \aleph_\omega$ .

4) Still we can prove: for some  $\ell \in \{0, 1, 2, 3\}$ ,  $(\forall \lambda), \text{IC}_{\text{chr}}(\lambda, \aleph_\omega, \aleph_\ell)$ . Can we use  $X \subseteq [\mu]^n$ . Use [Sh:620]?

5) Can we use an example of part (3) on some  $\kappa = \text{cf}(\kappa) < \beth_u$  and use...

6) Can we take stationary  $S \subseteq S_\theta^\lambda$  not reflecting  $S \notin I_\theta[\lambda]$ , but on a partial square of it.

## REFERENCES

- [EH74] Paul Erdős and Andras Hajnal, *Solved and Unsolved Problems in set theory*, Proc. of the Symp. in honor of Tarksi's seventieth birthday in Berkeley 1971 (Leon Henkin, ed.), Proc. Symp in Pure Math., vol. XXV, 1974, pp. 269–287.
- [Sh:g] Saharon Shelah, *Cardinal Arithmetic*, Oxford Logic Guides, vol. 29, Oxford University Press, 1994.
- [Sh:309] ———, *Black Boxes*, , 0812.0656. 0812.0656. 0812.0656.
- [Sh:347] ———, *Incompactness for chromatic numbers of graphs*, A tribute to Paul Erdős, Cambridge Univ. Press, Cambridge, 1990, pp. 361–371.
- [Sh:620] ———, *Special Subsets of  ${}^{cf(\mu)}\mu$* , *Boolean Algebras and Maharam measure Algebras*, Topology and its Applications **99** (1999), 135–235, 8th Prague Topological Symposium on General Topology and its Relations to Modern Analysis and Algebra, Part II (1996). math.LO/9804156.

EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM, 91904, ISRAEL, AND, DEPARTMENT OF MATHEMATICS, HILL CENTER - BUSCH CAMPUS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019 USA

*E-mail address:* [shelah@math.huji.ac.il](mailto:shelah@math.huji.ac.il)

*URL:* <http://shelah.logic.at>