

Structured Preconditioners for Nonsingular Matrices of Block Two-by-Two Structures *

Zhong-Zhi Bai

*State Key Laboratory of Scientific/Engineering Computing
Institute of Computational Mathematics and Scientific/Engineering Computing
Academy of Mathematics and System Sciences
Chinese Academy of Sciences, P.O. Box 2719, Beijing 100080, P.R.China
Email: bzz@lsec.cc.ac.cn*

Abstract

For the large sparse block two-by-two real nonsingular matrices, we establish a general framework of practical and efficient structured preconditioners through matrix transformation and matrix approximations. For the specific versions such as modified block Jacobi-type, modified block Gauss-Seidel-type, and modified block unsymmetric (symmetric) Gauss-Seidel-type preconditioners, we precisely describe their concrete expressions and deliberately analyze eigenvalue distributions and positive definiteness of the preconditioned matrices. Also, we show that when these structured preconditioners are employed to precondition the Krylov subspace methods such as GMRES and restarted GMRES, fast and effective iteration solvers can be obtained for the large sparse systems of linear equations with block two-by-two coefficient matrices. In particular, these structured preconditioners can lead to efficient and high-quality preconditioning matrices for some typical matrices from the real-world applications.

Keywords: Block Two-by-Two Matrix, Preconditioner, Modified Block Relaxation Iteration, Eigenvalue Distribution, Positive Definiteness.

AMS(MOS) Subject Classifications: 65F10, 65F50; CR: G1.3.

1 Introduction

Let \mathbb{R}^n represent the real n -dimensional vector space, and $\mathbb{R}^{n \times n}$ the real $n \times n$ matrix space. Consider iterative solution of the large sparse system of linear equations

$$Ax = b, \quad A \in \mathbb{R}^{n \times n} \text{ nonsingular} \quad \text{and} \quad x, b \in \mathbb{R}^n. \quad (1.1)$$

In this paper, we will study algorithmic constructions and theoretical properties of practical and efficient structured preconditioners to the matrix $A \in \mathbb{R}^{n \times n}$ which is of the block two-by-two

*Subsidized by The Special Funds For Major State Basic Research Projects (No. G1999032803) and The National Natural Science Foundation (No. 10471146), P.R. China

structure

$$A = \begin{bmatrix} B & E \\ F & C \end{bmatrix}, \quad (1.2)$$

where $B \in \mathbb{R}^{p \times p}$ nonsingular, $C \in \mathbb{R}^{q \times q}$, $E \in \mathbb{R}^{p \times q}$ and $F \in \mathbb{R}^{q \times p}$, with $p \geq q$, such that $A \in \mathbb{R}^{n \times n}$ is nonsingular. Evidently, when the matrix block B is nonsingular, the matrix A is nonsingular if and only if its Schur complement $S_A = C - FB^{-1}E$ is nonsingular.

Linear systems of the form (1.1)-(1.2) arise in a variety of scientific and engineering applications, including computational fluid dynamics[21, 23, 26], mixed finite element approximation of elliptic partial differential equations [16, 38], optimization[25, 30, 34], optimal control[13], weighted and equality constrained least squares estimation[14], stationary semiconductor device[36, 42, 43], structural analysis[44], electrical networks[44], inversion of geophysical data[31], and so on.

As we have known, preconditioned Krylov subspace methods[40] are efficient iterative solvers for the system of linear equations (1.1)-(1.2), and effective and high-quality preconditioners play a crucial role to guarantee their fast convergence and economical costs. A number of structured preconditioners have been studied in the literature for some special cases of the block two-by-two matrix A in (1.2). Besides specialized incomplete factorization preconditioners[17, 18] we mention, among others, algebraic multilevel iteration preconditioners[2, 3, 4, 5, 12], block and approximate Schur complement preconditioners[21, 23], splitting iteration preconditioners[15, 19, 22, 28, 29, 39, 45], block definite and indefinite preconditioners [24, 34, 38, 10], and block triangular preconditioners[35, 37, 10]. Theoretical analyses and experimental results have shown that these preconditioners may lead to nicely clustered eigenvalue distributions of the preconditioned matrices and, hence, result in fast convergence of the preconditioned Krylov subspace iteration methods for solving the large sparse system of linear equations (1.1)-(1.2). However, exact inversions of the matrix block B or C , as well as the Schur complement S_A , are demanded for most of these preconditioners, which makes them less practical and effective in actual applications.

In this paper, by sufficiently utilizing the matrix structure and property, we first establish a general framework of a class of practical and efficient structured preconditioners to the matrix $A \in \mathbb{R}^{n \times n}$ in (1.2) through matrix transformation and several steps of matrix approximations; these preconditioners can avoid the exact inversions of the matrix blocks B and C , as well as the Schur complement S_A , and cover the known preconditioners mentioned previously as special cases. Then, with this framework we further present a family of practical and efficient preconditioners by technically combining it with the modified block relaxation iterations[6, 7], which includes the modified block Jacobi-type, the modified block Gauss-Seidel-type and the modified block unsymmetric (symmetric) Gauss-Seidel-type preconditioners as typical examples. Moreover, we particularly discuss the eigenvalue distributions and the positive definiteness of the preconditioned matrices with respect to the modified block Jacobi-, the modified block Gauss-Seidel-, and the modified block unsymmetric (symmetric) Gauss-Seidel-type preconditioners to the block two-by-two matrix A , and deliberately address the applications of these preconditioners to three classes of real-world matrices, i.e., the symmetric positive definite matrix, the saddle point matrix and the Hamiltonian matrix. Besides, we show that when these structured preconditioners are employed to precondition the Krylov subspace methods such as GMRES or restarted GMRES, fast and effective iteration solvers can be obtained for the large sparse system

of linear equations (1.1)-(1.2).

The organization of this paper is as follows. After establishing the general framework of the structured preconditioners in Section 2, we present the modified block splitting iteration preconditioners and study the eigenvalue distributions and the positive definiteness of the corresponding preconditioned matrices in Section 3; connections of these preconditioners to Krylov subspace iteration methods are also briefly discussed in this section. Specifications of these preconditioners to three classes of real-world matrices are investigated in Section 4. Finally, in Section 5, we use a brief conclusion and several remarks to end the paper.

2 General Framework of the Structured Preconditioners

The construction of our structured preconditioners basically includes the following three steps: Firstly, seek two nonsingular block two-by-two matrices $P, Q \in \mathbb{R}^{n \times n}$ such that P and Q are easily invertible and $A = PHQ$ holds for a block two-by-two matrix $H \in \mathbb{R}^{n \times n}$ of certain good properties; Secondly, approximate the matrix H by another block two-by-two matrix $\overline{W} \in \mathbb{R}^{n \times n}$ by dropping some higher-order small block quantities; And thirdly, approximate the matrix \overline{W} further by another block two-by-two matrix $W \in \mathbb{R}^{n \times n}$ that is also easily invertible. Then, the resulting preconditioners are of the form $M = PWQ$. See [9, 11].

Let $L_B, R_B \in \mathbb{R}^{p \times p}$ and $L_C, R_C \in \mathbb{R}^{q \times q}$ be nonsingular matrices such that

$$L_B^{-1} B R_B^{-1} = J_B \quad \text{and} \quad L_C^{-1} C R_C^{-1} = J_C, \quad (2.1)$$

or equivalently,

$$B = L_B J_B R_B \quad \text{and} \quad C = L_C J_C R_C, \quad (2.2)$$

where $J_B \in \mathbb{R}^{p \times p}$ is a matrix approximating to the identity matrix $I_B \in \mathbb{R}^{p \times p}$, and $J_C \in \mathbb{R}^{q \times q}$ is a matrix approximating to the identity matrix $I_C \in \mathbb{R}^{q \times q}$ when it is positive definite and approximating to $-I_C \in \mathbb{R}^{q \times q}$ when it is negative definite. For simplicity, in the sequel we will abbreviate the identity matrices I_B and I_C as I , with its dimension being inferred from the context.

Evidently, L_B, R_B and L_C, R_C can be considered as *split preconditioners* to the matrix blocks B and C , respectively, whose preconditioning properties can be measured by the approximation degrees of the matrices J_B and $\pm J_C$ to the identity matrix I . There are many possible choices of the matrices L_B, R_B and L_C, R_C . For example, they may be the incomplete lower-upper triangular factors[2, 40], the incomplete orthogonal triangular factors[8], the approximate inverse preconditioners[40], the splitting iteration matrices[2, 6, 7, 27], the multigrid or the algebraic multilevel approximations[2, 3, 4, 5, 12], or even technical combinations of the abovementioned matrices, to the matrix blocks B and C , respectively.

In particular, when $C \in \mathbb{R}^{q \times q}$ is singular, besides the possible choices mentioned above, we may choose L_C and R_C according to the following cases:

- (i) if C is a symmetric positive semidefinite matrix, we may let $L_C = R_C = I$. Hence, $J_C = C$ is also symmetric positive semidefinite;
- (ii) if C is a symmetric negative semidefinite matrix, we may let $L_C = -I$ and $R_C = I$ (or $L_C = I$ and $R_C = -I$). Hence, $J_C = -C$ is symmetric positive semidefinite. Or we may let $L_C = R_C = I$. Hence, $J_C = C$ is also symmetric negative semidefinite;

(iii) if C is a general singular matrix, we may let $L_C = R_C = I$. Hence, $J_C = C$ is also singular.

To construct a high-quality structured preconditioner to the block two-by-two matrix $A \in \mathbb{R}^{n \times n}$, we introduce matrices

$$D_L = \begin{bmatrix} L_B & O \\ O & L_C \end{bmatrix}, \quad D_R = \begin{bmatrix} R_B & O \\ O & R_C \end{bmatrix}$$

and

$$\bar{E} = L_B^{-1} E R_C^{-1}, \quad \bar{F} = L_C^{-1} F R_B^{-1}, \quad (2.3)$$

where O denotes the zero matrix. Then from (2.2) we have

$$\begin{aligned} A &= \begin{bmatrix} B & E \\ F & C \end{bmatrix} = \begin{bmatrix} L_B J_B R_B & E \\ F & L_C J_C R_C \end{bmatrix} \\ &= \begin{bmatrix} L_B & O \\ O & L_C \end{bmatrix} \begin{bmatrix} J_B & L_B^{-1} E R_C^{-1} \\ L_C^{-1} F R_B^{-1} & J_C \end{bmatrix} \begin{bmatrix} R_B & O \\ O & R_C \end{bmatrix} \\ &:= D_L \bar{A} D_R, \end{aligned}$$

where

$$\bar{A} := \begin{bmatrix} J_B & \bar{E} \\ \bar{F} & J_C \end{bmatrix}.$$

Furthermore, we can find a unit lower triangular matrix $L \in \mathbb{R}^{n \times n}$ and a unit upper triangular matrix $U \in \mathbb{R}^{n \times n}$ of block two-by-two structures such that $H = L \bar{A} U$ is block-diagonally dominant as far as possible, and may also possess some other desired good properties.

In fact, if we let

$$L = \begin{bmatrix} I & O \\ L_{21} & I \end{bmatrix} \quad \text{and} \quad U = \begin{bmatrix} I & U_{12} \\ O & I \end{bmatrix},$$

then by concrete computations we obtain

$$\begin{aligned} H &= \begin{bmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{bmatrix} := L \bar{A} U \\ &= \begin{bmatrix} I & O \\ L_{21} & I \end{bmatrix} \begin{bmatrix} J_B & \bar{E} \\ \bar{F} & J_C \end{bmatrix} \begin{bmatrix} I & U_{12} \\ O & I \end{bmatrix} \end{aligned}$$

with

$$\begin{cases} H_{11} = J_B, & H_{12} = J_B U_{12} + \bar{E}, \\ H_{21} = L_{21} J_B + \bar{F}, & H_{22} = J_C + L_{21} J_B U_{12} + L_{21} \bar{E} + \bar{F} U_{12}, \end{cases}$$

and

$$A = D_L \bar{A} D_R = D_L (L^{-1} H U^{-1}) D_R = (D_L L^{-1}) H (U^{-1} D_R) := P H Q$$

with

$$P := D_L L^{-1} = \begin{bmatrix} L_B & O \\ O & L_C \end{bmatrix} \begin{bmatrix} I & O \\ -L_{21} & I \end{bmatrix} = \begin{bmatrix} L_B & O \\ -L_C L_{21} & L_C \end{bmatrix} \quad (2.4)$$

and

$$Q := U^{-1} D_R = \begin{bmatrix} I & -U_{12} \\ O & I \end{bmatrix} \begin{bmatrix} R_B & O \\ O & R_C \end{bmatrix} = \begin{bmatrix} R_B & -U_{12} R_C \\ O & R_C \end{bmatrix}. \quad (2.5)$$

We can now choose the matrices L and U such that either of the following two principles is satisfied as far as possible:

(P_1) the matrix H is block-diagonally dominant and symmetric;

(P_2) the matrix H is block-diagonally dominant and skew-symmetric.

Because if the matrix H satisfies either of the principles (P_1) and (P_2), we can easily construct a good approximation to it, and hence, obtain a high-quality preconditioner M to the original matrix A .

According to both (P_1) and (P_2), we can take L_{21} and U_{12} such that

$$\begin{cases} H_{21} = L_{21} J_B + \overline{F} \approx (J_B U_{12} + \overline{E})^T = \pm H_{12}^T, \\ H_{21} = L_{21} J_B + \overline{F} \approx O. \end{cases}$$

Recalling that $J_B \approx I$, we can let

$$L_{21} = -\overline{F} \quad \text{and} \quad U_{12} = -\overline{E}.$$

Thus, for both cases, it follows from (2.4) and (2.5) that the matrices P and Q have the following expressions:

$$P = \begin{bmatrix} L_B & O \\ L_C \overline{F} & L_C \end{bmatrix}, \quad Q = \begin{bmatrix} R_B & \overline{E} R_C \\ O & R_C \end{bmatrix}. \quad (2.6)$$

Therefore, for these choices of the matrices P and Q , we have

$$\begin{aligned} H &= \begin{bmatrix} J_B & (I - J_B) \overline{E} \\ \overline{F} (I - J_B) & J_C - \overline{F} \overline{E} - \overline{F} (I - J_B) \overline{E} \end{bmatrix} \\ &\approx \begin{bmatrix} J_B & (I - J_B) \overline{E} \\ \overline{F} (I - J_B) & J_C - \overline{F} \overline{E} \end{bmatrix} := \overline{W}. \end{aligned} \quad (2.7)$$

Because the nonsingularity of the matrix A implies that the matrix \overline{A} and its Schur complement $S_{\overline{A}} := J_C - \overline{F} J_B^{-1} \overline{E}$ are nonsingular, and

$$J_C - \overline{F} \overline{E} = S_{\overline{A}} + \overline{F} (I - J_B) J_B^{-1} \overline{E} \quad (2.8)$$

and the Schur complement of \overline{W} is

$$S_{\overline{W}} := J_C - \overline{F} \overline{E} - \overline{F} (I - J_B) J_B^{-1} (I - J_B) \overline{E} = S_{\overline{A}} - \overline{F} (I - J_B) J_B^{-1} \overline{E},$$

we immediately know that when

$$\|I - J_B\|_2 < \max \left\{ \frac{1}{1 + \|\overline{E}\|_2 \|S_A^{-1} \overline{F}\|_2}, \frac{1}{1 + \|\overline{E} S_A^{-1}\|_2 \|\overline{F}\|_2} \right\}, \quad (2.9)$$

both matrices $J_C - \overline{F}\overline{E}$ and \overline{W} are nonsingular.

Now, if we let $W \in \mathbb{R}^{n \times n}$ be a nonsingular “replacement” of the matrix \overline{W} , or in other words, a “replacement” to the matrix H , then the matrix

$$M = PWQ \quad (2.10)$$

is a natural preconditioner to the original matrix $A \in \mathbb{R}^{n \times n}$. And under the condition (2.9) this preconditioner is well defined.

Note that here we use the term “replacement” other than “approximation”. This is because sometimes we may choose the matrix W being not an approximation to \overline{W} in the usual sense so that the obtained preconditioner and the preconditioned matrix can possess some desired properties such as positive definiteness and, hence, a specified Krylov subspace iteration method may exploit its efficiency sufficiently.

If M is used as a left preconditioner to A , then

$$M^{-1}A = (PWQ)^{-1}(PHQ) = Q^{-1}(W^{-1}H)Q := Q^{-1}K_LQ \quad (2.11)$$

with

$$K_L = W^{-1}H. \quad (2.12)$$

Therefore, the preconditioning property of M to A is determined by the properties of the matrices K_L and Q . If M is used as a right preconditioner to A , then

$$AM^{-1} = (PHQ)(PWQ)^{-1} = P(HW^{-1})P^{-1} := PK_RP^{-1} \quad (2.13)$$

with

$$K_R = HW^{-1}. \quad (2.14)$$

Therefore, the preconditioning property of M to A is determined by the properties of the matrices K_R and P . In general, if the matrix W admits a split form

$$W = W_L W_R, \quad (2.15)$$

then (2.10) straightforwardly leads to a split preconditioner

$$M = (PW_L)(W_RQ) := M_L M_R, \quad \text{with } M_L = PW_L \quad \text{and} \quad M_R = W_RQ, \quad (2.16)$$

to the original matrix A . Because

$$M_L^{-1}AM_R^{-1} = (PW_L)^{-1}(PHQ)(W_RQ)^{-1} = W_L^{-1}HW_R^{-1} := K, \quad (2.17)$$

we see that the preconditioning property of M to A is determined by the property of the matrix K .

Evidently, the matrices K_L , K_R and K are similar, and hence, they have exactly the same spectrum. However, the eigenvectors of these kinds of preconditioned matrices are usually quite different, which may lead to different performance results of the corresponding preconditioned Krylov subspace iteration methods.

In actual applications, when the matrix M defined in (2.10) is employed as a preconditioner to some Krylov subspace iteration method for solving the block two-by-two system of linear equations (1.1), we need to solve a generalized residual equation of the form

$$Mz = r \quad (2.18)$$

at each iteration step, where r is the current residual vector. By making use of the two-by-two block structure of M , we can obtain the following practical procedure for computing the generalized residual vector $z = M^{-1}r$.

PROCEDURE FOR COMPUTING THE GENERALIZED RESIDUAL VECTOR. Let $r = (r_1^T, r_2^T)^T$, $z = (z_1^T, z_2^T)^T$ and $w = (w_1^T, w_2^T)^T$, with $r_1, z_1, w_1 \in \mathbb{R}^p$ and $r_2, z_2, w_2 \in \mathbb{R}^q$.

1. Solve $L_B t_1 = r_1$ and $L_C t_2 = r_2$ to get t_1 and t_2 , and let $t_2 := t_2 + \overline{F}t_1$;
2. Solve $Ww = t$ to get w , with $t = (t_1^T, t_2^T)^T$;
3. Solve $R_C z_2 = w_2$ and $R_B z_1 = w_1 - \overline{E}w_2$ to get z_1 and z_2 .

When the approximation matrix $W \in \mathbb{R}^{n \times n}$ to the matrix $\overline{W} \in \mathbb{R}^{n \times n}$ is specified, a concrete procedure for computing the generalized residual vector $z \in \mathbb{R}^n$ defined by (2.18) can be straightforwardly obtained from this procedure.

Usually, the matrix $W \in \mathbb{R}^{n \times n}$ may involve information about the matrices J_B , J_C , \overline{E} and \overline{F} . Therefore, to solve the linear system $Ww = t$ we may need to compute the vectors

$$\begin{cases} \overline{w}_1 &= J_B w_1 = L_B^{-1} B R_B^{-1} w_1, & \tilde{w}_1 &= \overline{F} w_1 = L_C^{-1} F R_B^{-1} w_1, \\ \overline{w}_2 &= J_C w_2 = L_C^{-1} C R_C^{-1} w_2, & \tilde{w}_2 &= \overline{E} w_2 = L_B^{-1} E R_C^{-1} w_2. \end{cases}$$

These vectors can be economically computed by the following formulas:

1. Solve $R_B t_1 = w_1$;
2. Solve $L_B \overline{w}_1 = B t_1$, $L_C \tilde{w}_1 = F t_1$;
3. Solve $R_C t_2 = w_2$;
4. Solve $L_C \overline{w}_2 = C t_2$, $L_B \tilde{w}_2 = E t_2$.

3 Several Practical Structured Preconditioners

In this section, we will construct three classes of structured approximations W to the block two-by-two matrix \overline{W} , or in other words, to the block two-by-two matrix H in (2.7), by making use of the modified block Jacobi, the modified block Gauss-Seidel and the modified block

unsymmetric Gauss-Seidel splittings of \overline{W} . See [6, 7] for details. Therefore, three types of structured preconditioners to the original block two-by-two matrix $A \in \mathbb{R}^{n \times n}$, called as the *modified block Jacobi-type (MBJ-type)* preconditioner, the *modified block Gauss-Seidel-type (MBGS-type)* preconditioner and the *modified block unsymmetric Gauss-Seidel-type (MBUGS-type)* preconditioner, can be obtained, correspondingly.

To analyze the spectral property of the preconditioned matrices with respect to the above-mentioned preconditioners, we need the following two basic facts.

Lemma 3.1 *Let $L \in \mathbb{R}^{(p+q) \times (p+q)}$ and $U \in \mathbb{R}^{(p+q) \times (p+q)}$ be unit lower and upper triangular matrices of the block two-by-two forms*

$$L = \begin{bmatrix} I & O \\ L_{21} & I \end{bmatrix} \quad \text{and} \quad U = \begin{bmatrix} I & U_{12} \\ O & I \end{bmatrix},$$

where $L_{21} \in \mathbb{R}^{q \times p}$ and $U_{12} \in \mathbb{R}^{p \times q}$. Let

$$\gamma(t) = \left[1 + \frac{1}{2}t \left(t + \sqrt{t^2 + 4} \right) \right]^{\frac{1}{2}} \quad (3.1)$$

be a monotone increasing function with respect to t in the interval $[0, +\infty)$. Then it holds that

$$\|L\|_2 = \gamma(\|L_{21}\|_2) \quad \text{and} \quad \|U\|_2 = \gamma(\|U_{12}\|_2).$$

Proof. By direct computations we have

$$L^T L = \begin{bmatrix} I & L_{21}^T \\ O & I \end{bmatrix} \begin{bmatrix} I & O \\ L_{21} & I \end{bmatrix} = \begin{bmatrix} I + L_{21}^T L_{21} & L_{21}^T \\ L_{21} & I \end{bmatrix}.$$

Without loss of generality, we assume $p \geq q$. From Theorem 2.5.2 in [27, page 70] we know that the matrix L_{21} admits a *singular value decomposition (SVD)*, i.e., there exist two orthogonal matrices $V_1 \in \mathbb{R}^{q \times q}$ and $V_2 \in \mathbb{R}^{p \times p}$ and a matrix $\tilde{\Sigma} = [\Sigma, O] \in \mathbb{R}^{q \times p}$, with $\Sigma = \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_q) \in \mathbb{R}^{q \times q}$ being a nonnegative diagonal matrix having the maximum diagonal entry $\sigma_1 = \|L_{21}\|_2$, such that $L_{21} = V_1^T \tilde{\Sigma} V_2$ holds. Define

$$V = \begin{bmatrix} V_2 & O \\ O & V_1 \end{bmatrix}.$$

Then V is an orthogonal matrix, too. It follows from concrete computations that

$$L^T L = V^T \begin{bmatrix} I + \Sigma^2 & O & \Sigma \\ O & I & O \\ \Sigma & O & I \end{bmatrix} V.$$

Therefore, detailed analysis shows that the eigenvalues of the matrix $L^T L$ are 1 with multiplicity $p - q$, and

$$1 + \frac{1}{2}\sigma_k \left(\sigma_k \pm \sqrt{\sigma_k^2 + 4} \right), \quad k = 1, 2, \dots, q.$$

It then follows straightforwardly that the spectral radius of the matrix $L^T L$, say $\rho(L^T L)$, is given by

$$\begin{aligned}\rho(L^T L) &= 1 + \frac{1}{2}\sigma_1 \left(\sigma_1 + \sqrt{\sigma_1^2 + 4} \right) \\ &= 1 + \frac{1}{2}\|L_{21}\|_2 \left(\|L_{21}\|_2 + \sqrt{\|L_{21}\|_2^2 + 4} \right),\end{aligned}$$

and therefore,

$$\|L\|_2 = \rho(L^T L)^{\frac{1}{2}} = \left[1 + \frac{1}{2}\|L_{21}\|_2 \left(\|L_{21}\|_2 + \sqrt{\|L_{21}\|_2^2 + 4} \right) \right]^{\frac{1}{2}} = \gamma(\|L_{21}\|_2).$$

The proof of the second equality can be demonstrated in a similar fashion. \square

We remark that for the real one-variable function $\gamma(t)$ defined by (3.1), the estimate $\gamma(t) \leq t+1$ holds for all $t \in [0, +\infty)$ because of $\sqrt{t^2+4} \leq t+2$ and $\sqrt{t^2+t+1} \leq t+1$.

Lemma 3.2 *Let $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathbb{C}^{n \times n}$ be a diagonal matrix, and $Y \in \mathbb{C}^{n \times n}$ a given matrix, where $\mathbb{C}^{n \times n}$ represents the complex $n \times n$ matrix space. If there exists a positive constant ρ_y such that $\|\Lambda - Y\|_2 \leq \rho_y$, then all eigenvalues of the matrix Y are located within $\cup_{i=1}^n \mathcal{N}(\lambda_i, \rho_y)$, where $\mathcal{N}(\lambda_i, \rho_y)$ denotes the circle having center λ_i and radius ρ_y on the complex plane.*

Proof. Let λ be an eigenvalue of the matrix $Y \in \mathbb{C}^{n \times n}$ and v be the corresponding normalized eigenvector. Then we have $(\Lambda - Y)v = (\Lambda - \lambda I)v$. Hence,

$$\|(\Lambda - \lambda I)v\|_2 = \|(\Lambda - Y)v\|_2 \leq \|\Lambda - Y\|_2 \leq \rho_y.$$

It then follows that $\|\Lambda - \lambda I\|_2 \leq \rho_y$. Therefore, it holds that $|\lambda - \lambda_i| \leq \rho_y$ ($i = 1, 2, \dots, n$), or equivalently, $\lambda \in \cup_{i=1}^n \mathcal{N}(\lambda_i, \rho_y)$. \square

For the simplicity of our statements, in the sequel we always use $\gamma : (0, +\infty) \rightarrow (0, +\infty)$ to represent the function defined by (3.1). For the matrices J_B in (2.1) and $\overline{E}, \overline{F}$ in (2.3), we write

$$\overline{\Delta}_1 = \overline{F}(I - J_B)\overline{E} \quad \text{and} \quad \overline{\Delta}_2 = \overline{F}(I - J_B)^2\overline{E},$$

and denote the (2, 2)-block entry of the matrix \overline{W} in (2.7) by \overline{S} , i.e.,

$$\overline{S} = J_C - \overline{F}\overline{E}. \tag{3.2}$$

Assume \overline{W} and \overline{S} be nonsingular, let S be a nonsingular matrix that is a *replacement* to \overline{S} (e.g., $S = \pm(I - \overline{F}\overline{E})$ or $S = \pm(I - \text{diag}(\overline{F}\overline{E}))$, etc.), and define the quantities

$$\Theta = \|\overline{E}\|_2, \quad \Gamma = \|\overline{F}\|_2, \quad \Theta_s = \|\overline{E}S^{-1}\|_2, \quad \Gamma_s = \|S^{-1}\overline{F}\|_2.$$

In addition, in the case that S is an approximation to \overline{S} , we define the quantities

$$\epsilon_L = \max\{\|I - J_B\|_2, \|I - S^{-1}\overline{S}\|_2\}, \quad \epsilon_R = \max\{\|I - J_B\|_2, \|I - \overline{S}S^{-1}\|_2\};$$

and in the case that S is an approximation to $-\overline{S}$, instead of ϵ_L and ϵ_R we use the quantities

$$\tilde{\epsilon}_L = \max\{\|I - J_B\|_2, \|I + S^{-1}\overline{S}\|_2\}, \quad \tilde{\epsilon}_R = \max\{\|I - J_B\|_2, \|I + \overline{S}S^{-1}\|_2\}.$$

For two positive constants $\rho_L^{(\xi)}$ and $\rho_R^{(\xi)}$ to be specified later, we use $\mathcal{N}^{(\xi)}$ to denote the circle having center $(1, 0)$ and radius $\rho^{(\xi)} := \min\{\rho_L^{(\xi)}\epsilon_L, \rho_R^{(\xi)}\epsilon_R\}$, and use $\widetilde{\mathcal{N}}^{(\xi)}$ to denote the union of the two circles having centers $(-1, 0)$ and $(1, 0)$ and radius $\widetilde{\rho}^{(\xi)} := \min\{\rho_L^{(\xi)}\widetilde{\epsilon}_L, \rho_R^{(\xi)}\widetilde{\epsilon}_R\}$, on the complex plane, respectively.

By making use of the above notations, the nonsingularity of the matrices \overline{S} and \overline{W} can be precisely described by the following lemma.

Lemma 3.3 *The matrices \overline{S} and \overline{W} are nonsingular, provided either of the following conditions holds:*

- (1) S is an approximation to \overline{S} , and
 - (a) $\epsilon_L < 1 + \Theta\Gamma_s - \sqrt{\Theta\Gamma_s(\Theta\Gamma_s + 2)}$, or
 - (b) $\epsilon_R < 1 + \Theta_s\Gamma - \sqrt{\Theta_s\Gamma(\Theta_s\Gamma + 2)}$;
- (2) S is an approximation to $-\overline{S}$, and
 - (a) $\widetilde{\epsilon}_L < 1 + \Theta\Gamma_s - \sqrt{\Theta\Gamma_s(\Theta\Gamma_s + 2)}$, or
 - (b) $\widetilde{\epsilon}_R < 1 + \Theta_s\Gamma - \sqrt{\Theta_s\Gamma(\Theta_s\Gamma + 2)}$.

Proof. We only prove (1a), as the other conclusions can be demonstrated analogously.

Because $\|I - J_B\|_2 \leq \epsilon_L < 1$, it holds that

$$\|J_B^{-1}\|_2 = \|[I - (I - J_B)]^{-1}\|_2 \leq \frac{1}{1 - \|I - J_B\|_2}.$$

From (2.8) we have

$$S_{\overline{A}} = \overline{S} - \overline{F}(I - J_B)J_B^{-1}\overline{E}.$$

Hence,

$$\begin{aligned} \|I - S^{-1}S_{\overline{A}}\|_2 &\leq \|I - S^{-1}\overline{S}\|_2 + \|\overline{E}\|_2\|S^{-1}\overline{F}\|_2 \cdot \frac{\|I - J_B\|_2}{1 - \|I - J_B\|_2} \\ &\leq \left(1 + \frac{\Theta\Gamma_s}{1 - \|I - J_B\|_2}\right) \cdot \max\{\|I - J_B\|_2, \|I - S^{-1}\overline{S}\|_2\} \\ &\leq \left(1 + \frac{\Theta\Gamma_s}{1 - \epsilon_L}\right) \epsilon_L \\ &< 1. \end{aligned}$$

It then follows that

$$\|S_{\overline{A}}^{-1}\overline{F}\|_2 \leq \frac{\|S^{-1}\overline{F}\|_2}{1 - \|I - S^{-1}S_{\overline{A}}\|_2} \leq \frac{\Gamma_s(1 - \epsilon_L)}{(1 - \epsilon_L)^2 - \Theta\Gamma_s\epsilon_L}.$$

Now, we easily see that (2.9) holds when

$$\epsilon_L < \frac{1}{1 + \frac{\Theta\Gamma_s(1 - \epsilon_L)}{(1 - \epsilon_L)^2 - \Theta\Gamma_s\epsilon_L}} = \frac{(1 - \epsilon_L)^2 - \Theta\Gamma_s\epsilon_L}{(1 - \epsilon_L)^2 - \Theta\Gamma_s\epsilon_L + \Theta\Gamma_s(1 - \epsilon_L)},$$

or equivalently, $\epsilon_L^2 - 2(1 + \Theta\Gamma_s)\epsilon_L + 1 > 0$. Therefore, when

$$\epsilon_L < 1 + \Theta\Gamma_s - \sqrt{\Theta\Gamma_s(\Theta\Gamma_s + 2)},$$

the matrices \overline{S} and \overline{W} are nonsingular. \square

We first consider the case that $S \approx \overline{S}$. The case that $S \approx -\overline{S}$ will be discussed in Section 3.4.

3.1 The MBJ-type Preconditioners

If the matrix $W \in \mathbb{R}^{n \times n}$ in (2.10) is taken to be the modified block Jacobi splitting matrix[6, 7] of the matrix \overline{W} in (2.7), i.e.,

$$W^{(J)} := W = \begin{bmatrix} I & O \\ O & S \end{bmatrix}, \quad (3.3)$$

then we obtain the *modified block Jacobi-type (MBJ-type)* preconditioner $M = PW^{(J)}Q$ to the original matrix $A \in \mathbb{R}^{n \times n}$. Note that when $S \in \mathbb{R}^{q \times q}$ is symmetric positive definite, $W^{(J)}$ is a symmetric positive definite matrix, and when $S \in \mathbb{R}^{q \times q}$ is symmetric negative definite, $W^{(J)}$ is a symmetric indefinite matrix.

The following theorem describes the eigenvalue distribution of the preconditioned matrix with respect to the MBJ-type preconditioner.

Theorem 3.1 *Let $M = PW^{(J)}Q$ be the MBJ-type preconditioner to the block two-by-two matrix $A = PHQ \in \mathbb{R}^{n \times n}$ in (1.2), where P and Q are given by (2.6), H is given by (2.7), and $W^{(J)}$ is defined by (3.3). Denote by $K_L = W^{(J)^{-1}}H$ and $K_R = HW^{(J)^{-1}}$, respectively. Then it holds that*

$$(i) \|I - K_L\|_2 \leq \rho_L^{(J)} \epsilon_L, \text{ with } \rho_L^{(J)} = \gamma(\Theta) \cdot \gamma(\Gamma_s); \text{ and}$$

$$(ii) \|I - K_R\|_2 \leq \rho_R^{(J)} \epsilon_R, \text{ with } \rho_R^{(J)} = \gamma(\Theta_s) \cdot \gamma(\Gamma).$$

It follows from Lemma 3.2 as well as (2.11) and (2.13) that the eigenvalues of the matrices $M^{-1}A$ and AM^{-1} are located within a circle having center $(1, 0)$ and radii $\rho_L^{(J)} \epsilon_L$ and $\rho_R^{(J)} \epsilon_R$, respectively, and therefore, they are all within the circle $\mathcal{N}^{(J)}$.

Proof. We only prove (i), as (ii) can be verified analogously.

From (2.7) and (3.3) we have

$$K_L = W^{(J)^{-1}}H = \begin{bmatrix} J_B & (I - J_B)\overline{E} \\ S^{-1}\overline{F}(I - J_B) & S^{-1}\overline{S} - S^{-1}\overline{\Delta}_1 \end{bmatrix}.$$

Hence,

$$I - K_L = \begin{bmatrix} I & O \\ -S^{-1}\overline{F} & I \end{bmatrix} \begin{bmatrix} I - J_B & O \\ O & I - S^{-1}\overline{S} \end{bmatrix} \begin{bmatrix} I & -\overline{E} \\ O & I \end{bmatrix}.$$

By making use of Lemma 3.1 we can immediately obtain

$$\begin{aligned}\|I - K_L\|_2 &\leq \gamma(\|S^{-1}\bar{F}\|_2) \cdot \gamma(\|\bar{E}\|_2) \cdot \max\{\|I - J_B\|_2, \|I - S^{-1}\bar{S}\|_2\} \\ &= \gamma(\Theta) \cdot \gamma(\Gamma_s) \cdot \max\{\|I - J_B\|_2, \|I - S^{-1}\bar{S}\|_2\} \\ &= \rho_L^{(J)} \epsilon_L.\end{aligned}$$

□

Furthermore, when the matrix J_B is positive definite, we can demonstrate the positive definiteness of the matrices $K_L = W^{(J)^{-1}}H$ and $K_R = HW^{(J)^{-1}}$.

Theorem 3.2 *Let the matrix J_B be positive definite. Then*

(i) *the matrix $K_L = W^{(J)^{-1}}H$ is positive definite, provided $\epsilon_L < \delta_L^{(J)}$, where*

$$\delta_L^{(J)} = \frac{2\left(\Theta\Gamma_s + 2 - \sqrt{\Theta^2\Gamma_s^2 + (\Theta + \Gamma_s)^2}\right)}{4 - (\Theta - \Gamma_s)^2} < 1;$$

(ii) *the matrix $K_R = HW^{(J)^{-1}}$ is positive definite, provided $\epsilon_R < \delta_R^{(J)}$, where*

$$\delta_R^{(J)} = \frac{2\left(\Theta_s\Gamma + 2 - \sqrt{\Theta_s^2\Gamma^2 + (\Theta_s + \Gamma)^2}\right)}{4 - (\Theta_s - \Gamma)^2} < 1.$$

Proof. We only prove the validity of (i), as (ii) can be demonstrated similarly.

Some straightforward computations immediately show that $\delta_L^{(J)} < 1$. Denote by

$$T = \begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix} \equiv \frac{1}{2}(K_L + K_L^T).$$

Then from the proof of Theorem 3.1 we easily obtain

$$\begin{cases} T_{11} &= \frac{1}{2}(J_B + J_B^T), \\ T_{12} &= \frac{1}{2}[(I - J_B)\bar{E} + (I - J_B^T)\bar{E}^T S^{-T}], \\ T_{21} &= \frac{1}{2}[S^{-1}\bar{F}(I - J_B) + \bar{E}^T(I - J_B^T)], \\ T_{22} &= \frac{1}{2}[S^{-1}\bar{S} + \bar{S}^T S^{-T}] - \frac{1}{2}[S^{-1}\bar{\Delta}_1 + \bar{\Delta}_1^T S^{-T}]. \end{cases}$$

Because J_B is positive definite, we know that its symmetric part $\frac{1}{2}(J_B + J_B^T)$ is symmetric positive definite. Therefore, the matrix T is symmetric positive definite if and only if so is its Schur complement $S_T := T_{22} - T_{21}T_{11}^{-1}T_{12}$.

Since

$$\|I - T_{11}\|_2 \leq \|I - J_B\|_2 \leq \epsilon_L < \delta_L^{(J)} < 1,$$

we have

$$\|T_{11}^{-1}\|_2 = \|[I - (I - T_{11})]^{-1}\|_2 \leq \frac{1}{1 - \|I - J_B\|_2}.$$

By direct computations we immediately obtain

$$\begin{aligned}\|T_{12}\|_2 = \|T_{21}\|_2 &\leq \frac{1}{2} \left(\|S^{-1}\overline{F}\|_2 \|I - J_B\|_2 + \|\overline{E}^T\|_2 \|I - J_B^T\|_2 \right) \\ &= \frac{1}{2} (\Theta + \Gamma_s) \|I - J_B\|_2\end{aligned}$$

and

$$\begin{aligned}\|I - T_{22}\|_2 &\leq \frac{1}{2} \left(\|I - S^{-1}\overline{S}\|_2 + \|I - \overline{S}^T S^{-T}\|_2 \right) \\ &\quad + \frac{1}{2} \left(\|S^{-1}\overline{\Delta}_1\|_2 + \|\overline{\Delta}_1^T S^{-T}\|_2 \right) \\ &\leq \|I - S^{-1}\overline{S}\|_2 + \|S^{-1}\overline{F}\|_2 \|\overline{E}\|_2 \|I - J_B\|_2 \\ &\leq (1 + \Theta\Gamma_s) \cdot \max\{\|I - J_B\|_2, \|I - S^{-1}\overline{S}\|_2\} \\ &= (1 + \Theta\Gamma_s)\epsilon_L.\end{aligned}$$

It then follows that

$$\begin{aligned}\min_{x \neq 0} \frac{\langle x, S_T x \rangle}{\langle x, x \rangle} &\geq 1 - \max_{x \neq 0} \frac{\langle x, (I - T_{22})x \rangle}{\langle x, x \rangle} - \max_{x \neq 0} \frac{\langle x, T_{21}T_{11}^{-1}T_{12}x \rangle}{\langle x, x \rangle} \\ &\geq 1 - \|I - T_{22}\|_2 - \|T_{21}T_{11}^{-1}T_{12}\|_2 \\ &\geq 1 - \left((1 + \Theta\Gamma_s)\epsilon_L + \frac{(\Theta + \Gamma_s)^2 \|I - J_B\|_2^2}{4(1 - \|I - J_B\|_2)} \right) \\ &\geq 1 - \left(1 + \Theta\Gamma_s + \frac{(\Theta + \Gamma_s)^2}{4(1 - \|I - J_B\|_2)} \cdot \|I - J_B\|_2 \right) \epsilon_L \\ &\geq 1 - \left(1 + \Theta\Gamma_s + \frac{(\Theta + \Gamma_s)^2}{4(1 - \epsilon_L)} \cdot \epsilon_L \right) \epsilon_L.\end{aligned}$$

Noticing that

$$\left(1 + \Theta\Gamma_s + \frac{(\Theta + \Gamma_s)^2}{4(1 - \epsilon_L)} \cdot \epsilon_L \right) \epsilon_L < 1$$

holds if and only if

$$4\Theta\Gamma_s\epsilon_L + (\Theta - \Gamma_s)^2\epsilon_L^2 < 4(1 - \epsilon_L)^2,$$

or equivalently,

$$\epsilon_L < \frac{2 \left(\Theta\Gamma_s + 2 - \sqrt{\Theta^2\Gamma_s^2 + (\Theta + \Gamma_s)^2} \right)}{4 - (\Theta - \Gamma_s)^2},$$

we therefore know that $\min_{x \neq 0} \frac{\langle x, S_T x \rangle}{\langle x, x \rangle} > 0$ holds true when $\epsilon_L < \delta_L^{(J)}$. Hence, S_T is a symmetric positive definite matrix, and K_L is a positive definite matrix. \square

3.2 The MBGS-type Preconditioners

If the matrix $W \in \mathbb{R}^{n \times n}$ in (2.10) is taken to be the modified block Gauss-Seidel splitting matrix[6, 7] of the matrix \overline{W} in (2.7), i.e.,

$$W^{(GS)} := W = \begin{bmatrix} I & O \\ \overline{F}(I - J_B) & S \end{bmatrix}, \quad (3.4)$$

then we obtain the *modified block Gauss-Seidel-type (MBGS-type)* preconditioner $M = PW^{(GS)}Q$ to the original matrix $A \in \mathbb{R}^{n \times n}$.

The following theorem describes the eigenvalue distribution of the preconditioned matrix with respect to the MBGS-type preconditioner.

Theorem 3.3 *Let $M = PW^{(GS)}Q$ be the MBGS-type preconditioner to the block two-by-two matrix $A = PHQ \in \mathbb{R}^{n \times n}$ in (1.2), where P and Q are given by (2.6), H is given by (2.7), and $W^{(GS)}$ is defined by (3.4). Denote by $K_L = W^{(GS)-1}H$ and $K_R = HW^{(GS)-1}$, respectively. Then it holds that*

- (i) $\|I - K_L\|_2 \leq \rho_L^{(GS)} \epsilon_L$, with $\rho_L^{(GS)} = \gamma(\Theta) \cdot \gamma(\Gamma_s \|I - J_B\|_2) \cdot (1 + \Theta \Gamma_s)$; and
- (ii) $\|I - K_R\|_2 \leq \rho_R^{(GS)} \epsilon_R$, with $\rho_R^{(GS)} = \gamma(\Theta_s) \cdot \gamma(\Gamma \|I - J_B\|_2) \cdot (1 + \Theta_s \Gamma)$.

It follows from Lemma 3.2 as well as (2.11) and (2.13) that the eigenvalues of the matrices $M^{-1}A$ and AM^{-1} are located within a circle having center $(1, 0)$ and radii $\rho_L^{(GS)} \epsilon_L$ and $\rho_R^{(GS)} \epsilon_R$, respectively, and therefore, they are all within the circle $\mathcal{N}^{(GS)}$.

Proof. We only prove (i), as (ii) can be verified analogously.

From (2.7) and (3.4) we have

$$K_L = W^{(GS)-1}H = \begin{bmatrix} J_B & (I - J_B)\overline{E} \\ S^{-1}\overline{F}(I - J_B)^2 & S^{-1}\overline{S} - S^{-1}(\overline{\Delta}_1 + \overline{\Delta}_2) \end{bmatrix}.$$

Hence,

$$I - K_L = \begin{bmatrix} I & O \\ -S^{-1}\overline{F}(I - J_B) & I \end{bmatrix} \begin{bmatrix} I - J_B & O \\ O & I - S^{-1}\overline{S} + S^{-1}\overline{\Delta}_1 \end{bmatrix} \begin{bmatrix} I & -\overline{E} \\ O & I \end{bmatrix}.$$

By making use of Lemma 3.1 we can immediately obtain

$$\begin{aligned} \|I - K_L\|_2 &\leq \gamma(\|S^{-1}\overline{F}(I - J_B)\|_2) \cdot \gamma(\|\overline{E}\|_2) \\ &\quad \cdot \max\{\|I - J_B\|_2, \|I - S^{-1}\overline{S}\|_2 + \|S^{-1}\overline{F}\|_2 \|\overline{E}\|_2 \|I - J_B\|_2\} \\ &\leq \gamma(\Theta) \cdot \gamma(\Gamma_s \|I - J_B\|_2) \cdot (1 + \Theta \Gamma_s) \\ &\quad \cdot \max\{\|I - J_B\|_2, \|I - S^{-1}\overline{S}\|_2\} \\ &= \rho_L^{(GS)} \epsilon_L. \end{aligned}$$

□

Furthermore, when the matrix J_B is positive definite, we can demonstrate the positive definiteness of the matrices $K_L = W^{(GS)-1}H$ and $K_R = HW^{(GS)-1}$.

Theorem 3.4 *Let the matrix J_B be positive definite. Then*

(i) *the matrix $K_L = W^{(GS)^{-1}}H$ is positive definite, provided $\epsilon_L < \delta_L^{(GS)}$, where*

$$\delta_L^{(GS)} = \frac{2 \left(\Theta \Gamma_s + 2 - \sqrt{\Theta^2 \Gamma_s^2 + 4\Theta \Gamma_s + (\Theta + \Gamma_s)^2} \right)}{4 - (\Theta + \Gamma_s)^2} < 1;$$

(ii) *the matrix $K_R = HW^{(GS)^{-1}}$ is positive definite, provided $\epsilon_R < \delta_R^{(GS)}$, where*

$$\delta_R^{(GS)} = \frac{2 \left(\Theta_s \Gamma + 2 - \sqrt{\Theta_s^2 \Gamma^2 + 4\Theta_s \Gamma + (\Theta_s + (1 + \Theta_s \Gamma)\Gamma)^2} \right)}{4 - (\Theta_s + (1 + \Theta_s \Gamma)\Gamma)^2} < 1.$$

Proof. We only prove the validity of (i), as (ii) can be demonstrated similarly.

Some straightforward computations immediately show that $\delta_L^{(GS)} < 1$. Denote by

$$T = \begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix} \equiv \frac{1}{2}(K_L + K_L^T).$$

Then from the proof of Theorem 3.3 we can easily obtain

$$\begin{cases} T_{11} &= \frac{1}{2}(J_B + J_B^T), \\ T_{12} &= \frac{1}{2}[(I - J_B)\overline{E} + (I - J_B^T)^2 \overline{F}^T S^{-T}], \\ T_{21} &= \frac{1}{2}[S^{-1} \overline{F}(I - J_B)^2 + \overline{E}^T (I - J_B^T)], \\ T_{22} &= \frac{1}{2}[S^{-1} \overline{S} + \overline{S}^T S^{-T}] - \frac{1}{2}[S^{-1} \overline{\Delta}_1 + \overline{\Delta}_1^T S^{-T}] - \frac{1}{2}[S^{-1} \overline{\Delta}_2 + \overline{\Delta}_2^T S^{-T}]. \end{cases}$$

Because J_B is positive definite, we know that its symmetric part $\frac{1}{2}(J_B + J_B^T)$ is symmetric positive definite. Therefore, the matrix T is symmetric positive definite if and only if so is its Schur complement $S_T := T_{22} - T_{21}T_{11}^{-1}T_{12}$.

Since

$$\|I - T_{11}\|_2 \leq \|I - J_B\|_2 \leq \epsilon_L < \delta_L^{(GS)} < 1,$$

we have

$$\|T_{11}^{-1}\|_2 = \|[I - (I - T_{11})]^{-1}\|_2 \leq \frac{1}{1 - \|I - J_B\|_2}.$$

By direct computations we immediately get

$$\begin{aligned} \|T_{12}\|_2 = \|T_{21}\|_2 &\leq \frac{1}{2} \left(\|S^{-1} \overline{F}(I - J_B)^2\|_2 + \|\overline{E}^T (I - J_B^T)\|_2 \right) \\ &\leq \frac{1}{2} (\Theta + \Gamma_s \|I - J_B\|_2) \|I - J_B\|_2 \end{aligned}$$

and

$$\|I - T_{22}\|_2 \leq \frac{1}{2} \left(\|I - S^{-1} \overline{S}\|_2 + \|I - \overline{S}^T S^{-T}\|_2 \right)$$

$$\begin{aligned}
& + \frac{1}{2} \left(\|S^{-1}\overline{F}(I - J_B)\overline{E}\|_2 + \|\overline{E}^T(I - J_B^T)\overline{F}^T S^{-T}\|_2 \right) \\
& + \frac{1}{2} \left(\|S^{-1}\overline{F}(I - J_B)^2\overline{E}\|_2 + \|\overline{E}^T(I - J_B^T)^2\overline{F}^T S^{-T}\|_2 \right) \\
\leq & \|I - S^{-1}\overline{S}\|_2 + \Theta\Gamma_s(1 + \|I - J_B\|_2)\|I - J_B\|_2 \\
\leq & [1 + \Theta\Gamma_s(1 + \|I - J_B\|_2)] \cdot \max\{\|I - J_B\|_2, \|I - S^{-1}\overline{S}\|_2\} \\
\leq & [1 + \Theta\Gamma_s(1 + \epsilon_L)]\epsilon_L.
\end{aligned}$$

It then follows that

$$\begin{aligned}
\min_{x \neq 0} \frac{\langle x, S_T x \rangle}{\langle x, x \rangle} & \geq 1 - \max_{x \neq 0} \frac{\langle x, (I - T_{22})x \rangle}{\langle x, x \rangle} - \max_{x \neq 0} \frac{\langle x, T_{21}T_{11}^{-1}T_{12}x \rangle}{\langle x, x \rangle} \\
& \geq 1 - \|I - T_{22}\|_2 - \|T_{21}T_{11}^{-1}T_{12}\|_2 \\
& \geq 1 - [1 + \Theta\Gamma_s(1 + \epsilon_L)]\epsilon_L - \frac{(\Theta + \Gamma_s\|I - J_B\|_2)^2\|I - J_B\|_2^2}{4(1 - \|I - J_B\|_2)} \\
& \geq 1 - \left(1 + \Theta\Gamma_s(1 + \epsilon_L) + \frac{(\Theta + \Gamma_s\epsilon_L)^2}{4(1 - \epsilon_L)} \cdot \epsilon_L \right) \epsilon_L.
\end{aligned}$$

Noticing that

$$\left(1 + \Theta\Gamma_s(1 + \epsilon_L) + \frac{(\Theta + \Gamma_s\epsilon_L)^2}{4(1 - \epsilon_L)} \cdot \epsilon_L \right) \epsilon_L < 1$$

holds if and only if

$$4\Theta\Gamma_s(1 - \epsilon_L^2)\epsilon_L + (\Theta + \Gamma_s\epsilon_L)^2\epsilon_L^2 < 4(1 - \epsilon_L)^2.$$

And this inequality holds when

$$4\Theta\Gamma_s\epsilon_L + (\Theta + \Gamma_s)^2\epsilon_L^2 < 4(1 - \epsilon_L)^2,$$

or equivalently,

$$\epsilon_L < \frac{2 \left(\Theta\Gamma_s + 2 - \sqrt{\Theta^2\Gamma_s^2 + 4\Theta\Gamma_s + (\Theta + \Gamma_s)^2} \right)}{4 - (\Theta + \Gamma_s)^2}.$$

Therefore, we know that $\min_{x \neq 0} \frac{\langle x, S_T x \rangle}{\langle x, x \rangle} > 0$ holds true when $\epsilon_L < \delta_L^{(GS)}$. Hence, S_T is a symmetric positive definite matrix, and K_L is a positive definite matrix. \square

Alternatively, if the matrix $W \in \mathbb{R}^{n \times n}$ in (2.10) is taken to be the modified block Gauss-Seidel splitting matrix[6, 7] of the matrix \overline{W} in (2.7), i.e.,

$$W^{(GS)} := W = \begin{bmatrix} I & (I - J_B)\overline{E} \\ O & S \end{bmatrix}, \quad (3.5)$$

then we obtain another *modified block Gauss-Seidel-type (MBGS-type)* preconditioner $M = PW^{(GS)}Q$ to the original matrix $A \in \mathbb{R}^{n \times n}$. Exactly following the demonstrations of Theorems 3.3 and 3.4, we can obtain the following results for the eigenvalue distribution and the positive definiteness of the preconditioned matrix with respect to the MBGS-type preconditioner (3.5).

Theorem 3.5 *Let $M = PW^{(GS)}Q$ be the MBGS-type preconditioner to the block two-by-two matrix $A = PHQ \in \mathbb{R}^{n \times n}$ in (1.2), where P and Q are given by (2.6), H is given by (2.7), and $W^{(GS)}$ is defined by (3.5). Denote by $K_L = W^{(GS)^{-1}}H$ and $K_R = HW^{(GS)^{-1}}$, respectively. Then it holds that*

- (i) $\|I - K_L\|_2 \leq \rho_L^{(GS)} \epsilon_L$, with $\rho_L^{(GS)} = \gamma(\Theta\|I - J_B\|_2) \cdot \gamma(\Gamma_s) \cdot (1 + \Theta\Gamma_s)$; and
- (ii) $\|I - K_R\|_2 \leq \rho_R^{(GS)} \epsilon_R$, with $\rho_R^{(GS)} = \gamma(\Theta_s\|I - J_B\|_2) \cdot \gamma(\Gamma) \cdot (1 + \Theta_s\Gamma)$.

It follows from Lemma 3.2 as well as (2.11) and (2.13) that the eigenvalues of the matrices $M^{-1}A$ and AM^{-1} are located within a circle having center $(1, 0)$ and radii $\rho_L^{(GS)} \epsilon_L$ and $\rho_R^{(GS)} \epsilon_R$, respectively, and therefore, they are all within the circle $\mathcal{N}^{(GS)}$.

Theorem 3.6 *Let the matrix J_B be positive definite. Then*

- (i) *the matrix $K_L = W^{(GS)^{-1}}H$ is positive definite, provided $\epsilon_L < \delta_L^{(GS)}$, where*

$$\delta_L^{(GS)} = \frac{2 \left(\Theta\Gamma_s + 2 - \sqrt{\Theta^2\Gamma_s^2 + 4\Theta\Gamma_s + (\Gamma_s + (1 + \Theta\Gamma_s)\Theta)^2} \right)}{4 - (\Gamma_s + (1 + \Theta\Gamma_s)\Theta)^2} < 1;$$

- (ii) *the matrix $K_R = HW^{(GS)^{-1}}$ is positive definite, provided $\epsilon_R < \delta_R^{(GS)}$, where*

$$\delta_R^{(GS)} = \frac{2 \left(\Theta_s\Gamma + 2 - \sqrt{\Theta_s^2\Gamma^2 + 4\Theta_s\Gamma + (\Theta_s + \Gamma)^2} \right)}{4 - (\Theta_s + \Gamma)^2} < 1.$$

3.3 The MBUGS-type Preconditioners

If the matrix $W \in \mathbb{R}^{n \times n}$ in (2.10) is taken to be the modified block unsymmetric Gauss-Seidel splitting matrix[6, 7] of the matrix \overline{W} in (2.7), i.e.,

$$W^{(UGS)} := W = \begin{bmatrix} I & (I - J_B)\overline{E} \\ O & S \end{bmatrix} \begin{bmatrix} I & O \\ O & S \end{bmatrix}^{-1} \begin{bmatrix} I & O \\ \overline{F}(I - J_B) & S \end{bmatrix}, \quad (3.6)$$

then we obtain the *modified block unsymmetric Gauss-Seidel-type (MBUGS-type)* preconditioner $M = PW^{(UGS)}Q$ to the original matrix $A \in \mathbb{R}^{n \times n}$.

The following theorem describes the eigenvalue distribution of the preconditioned matrix with respect to the MBUGS-type preconditioner.

Theorem 3.7 *Let $M = PW^{(UGS)}Q$ be the MBUGS-type preconditioner to the block two-by-two matrix $A = PHQ \in \mathbb{R}^{n \times n}$ in (1.2), where P and Q are given by (2.6), H is given by (2.7), and $W^{(UGS)}$ is defined by (3.6). Denote by $K_L = W^{(UGS)^{-1}}H$ and $K_R = HW^{(UGS)^{-1}}$, respectively. Then it holds that*

- (i) $\|I - K_L\|_2 \leq \rho_L^{(UGS)} \epsilon_L$, with
- $$\rho_L^{(UGS)} = \gamma(\Gamma_s\|I - J_B\|_2) \cdot [\gamma(\Theta\|I - J_B\|_2) + \Theta\Gamma_s\|I - J_B\|_2] \cdot (1 + \Theta\Gamma_s); \text{ and}$$

(ii) $\|I - K_R\|_2 \leq \rho_R^{(UGS)} \epsilon_R$, with

$$\rho_R^{(UGS)} = \gamma(\Theta_s \|I - J_B\|_2) \cdot [\gamma(\Gamma \|I - J_B\|_2) + \Theta_s \Gamma \|I - J_B\|_2] \cdot (1 + \Theta_s \Gamma).$$

It follows from Lemma 3.2 as well as (2.11) and (2.13) that the eigenvalues of the matrices $M^{-1}A$ and AM^{-1} are located within a circle having center $(1, 0)$ and radii $\rho_L^{(UGS)} \epsilon_L$ and $\rho_R^{(UGS)} \epsilon_R$, respectively, and therefore, they are all within the circle $\mathcal{N}^{(UGS)}$.

Proof. It is analogous to the proofs of Theorems 3.1 and 3.3, and is hence omitted. \square

Furthermore, when the matrix J_B is positive definite, we can demonstrate the positive definiteness of the matrices $K_L = W^{(UGS)^{-1}}H$ and $K_R = HW^{(UGS)^{-1}}$.

Theorem 3.8 *Let the matrix J_B be positive definite. Then*

(i) *the matrix $K_L = W^{(UGS)^{-1}}H$ is positive definite, provided $\epsilon_L < \delta_L^{(UGS)}$, where*

$$\delta_L^{(UGS)} = \frac{2 \left(\Theta \Gamma_s + 2 - \sqrt{\Theta^2 \Gamma_s^2 + 4\Theta \Gamma_s + (1 + \Theta \Gamma_s)^2 (\Theta + \Gamma_s)^2} \right)}{4 - (1 + \Theta \Gamma_s)^2 (\Theta + \Gamma_s)^2} < 1;$$

(ii) *the matrix $K_R = HW^{(UGS)^{-1}}$ is positive definite, provided $\epsilon_R < \delta_R^{(UGS)}$, where*

$$\delta_R^{(UGS)} = \frac{2 \left(\Theta_s \Gamma + 2 - \sqrt{\Theta_s^2 \Gamma^2 + 4\Theta_s \Gamma + (1 + \Theta_s \Gamma)^2 (\Theta_s + \Gamma)^2} \right)}{4 - (1 + \Theta_s \Gamma)^2 (\Theta_s + \Gamma)^2} < 1.$$

Proof. It is analogous to the proofs of Theorems 3.2 and 3.4, and is hence omitted. \square

Alternatively, if the matrix $W^{(UGS)}$ defined by (3.6) is considered to possess the split form $W^{(UGS)} = W_L^{(UGS)} W_R^{(UGS)}$, with

$$W_L^{(UGS)} = \begin{bmatrix} I & (I - J_B) \overline{E} S^{-1} \\ O & I \end{bmatrix}, \quad W_R^{(UGS)} = \begin{bmatrix} I & O \\ \overline{F} (I - J_B) & S \end{bmatrix} \quad (3.7)$$

or

$$W_L^{(UGS)} = \begin{bmatrix} I & (I - J_B) \overline{E} \\ O & S \end{bmatrix}, \quad W_R^{(UGS)} = \begin{bmatrix} I & O \\ S^{-1} \overline{F} (I - J_B) & I \end{bmatrix}, \quad (3.8)$$

then we can obtain other *modified block unsymmetric Gauss-Seidel-type* preconditioners $M = M_L^{(UGS)} M_R^{(UGS)}$ to the original matrix $A \in \mathbb{R}^{n \times n}$, where

$$M_L^{(UGS)} = P W_L^{(UGS)} \quad \text{and} \quad M_R^{(UGS)} = W_R^{(UGS)} Q,$$

and P and Q are given by (2.6). Exactly following the demonstrations of Theorems 3.7 and 3.8, we can obtain the results about the eigenvalue distributions and the positive definiteness of the preconditioned matrices with respect to the MBUGS-type preconditioners (3.7)-(3.8).

We remark that when $\overline{F} = \overline{E}^T$, the above-discussed modified block unsymmetric Gauss-Seidel-type preconditioners naturally reduce to the *modified block symmetric Gauss-Seidel-type (MBSGS-type)* preconditioners to the matrix $A \in \mathbb{R}^{n \times n}$ in (1.2), correspondingly.

3.4 The Case $S \approx -\bar{S}$

In the case that \bar{S} is negative definite, we may let S be an approximation to $-\bar{S}$ in order to obtain a preconditioner of positive definiteness in nature. Hence, some specified preconditioned Krylov subspace iteration method can exploit its efficiency sufficiently.

When $S \approx -\bar{S}$, for the MBJ-, the MBGS-, and the MBUGS-type preconditioners discussed above, we can demonstrate that the eigenvalues of the preconditioned matrices are, correspondingly, located within two circles having center $(-1, 0)$ and $(1, 0)$ in the complex plane. This results are precisely summarized in the following theorem. Since their proofs are essentially the same to those of Theorems 3.1, 3.3, 3.5 and 3.7 with only the identity matrix I being replaced by the matrix

$$J := \begin{bmatrix} I & O \\ O & -I \end{bmatrix}, \quad I \in \mathbb{R}^{p \times p} \quad \text{and} \quad -I \in \mathbb{R}^{q \times q},$$

we only state the theorem but omit its proof.

Theorem 3.9 *Let $M = PWQ \in \mathbb{R}^{n \times n}$ in (2.10) be the preconditioner to the block two-by-two matrix $A = PHQ \in \mathbb{R}^{n \times n}$ in (1.2), with P and Q being given by (2.6) and H being given by (2.7). Denote by $K_L = W^{-1}H$ and $K_R = HW^{-1}$, respectively.*

(i) *If $W = W^{(J)}$ is defined by (3.3), then*

$$\|J - K_L\|_2 \leq \rho_L^{(J)} \tilde{\epsilon}_L, \quad \|J - K_R\|_2 \leq \rho_R^{(J)} \tilde{\epsilon}_R,$$

where $\rho_L^{(J)}$ and $\rho_R^{(J)}$ are the same as in Theorem 3.1;

(ii) *If $W = W^{(GS)}$ is defined by (3.4), then*

$$\|J - K_L\|_2 \leq \rho_L^{(GS)} \tilde{\epsilon}_L, \quad \|J - K_R\|_2 \leq \rho_R^{(GS)} \tilde{\epsilon}_R,$$

where $\rho_L^{(GS)}$ and $\rho_R^{(GS)}$ are the same as in Theorem 3.3;

(iii) *If $W = W^{(GS)}$ is defined by (3.5), then*

$$\|J - K_L\|_2 \leq \rho_L^{(GS)} \tilde{\epsilon}_L, \quad \|J - K_R\|_2 \leq \rho_R^{(GS)} \tilde{\epsilon}_R,$$

where $\rho_L^{(GS)}$ and $\rho_R^{(GS)}$ are the same as in Theorem 3.5;

(iv) *If $W = W^{(UGS)}$ is defined by (3.6), then*

$$\|J - K_L\|_2 \leq \rho_L^{(UGS)} \tilde{\epsilon}_L, \quad \|J - K_R\|_2 \leq \rho_R^{(UGS)} \tilde{\epsilon}_R,$$

where $\rho_L^{(UGS)}$ and $\rho_R^{(UGS)}$ are the same as in Theorem 3.7.

It follows from Lemma 3.2 as well as (2.11) and (2.13) that the eigenvalues of the preconditioned matrix $M^{-1}A$ are located within the union of two circles having centers $(-1, 0)$ and $(1, 0)$ and radius $\rho_L^{(\xi)} \tilde{\epsilon}_L$, and those of the preconditioned matrix AM^{-1} are located within the union of two circles having centers $(-1, 0)$ and $(1, 0)$ and radius $\rho_R^{(\xi)} \tilde{\epsilon}_R$, respectively. Therefore, they are all within $\tilde{\mathcal{N}}^{(\xi)}$. Here, $\xi = J, GS$ and UGS .

We observe from the demonstrations of Theorems 3.1-3.9 that when $J_B = I$ or $S = \bar{S}$, the results in these theorems can be considerably improved and accurated.

3.5 Connections to Krylov Subspace Methods

The preconditioning matrix M defined in (2.10) can be used to accelerate the Krylov subspace methods such as GMRES or its restarted variant GMRES(m) [41, 40] for solving the large sparse system of linear equations (1.1)-(1.2). This preconditioning matrix can be used as a left (see (2.11)-(2.12)), a right (see (2.13)-(2.14)), or a split (see (2.15)-(2.17)) preconditioner to the system of linear equations (1.1). The obtained equivalent linear systems can be solved by GMRES or GMRES(m).

Assume that the coefficient matrices \mathbf{A} of the above preconditioned linear systems are diagonalizable, i.e., there exist a nonsingular matrix $\mathbf{X} \in \mathbb{C}^{n \times n}$ and a diagonal matrix $\mathbf{D} \in \mathbb{C}^{n \times n}$ such that $\mathbf{A} = \mathbf{X}\mathbf{D}\mathbf{X}^{-1}$. Then it is well known from [41, Theorem 4] that the residual norm $\|\mathbf{r}^{(k)}\|_2$ at the k -th step of the preconditioned GMRES is bounded by $\|\mathbf{r}^{(k)}\|_2 \leq \kappa(\mathbf{X})\varepsilon^{(k)}\|\mathbf{r}^{(0)}\|_2$, where $\kappa(\mathbf{X})$ is the Euclidean condition number of \mathbf{X} and $\varepsilon^{(k)} := \min_{\mathcal{P} \in \mathcal{P}_k} \max_{\lambda_i \in \sigma(\mathbf{A})} |\mathcal{P}(\lambda_i)|$. Here, \mathcal{P}_k denotes the set of all polynomials $\mathcal{P}(\lambda)$ of degree not greater than k such that $\mathcal{P}(0) = 1$, and $\sigma(\mathbf{A})$ denotes the spectrum of the matrix \mathbf{A} .

Consider \bar{S} defined by (3.2), see also (2.1) and (2.3). When the matrix S is an approximation to the matrix \bar{S} , from Theorems 3.1, 3.3, 3.5 and 3.7 we know that all eigenvalues of the matrix \mathbf{A} are contained in either of the circles $\mathcal{N}^{(\xi)}$, $\xi = J, GS$ and UGS . Therefore, when $\rho^{(\xi)} < 1$, a special case of Theorem 5 in [41] implies that $\varepsilon^{(k)} \leq (\rho^{(\xi)})^k$, $\xi = J, GS$ and UGS .

Alternatively, the preconditioning matrix M can also be used as a left, a right, or a split preconditioner to the system of linear equations (1.1) to obtain a preconditioned linear system of coefficient matrix $\tilde{\mathbf{A}} = K_L, K_R$, or K , respectively. Because Theorems 3.2, 3.4, 3.6 and 3.8 guarantee the positive definiteness of the preconditioned matrix $\tilde{\mathbf{A}}$, it is known from [20] and [41, pp. 866] that the following error bound for the correspondingly preconditioned GMRES holds:

$$\|\mathbf{r}^{(k)}\|_2 \leq \left(1 - \frac{(\lambda_{\min}(\tilde{\mathbf{H}}))^2}{\lambda_{\max}(\tilde{\mathbf{A}}^T \tilde{\mathbf{A}})}\right)^{\frac{k}{2}} \|\mathbf{r}^{(0)}\|_2,$$

where $\tilde{\mathbf{H}} = \frac{1}{2}(\tilde{\mathbf{A}} + \tilde{\mathbf{A}}^T)$ denotes the symmetric part of the matrix $\tilde{\mathbf{A}}$, and $\lambda_{\min}(\cdot)$ and $\lambda_{\max}(\cdot)$ denote, respectively, the smallest and the largest eigenvalues of the corresponding matrix. This gives a guarantee for the convergence of the restarted preconditioned GMRES iteration, say PGMRES(m), for all m , when the coefficient matrix $\tilde{\mathbf{A}}$ is positive definite.

When the matrix S is an approximation to the matrix $-\bar{S}$, because the preconditioned matrix \mathbf{A} or $\tilde{\mathbf{A}}$ may be usually not positive definite, instead of GMRES and GMRES(m) we may use other Krylov subspace methods such as BiCGSTAB, QMR and TFQMR to solve the preconditioned linear systems. In particular, when the original coefficient matrix A is symmetric indefinite, MINRES is a possible candidate if a symmetric positive definite or indefinite preconditioner M is obtainable. See [2, 27, 40].

4 Applications to Three Typical Matrices

In this section, we will investigate the concretizations of the structured preconditioners established in Sections 2 and 3 to three special classes of matrices arising from the real-world applications.

4.1 The Symmetric Positive Definite Matrix

When the matrix blocks $B \in \mathbb{R}^{p \times p}$ and $C \in \mathbb{R}^{q \times q}$ are symmetric positive definite, $F = E^T$ and the Schur complement $S_A = C - E^T B^{-1} E$ is symmetric positive definite, the matrix $A \in \mathbb{R}^{n \times n}$ reduces to the block two-by-two symmetric positive definite matrix

$$A = \begin{bmatrix} B & E \\ E^T & C \end{bmatrix}.$$

This kind of matrices may arise in the red/black ordering of a symmetric positive definite linear system, or in discretization incorporated with domain decomposition technique of a boundary value problem of a self-adjoint elliptic partial differential equation, etc. See [2, 3, 6, 7, 27, 40].

Let $L_B \in \mathbb{R}^{p \times p}$ and $L_C \in \mathbb{R}^{q \times q}$ be nonsingular matrices such that either (2.1) or (2.2) holds with $R_B = L_B^T$ and $R_C = L_C^T$. Then from (2.10) and (2.6) we know that $M = PWQ$ is the structured preconditioner to the matrix A , where

$$P = \begin{bmatrix} L_B & O \\ E^T L_B^{-T} & L_C \end{bmatrix}, \quad Q = \begin{bmatrix} L_B^T & L_B^{-1} E \\ O & L_C^T \end{bmatrix} = P^T,$$

and $W \in \mathbb{R}^{n \times n}$ is an approximation to the matrix

$$\overline{W} = \begin{bmatrix} J_B & (I - J_B)\overline{E} \\ \overline{E}^T(I - J_B) & \overline{S} \end{bmatrix} \approx H$$

defined by (2.7), with $\overline{E} = L_B^{-1} E L_C^{-T}$ and $\overline{S} = J_C - \overline{E}^T \overline{E}$.

Note that \overline{S} and \overline{W} are symmetric positive definite. Let $S \in \mathbb{R}^{q \times q}$ be an approximation to the matrix $I - \overline{E}^T \overline{E} \approx \overline{S}$. To guarantee the symmetric positive definiteness of the preconditioning matrix M , we can choose W to be the modified block Jacobi splitting matrix in (3.3) or the modified block symmetric Gauss-Seidel splitting matrix in (3.6), obtaining the modified block Jacobi-type preconditioner or the modified block symmetric Gauss-Seidel-type preconditioner to the matrix A , respectively.

4.2 The Saddle Point Matrix

When the matrix block $B \in \mathbb{R}^{p \times p}$ is symmetric positive definite, $C = O$ and $F = \pm E^T$ is of full row rank, the matrix $A \in \mathbb{R}^{n \times n}$ reduces to the saddle point matrices

$$A^\pm = \begin{bmatrix} B & E \\ \pm E^T & O \end{bmatrix}.$$

This kind of matrices may arise in constrained optimization as well as least-squares, saddle-point and Stokes problems, without a regularizing/stabilizing term, etc. See [14, 16, 24, 25, 28, 37, 44].

Let $L_B \in \mathbb{R}^{p \times p}$ be a nonsingular matrix such that either (2.1) or (2.2) holds with $R_B = L_B^T$ and $L_C = R_C = I$. Then from (2.10) and (2.6) we know that $M^\pm = P^\pm W^\pm Q^\pm$ are the preconditioners to the matrices A^\pm , respectively, where

$$P^\pm = \begin{bmatrix} L_B & O \\ \pm E^T L_B^{-T} & I \end{bmatrix}, \quad Q^\pm = \begin{bmatrix} L_B^T & L_B^{-1} E \\ O & I \end{bmatrix},$$

and $W^\pm \in \mathbb{R}^{n \times n}$ are approximations to the matrices

$$\overline{W}^\pm = \begin{bmatrix} J_B & (I - J_B)\overline{E} \\ \pm \overline{E}^T (I - J_B) & \overline{S}^\pm \end{bmatrix} \approx H^\pm$$

defined by (2.7), with $\overline{E} = L_B^{-1}E$ and $\overline{S}^\pm = \mp \overline{E}^T \overline{E}$.

Let $S^\pm \in \mathbb{R}^{q \times q}$ be approximations to the matrices \overline{S}^\pm . By choosing the matrices W^\pm to be the modified block Jacobi splitting matrices in (3.3), the modified block Gauss-Seidel splitting matrices in (3.4) or (3.5), or the modified block unsymmetric Gauss-Seidel splitting matrices in (3.6), we can obtain the modified block Jacobi-type preconditioners, the modified block Gauss-Seidel-type preconditioners, or the modified block unsymmetric Gauss-Seidel-type preconditioners to the matrices A^\pm , respectively.

4.3 The Hamiltonian Matrix

When the matrix blocks $B \in \mathbb{R}^{p \times p}$ is symmetric positive definite and $C \in \mathbb{R}^{q \times q}$ is symmetric positive/negative definite (denoted by C_+/C_- , respectively), and $F = \mp E^T$, the matrix $A \in \mathbb{R}^{n \times n}$ reduces to the Hamiltonian matrices

$$A^\pm = \begin{bmatrix} B & E \\ \mp E^T & C_\pm \end{bmatrix}.$$

This kind of matrices may arise in stationary semiconductor devices [36, 43, 42], in constrained optimization as well as least-squares, saddle-point and Stokes problems, with a regularizing/stabilizing term [28].

Let $L_B \in \mathbb{R}^{p \times p}$ and $L_{C_\pm} \in \mathbb{R}^{q \times q}$ be nonsingular matrices such that either (2.1) or (2.2) holds with $R_B = L_B^T$ and $R_{C_\pm} = L_{C_\pm}^T$. Then from (2.10) and (2.6) we know that $M^\pm = P^\pm W^\pm Q^\pm$ are the preconditioners to the matrices A^\pm , where

$$P^\pm = \begin{bmatrix} L_B & O \\ \mp E^T L_B^{-T} & L_{C_\pm} \end{bmatrix}, \quad Q^\pm = \begin{bmatrix} L_B^T & L_B^{-1}E \\ O & L_{C_\pm}^T \end{bmatrix},$$

and $W^\pm \in \mathbb{R}^{n \times n}$ are approximations to the matrices

$$\overline{W}^\pm = \begin{bmatrix} J_B & (I - J_B)\overline{E}^\pm \\ \mp (\overline{E}^\pm)^T (I - J_B) & \overline{S}^\pm \end{bmatrix} \approx H^\pm$$

defined by (2.7), with $\overline{E}^\pm = L_B^{-1}EL_{C_\pm}^{-T}$ and $\overline{S}^\pm = J_C \pm (\overline{E}^\pm)^T \overline{E}^\pm$.

Let $S^\pm \in \mathbb{R}^{q \times q}$ be approximations to the matrices $I \pm (\overline{E}^\pm)^T \overline{E}^\pm \approx \overline{S}^\pm$. By choosing the matrices W^\pm to be the modified block Jacobi splitting matrices in (3.3), the modified block Gauss-Seidel splitting matrices in (3.4) or (3.5), or the modified block unsymmetric Gauss-Seidel splitting matrices in (3.6), we can obtain the modified block Jacobi-type preconditioners, the modified block Gauss-Seidel-type preconditioners, or the modified block unsymmetric Gauss-Seidel-type preconditioners to the matrices A^\pm , respectively.

4.4 An Illustrative Example

Let us consider the electromagnetic scattering problem from a large rectangular cavity on the x - y plane in which the medium is y -directional inhomogeneous. In the transverse magnetic polarization case, when the model Helmholtz equation with positive wave number is discretized by the five-point finite difference scheme with uniform stepsize h , we obtain a block two-by-two system of linear equations (1.1)-(1.2), in which

$$B = V \otimes I + I \otimes V - I \otimes \Omega \in \mathbb{R}^{p \times p}, \quad C = I - hG \in \mathbb{R}^{q \times q}, \quad E = I \otimes e_q \in \mathbb{R}^{p \times q}$$

and $F = -E^T$, where $h = \frac{1}{q+1}$, $p = q^2$, $\theta \geq 0$ is a real constant, e_q is the q -th unit vector in \mathbb{R}^q , I is the q -by- q identity matrix, $V = \text{tridiag}(-1 + \frac{1}{2}\theta h, 2, -1 - \frac{1}{2}\theta h) \in \mathbb{R}^{q \times q}$ is a tridiagonal matrix, $\Omega = h^2 \cdot \text{diag}(\omega_1^2, \omega_2^2, \dots, \omega_q^2) \in \mathbb{R}^{q \times q}$ is a nonnegative diagonal matrix, $G = (g_{ij}) \in \mathbb{R}^{q \times q}$, and \otimes denotes the Kronecker product. See [33, 1].

Concretely, in our computations we take $\theta = 1$, $\omega_i = 16\pi$ ($i = 1, 2, \dots, q$), and $g_{ij} = \frac{1}{(i+j)^2}$ ($i, j = 1, 2, \dots, q$).

Let $B \approx L_B R_B$ be an incomplete triangular factorization of the matrix block B , and $L_C = R_C = I$. Then we have

$$\overline{E} = L_B^{-1} E, \quad \overline{F} = -E^T R_B^{-1}, \quad J_C = C \quad \text{and} \quad \overline{S} = C - \overline{F} \overline{E}.$$

Now, by choosing $S = \text{band}_{\ell_b}(C) - \overline{F} \overline{E}$ with $\text{band}_{\ell_b}(C)$ being the band matrix of half-band width ℓ_b truncated from the matrix C , after straightforward computations we can obtain the results listed in Tables 1-4 for the discretization stepsizes $h = \frac{1}{16}, \frac{1}{24}, \frac{1}{32}$ and $\frac{1}{64}$, or equivalently, for the problem sizes $(p, q) = (225, 15), (529, 23), (961, 31)$ and $(3969, 63)$, respectively.

In Table 1 we list the half-band width ℓ_b , the quantities

$$\Theta = \|\overline{E}\|_2, \quad \Gamma = \|\overline{F}\|_2, \quad \Theta_s = \|\overline{E} S^{-1}\|_2 \quad \text{and} \quad \Gamma_s = \|S^{-1} \overline{F}\|_2$$

with respect to the matrix norms, and

$$\epsilon_L = \max\{\|I - J_B\|_2, \|I - S^{-1} \overline{S}\|_2\} \quad \text{and} \quad \epsilon_R = \max\{\|I - J_B\|_2, \|I - \overline{S} S^{-1}\|_2\}$$

with respect to the matrix approximation accuracies. For $\xi = J, GS$ and UGS , in Tables 2-4 we list the radii $\rho_L^{(\xi)} \epsilon_L$ and $\rho_R^{(\xi)} \epsilon_R$ of the circles centered at $(1, 0)$ where all eigenvalues of the matrices K_L and K_R are located within, $\rho^{(\xi)} = \min\{\rho_L^{(J)} \epsilon_L, \rho_R^{(J)} \epsilon_R\}$ and the radii $\rho_*^{(\xi)}$ of the smallest circles that include all eigenvalues of the corresponding preconditioned matrices (see Theorems 3.1, 3.3 and 3.7), and the quantities $\delta_L^{(\xi)}$ and $\delta_R^{(\xi)}$ that guarantee the positive definiteness of the preconditioned matrices K_L and K_R whenever $\epsilon_L < \delta_L^{(\xi)}$ and $\epsilon_R < \delta_R^{(\xi)}$ (see Theorems 3.2, 3.4 and 3.8), respectively.

Table 1: Quantities with respect to the preconditioned matrices

h	$\frac{1}{16}$	$\frac{1}{24}$	$\frac{1}{32}$	$\frac{1}{64}$
ℓ_b	2	4	6	30
Θ	11.9704	15.8432	25.3679	28.8844
Γ	6.05339	7.38068	8.95453	14.7829
Θ_s	5.77108	6.57523	11.0775	39.2410
Γ_s	2.82323	3.28938	3.93560	19.5354
ϵ_L	2.95e-3	1.52e-3	6.39e-04	5.75e-05
ϵ_R	4.13e-3	2.15e-3	7.89e-04	4.63e-05

Table 2: Bounds with respect to the MBJ-type preconditioner $M = PW^{(J)}Q$ with $W^{(J)}$ being defined by (3.3)

h	$\frac{1}{16}$	$\frac{1}{24}$	$\frac{1}{32}$	$\frac{1}{64}$
$\rho_L^{(J)}\epsilon_L$	0.111566	8.62e-02	6.78e-02	3.26e-02
$\rho_R^{(J)}\epsilon_R$	0.152507	0.108497	7.98e-02	2.70e-02
$\rho^{(J)}$	0.111566	8.62e-02	6.78e-02	2.70e-02
$\delta_L^{(J)}$	2.75e-02	1.82e-02	9.71e-03	1.77e-02
$\delta_R^{(J)}$	2.71e-02	1.98e-02	9.88e-03	1.72e-03
$\rho_*^{(J)}$	2.31e-03	1.24e-03	4.63e-04	3.93e-05

Table 3: Bounds with respect to the MBGS-type preconditioner $M = PW^{(GS)}Q$ with $W^{(GS)}$ being defined by (3.4)

h	$\frac{1}{16}$	$\frac{1}{24}$	$\frac{1}{32}$	$\frac{1}{64}$
$\rho_L^{(GS)}\epsilon_L$	3.55e-02	2.41e-02	1.62e-02	1.66e-03
$\rho_R^{(GS)}\epsilon_R$	2.45e-02	1.44e-02	8.81e-03	1.82e-03
$\rho^{(GS)}$	2.45e-02	1.44e-02	8.81e-03	1.66e-03
$\delta_L^{(GS)}$	2.69e-02	1.79e-02	9.63e-03	1.76e-03
$\delta_R^{(GS)}$	7.60e-03	4.69e-03	1.97e-03	2.17e-04
$\rho_*^{(GS)}$	2.31e-03	1.24e-03	4.63e-04	3.93e-05

Table 4: Bounds with respect to the MBUGS-type preconditioner $M = PW^{(UGS)}Q$ with $W^{(UGS)}$ being defined by (3.6)

h	$\frac{1}{16}$	$\frac{1}{24}$	$\frac{1}{32}$	$\frac{1}{64}$
$\rho_L^{(UGS)}\epsilon_L$	0.102518	8.06e-02	0.166135	3.25e-02
$\rho_R^{(UGS)}\epsilon_R$	0.148479	0.106365	0.216704	2.69e-02
$\rho^{(UGS)}$	0.102518	8.06e-02	0.166135	2.69e-02
$\delta_L^{(UGS)}$	3.62e-03	1.87e-03	6.54e-04	7.16e-05
$\delta_R^{(UGS)}$	4.32e-03	2.69e-03	9.47e-04	6.25e-05
$\rho_*^{(UGS)}$	2.31e-03	1.24e-03	4.63e-04	3.93e-05

The results in Tables 2-4 clearly show that

- (i) for $\xi = J, GS$ and UGS , $\rho_L^{(\xi)} \epsilon_L < 1$ and $\rho_R^{(\xi)} \epsilon_R < 1$. It follows that $\rho^{(\xi)} < 1$. As $\rho_L^{(\xi)} \epsilon_L$ and $\rho_R^{(\xi)} \epsilon_R$ are quite small, the eigenvalues of the preconditioned matrices, with respect to the MBJ-, the MBGS- and the MBUGS-type preconditioners, are tightly clustered around the point $(1, 0)$, see Theorems 3.1, 3.3 and 3.7. Hence, a Krylov subspace method such as GMRES, when applied to the preconditioned systems of linear equations, will achieve fast convergence, see Section 3.5;
- (ii) for $\xi = J, GS$ and UGS , $\epsilon_L < \delta_L^{(\xi)} < 1$ and $\epsilon_R < \delta_R^{(\xi)} < 1$. It follows that the preconditioned matrices, with respect to the MBJ-, the MBGS- and the MBUGS-type preconditioners, are positive definite, and the convergence of the restarted GMRES methods preconditioned by these preconditioners are guaranteed, see Theorems 3.2, 3.4 and 3.8 as well as Section 3.5;
- (iii) for $\xi = J, GS$ and UGS , $\rho_*^{(\xi)} \leq \rho^{(\xi)}$. This shows that the eigenvalues of the preconditioned matrices, with respect to the MBJ-, the MBGS- and the MBUGS-type preconditioners, are really located within the theoretically estimated circles centered at $(1, 0)$ with radii $\rho^{(\xi)}$ given in Theorems 3.1, 3.3 and 3.7, respectively.

In summary, this example shows that the conditions of our theorems are reasonable and the conclusions of them are correct.

5 Conclusion and Remarks

We have established a general framework of practical and efficient structured preconditioners to the large sparse block two-by-two nonsingular matrices. For several special cases associated with the modified block relaxation iteration methods, we have studied the eigenvalue distributions and the positive definiteness of the preconditioned matrices. Theoretical analyses have shown that this preconditioning technique can afford effective and high-quality preconditioners to the Krylov subspace iteration methods for solving large sparse systems of linear equations with block two-by-two coefficient matrices.

We remark that our preconditioning technique and the corresponding theory can be straightforwardly developed to the following cases:

- (a) The approximation matrix $W \in \mathbb{R}^{n \times n}$ in (2.10) that is generated by a multi-step variant of the modified block Jacobi, the modified block Gauss-Seidel or the modified block unsymmetric Gauss-Seidel splitting matrix of the matrix $\overline{W} \in \mathbb{R}^{n \times n}$ in (2.7)[6, 7];
- (b) Alternatively, the approximation matrix $W \in \mathbb{R}^{n \times n}$ in (2.10) that is generated by a single- or multiple-step variant of the modified block *successive overrelaxation* (**SOR**), the modified block unsymmetric SOR, the modified block *accelerated overrelaxation* (**AOR**) or the modified block unsymmetric AOR splitting matrix of the matrix $\overline{W} \in \mathbb{R}^{n \times n}$ in (2.7)[32, 6, 7];
- (c) And more generally, the approximation matrix $W \in \mathbb{R}^{n \times n}$ in (2.10) that is generated by any suitable direct or iterative method induced by the matrix $\overline{W} \in \mathbb{R}^{n \times n}$ in (2.7);

- (d) The matrix $A \in \mathbb{R}^{n \times n}$ that is of a general ℓ -by- ℓ block structure. More concretely, $A = (A_{i,j}) \in \mathbb{R}^{n \times n}$, where $A_{i,j} \in \mathbb{R}^{n_i \times n_j}$, $i, j = 1, 2, \dots, \ell$, and $n_i (i = 1, 2, \dots, \ell)$ are positive integers satisfying $n_1 + n_2 + \dots + n_\ell = n$.

For the structured preconditioners based on the relaxation iteration methods involving parameters, we can further optimize them through choices of the optimal parameters. In addition, we should point out that, although all results in this paper are demonstrated in $\|\cdot\|_2$ -norm, they trivially hold for other consistent matrix norms such as $\|\cdot\|_1$ -norm and $\|\cdot\|_\infty$ -norm.

Acknowledgments: The author is very much indebted to the referees for their constructive and valuable comments and suggestions which greatly improved the original version of this paper.

References

- [1] H. Ammari, G. Bao and A.W. Wood, An integral equation method for the electromagnetic scattering from cavities, *Math. Methods Appl. Sci.*, 23(2000), 1057-1072.
- [2] O. Axelsson, Iterative Solution Methods, *Cambridge University Press*, Cambridge, 1994.
- [3] Z.-Z. Bai, Parallel Iterative Methods for Large-Scale Systems of Algebraic Equations, *Ph.D. Thesis of Shanghai University of Science and Technology*, Shanghai, June 1993. (In Chinese)
- [4] Z.-Z. Bai, A class of hybrid algebraic multilevel preconditioning methods, *Appl. Numer. Math.*, 19(1996), 389-399.
- [5] Z.-Z. Bai, Parallel hybrid algebraic multilevel iterative methods, *Linear Algebra Appl.*, 267(1997), 281-315.
- [6] Z.-Z. Bai, A class of modified block SSOR preconditioners for symmetric positive definite systems of linear equations, *Adv. Comput. Math.*, 10(1999), 169-186.
- [7] Z.-Z. Bai, Modified block SSOR preconditioners for symmetric positive definite linear systems, *Ann. Operations Research*, 103(2001), 263-282.
- [8] Z.-Z. Bai, I.S. Duff and A.J. Wathen, A class of incomplete orthogonal factorization methods. I: methods and theories, *BIT*, 41(2001), 53-70.
- [9] Z.-Z. Bai and G.-Q. Li, Restrictively preconditioned conjugate gradient methods for systems of linear equations, *IMA J. Numer. Anal.*, 23(2003), 561-580.
- [10] Z.-Z. Bai and M.K. Ng, On inexact preconditioners for nonsymmetric matrices, *SIAM J. Sci. Comput.*, (2005), in press.
- [11] Z.-Z. Bai and Z.-Q. Wang, Restrictive preconditioners for conjugate gradient methods for symmetric positive definite linear systems, *J. Comput. Appl. Math.*, (2005), in press.
- [12] Z.-Z. Bai and D.-R. Wang, A class of new hybrid algebraic multilevel preconditioning methods, *Linear Algebra Appl.*, 260(1997), 223-255.

- [13] J.T. Betts, Practical Methods for Optimal Control Using Nonlinear Programming, *SIAM*, Philadelphia, PA, 2001.
- [14] A. Björck, Numerical Methods for Least Squares Problems, *SIAM*, Philadelphia, PA, 1996.
- [15] J.H. Bramble, J.E. Pasciak and A.T. Vassilev, Analysis of the inexact Uzawa algorithm for saddle point problems, *SIAM J. Numer. Anal.*, 34(1997), 1072-1092.
- [16] F. Brezzi and M. Fortin, Mixed and Hybrid Finite Element Methods, *Springer-Verlag*, New York etc., 1991.
- [17] I.S. Duff, N.I.M. Gould, J.K. Reid, J.A. Scott and K. Turner, The factorization of sparse symmetric indefinite matrices, *IMA J. Numer. Anal.*, 11(1991), 181-204.
- [18] I.S. Duff and J.K. Reid, Exploiting zeros on the diagonal in the direct solution of indefinite sparse symmetric linear systems, *ACM Trans. Math. Software*, 22(1996), 227-257.
- [19] N. Dyn and W.E. Ferguson, Jr., The numerical solution of equality constrained quadratic programming problems, *Math. Comput.*, 41(1983), 165-170.
- [20] S.C. Eisenstat, H.C. Elman and M.H. Schultz, Variational iterative methods for nonsymmetric systems of linear equations, *SIAM J. Numer. Anal.*, 20(1983), 345-357.
- [21] H.C. Elman, Preconditioners for saddle point problems arising in computational fluid dynamics, *Appl. Numer. Math.*, 43(2002), 75-89.
- [22] H.C. Elman and G.H. Golub, Inexact and preconditioned Uzawa algorithms for saddle point problems, *SIAM J. Numer. Anal.*, 31(1994), 1645-1661.
- [23] H.C. Elman, D.J. Silvester and A.J. Wathen, Performance and analysis of saddle point preconditioners for the discrete steady-state Navier-Stokes equations, *Numer. Math.*, 90(2002), 665-688.
- [24] B. Fischer, R. Ramage, D.J. Silvester and A.J. Wathen, Minimum residual methods for augmented systems, *BIT*, 38(1998), 527-543.
- [25] P.E. Gill, W. Murray and M.H. Wright, Practical Optimization, *Academic Press*, New York, NY, 1981.
- [26] R. Glowinski, Numerical Methods for Nonlinear Variational Problems, *Springer-Verlag*, New York etc., 1984.
- [27] G.H. Golub and C.F. Van Loan, Matrix Computations, 3rd Edition, *The Johns Hopkins University Press*, Baltimore and London, 1996.
- [28] G.H. Golub and A.J. Wathen, An iteration for indefinite systems and its application to the Navier-Stokes equations, *SIAM J. Sci. Comput.*, 19(1998), 530-539.
- [29] G.H. Golub, X. Wu and J.-Y. Yuan, SOR-like methods for augmented systems, *BIT*, 41(2001), 71-85.

- [30] N.I.M. Gould, M.E. Hribar and J. Nocedal, On the solution of equality constrained quadratic programming problems arising in optimization, *SIAM J. Sci. Comput.*, 23(2001), 1375-1394.
- [31] E. Haber, U.M. Ascher and D. Oldenburg, On optimization techniques for solving nonlinear inverse problems, *Inverse Problems*, 16(2000), 1263-1280.
- [32] A. Hadjidimos, Accelerated overrelaxation method, *Math. Comput.*, 32(1978), 149-157.
- [33] J. Jin, The Finite Element Method in Electromagnetics, *John Wiley & Sons*, New York, NY, 1993.
- [34] C. Keller, N.I.M. Gould and A.J. Wathen, Constrained preconditioning for indefinite linear systems, *SIAM J. Matrix Anal. Appl.*, 21(2000), 1300-1317.
- [35] A. Klawonn, Block-triangular preconditioners for saddle point problems with a penalty term, *SIAM J. Sci. Comput.*, 19(1998), 172-184.
- [36] P.A. Markowich, The Stationary Semiconductor Device Equations, *Springer-Verlag*, New York, 1986.
- [37] M.F. Murphy, G.H. Golub and A.J. Wathen, A note on preconditioning for indefinite linear systems, *SIAM J. Sci. Comput.*, 21(2000), 1969-1972.
- [38] I. Perugia and V. Simoncini, Block-diagonal and indefinite symmetric preconditioners for mixed finite element formulations, *Numer. Linear Algebra Appl.*, 7(2000), 585-616.
- [39] R.J. Plemmons, A parallel block iterative method applied to computations in structural analysis, *SIAM J. Alg. Disc. Meth.*, 7(1986), 337-347.
- [40] Y. Saad, Iterative Methods for Sparse Linear Systems, 2nd Edition, *SIAM*, Philadelphia, PA, 2003.
- [41] Y. Saad and M.H. Schultz, GMRES: a generalized minimal residual algorithm for solving nonsymmetric linear systems, *SIAM J. Sci. Stat. Comput.*, 7(1986), 856-869.
- [42] G.E. Sartoris, A 3D rectangular mixed finite element method to solve the stationary semiconductor equations, *SIAM J. Sci. Stat. Comput.*, 19(1998), 387-403.
- [43] S. Selberherr, Analysis and Simulation of Semiconductor Devices, *Springer-Verlag*, New York, 1984.
- [44] G. Strang, Introduction to Applied Mathematics, *Wellesley-Cambridge Press*, MA, 1986.
- [45] W. Zulenher, Analysis of iterative methods for saddle point problems: a unified approach, *Math. Comput.*, 71(2001), 479-505.