

Exponential Dichotomies and Homoclinic Orbits from Heteroclinic Cycles*

Tiejun Chen, Yu Xiang, Weiwei Cai, Yuxiao Chen

Yiyang Medical College Hunan Pro of China, Yiyang, China
Email: cwwlove@sina.com

Received February 13, 2012; revised April 20, 2012; accepted April 28, 2012

ABSTRACT

In this paper, we investigate the homoclinic bifurcations from a heteroclinic cycle by using exponential dichotomies. We give a Melnikov—type condition assuring the existence of homoclinic orbits from heteroclinic cycle. We improve some important results.

Keywords: Exponential Dichotomies; Homoclinic Orbits; Heteroclinic Cycle; Melnikov Function

1. Introduction

We consider the n -dimensional differential equations

$$\dot{x} = f(x, v, \varepsilon) \quad (1.1)$$

where $x \in R^n$, ε is a small parameter, $v \in R^2$ is a parameter. In studying the global bifurcation, we usually assume unperturbed differential equations

$$\dot{x} = f(x, 0, 0) \quad (1.2)$$

admits a hyperbolic equilibrium and a homoclinic orbit connecting it. It is the persistence of homoclinic orbit and heteroclinic that we usually study in global bifurcation, we refer to Wiggins [1], Palmer [2,3], Naudot [4] and Meyer and Sell [5]. But in studying the pulse solutions of some reaction-diffusion equations, we often meet the problem of homoclinic bifurcations from the heteroclinic cycles, refer to Kokubu [6], Chow, Deng and Terman [7], Gambaudo [8] and reference therein. Suppose Equation (1.2) has two hyperbolic equilibria p_1, p_2 and two homoclinic orbits q_1, q_2 and two homoclinic orbits $q_1(t), q_2(t)$.

If

$$\lim_{t \rightarrow -\infty} q_i(t) = p_i, \lim_{t \rightarrow +\infty} q_i(t) = p_{i+1}, i = 1, 2$$

(where we assume $q_1(t) \rightarrow q_3(t) \rightarrow p_1 = p_3$) then we say that $\Gamma = q_1(t) \cup q_2(t) \cup q_1 \cup q_2$ is a heteroclinic cycle consisting of $q_1(t), q_2(t), p_1$ and p_2 . The study of homoclinic bifurcation from a heteroclinic cycle is very important and interest not only from the point of view of

bifurcation theory itself but also from the point of view of application, we refer to Kokubu [6], Chow, Deng and Terman [7]. The main purpose of this paper is to investigate the homoclinic bifurcation from heteroclinic cycles by making use of exponential dichotomies and Melnikov technique. For convenience, we only discuss the case of heteroclinic cycles with length = 2. Using the theory of exponential dichotomies, Melnikov functions and Slinnikov change of variable, Kokubu [6] investigate the periodic and homoclinic bifurcations from a heteroclinic cycle. In Kokubu [6], he needs to divide the problem into critical and non-critical two cases. Moreover, he needs that the heteroclinic orbits approach the hyperbolic equilibria along the eigenspaces associated with the principal eigenvalues. Chow, Deng and Terman [7] also studied the same problem in the non-critical case by making use of Liapunov-schmidt method and Silnikov's changes of variable and Poincaré map and obtain some analytical results. Chow, Deng and Terman [7] also the conditions as in Kokubu [6]. Melnikov functions were not obtained in Chow, Deng and Terman [7]. Chow, Deng and Terman [9] studied the same problem as this paper, Kokubu [7] did not need to divide the problem into critical and non-critical two cases and unified the two cases and didn't need that the heteroclinic orbits approach the hyperbolic equilibria along the eigenspaces associated with the principal eigenvalues. The results of Chow, Deng and Terman [9] are weaker than those of Kokubu [6] and Chow, Deng and Terman [7] under weaker assumptions because of the topological approaches. The purpose of this paper is to improve the above results by an analytic method (Lin's method [10]). We can also unify the critical and non-critical cases and weak the condi-

*This work is supported by Program for Yiyang Municipal Bureau of Science and Technology of Hunan Pro in China (Ministry YKZ-2011-30)

tions of Kokubu [6], Chow, Deng and Terman [7,9]. Moreover, it is also an interesting to provide an analytic method of studying bifurcations of heteroclinic cycles. Many ideas of this paper come from Lin [10], Meyer and Sell [5], Kokubu [6] and Palmer [2,3]. But it should note that the results of this paper cannot be followed directly from these papers, much technique has been made. Let us finally mention the related results on the bifurcations of heteroclinic cycles. Sandstede [11] investigated the forced symmetry breaking of heteroclinic cycles. Guckenheimer and Holmes [12] discussed the spontaneous symmetry breaking of heteroclinic cycle. Krupa and Melbourne [13] studied the stability of heteroclinic cycle. On the other related results on heteroclinic cycles, we refer to the references of the above mentioned papers and good survey of Krupa [14]. The paper is organized as following. In Section 2, we give the main result; in Section 3, the proof of the main result is given.

The main tool used in this paper is theory of exponential dichotomies. We consider the linear differential equations

$$\dot{x} = A(t)x \tag{1.3}$$

where $x \in R^n$, $A(t)$ is a $n \times n$ continuous bounded matrix on R . We say Equation (1.3) admits an exponential dichotomy on interval J if there exist constants K, α , a projection P and the fundamental matrix $X(t)$ of Equation (1.3) satisfying;

$$\begin{aligned} |X(t)PX^{-1}(s)| &\leq Ke^{-\alpha(t-s)} t \geq s \\ |X(t)(I-P)X^{-1}(s)| &\leq Ke^{-\alpha(t-s)} \geq s \end{aligned}$$

for $t, s \in J$. On the theory of exponential dichotomies, refer to Coppel [15], Sacker and Sell [16] and Meyer and Sell [17]. On the relations between exponential dichotomies and homoclinic, heteroclinic bifurcations, we refer to Palmer [18] and Meyer and Sell [16].

2. Main Result

We consider differential equations

$$\dot{x} = f(x, v, \varepsilon) \tag{2.1}$$

where $x \in R^n, \varepsilon$ is small parameter, $v \in R^2$ is a parameter. $f : \Omega \times I_1^2 \times I_2 \rightarrow R^n$ is C_b^2 with respect to $(x, v, \varepsilon) \in \Omega \times I_1^2 \times I_2$, where $\Omega \subset R^2$ a compact subset, I_1 a small interval containing zero, $I_2 = [0, b]$ a small interval.

We assume C1. For $v = 0, \varepsilon = 0$, unperturbed equation

$$\dot{x} = f(x, 0, 0) \tag{2.2}$$

Admits two hyperbolic equilibriums p_1, p_2 and two heteroclinic orbits $q_1(t), q_2(t)$ connecting p_1, p_2 respectively (form a heteroclinic cycle), that is,

$$\lim_{t \rightarrow -\infty} q_2(t) = p_1, \lim_{t \rightarrow \infty} q_1(t) = p_2,$$

$$\lim_{t \rightarrow -\infty} q_1(t) = p_2, \lim_{t \rightarrow \infty} q_2(t) = p_1,$$

$$q_i(t) \in \Omega, i = 1, 2.$$

We denote the heteroclinic cycle by

$$\Gamma = q_1(t) \cup q_2(t) \cup q_1 \cup q_2.$$

We want to study under what conditions can a homoclinic orbit bifurcate from the heteroclinic cycle Γ as the second case of Kokubu [6]

C2. All real parts of the matrix $f_x(p_i, 0, 0) (i = 1, 2)$ are different from zero; and the number of eigenvalues with positive real parts is $m_i = m (< n) (i = 1, 2)$.

If the conditions C1 and C2 are satisfied then equation

$$\dot{x} = f_x(p_i, 0, 0)x, i = 1, 2. \tag{2.3}$$

admit an exponential dichotomy on both R_+ and R_- , and the sum of dimensions of stable and unstable subspaces is n . It follows from the roughness of exponential dichotomy that (refer to Zeng [12], Sacker and Sell [16], Coppel [15]) that the variational equations along $q_i(t)$

$$\dot{x} = f_x(q_i(t), 0, 0), i = 1, 2 \tag{2.4}$$

admit an exponential dichotomy on both R_+ and R_- , and the sum of dimensions of the stable and unstable subspaces is $m_1 + n - m_2 = n$. In the follows, because we want to the exponent of $e^{\alpha \ln \varepsilon} = \varepsilon^\alpha$ to be greater than 1, without loss of generality, we may assume the constants

$K, \alpha > 1$. Otherwise, we replace $l_n \varepsilon$ by $\frac{\alpha}{2} l_n \varepsilon$, then the exponent of $e^{\frac{\alpha^2}{2} \ln \varepsilon} = \varepsilon^2$ is greater than 1.

C3. The variational Equations (2.4) admit a unique (up to a scalar multiple) nontrivial bounded solution $\phi_i(t)$ on R .

Under the conditions C1, C2, C3, we can prove (refer to Zeng [12]) that the adjoint equations of equations of (2.3), (2.4)

$$\dot{\xi} = -g_x^*(q_i(t))\xi \quad i = 1, 2$$

also admit unique (up to a scalar multiple) nontrivial bounded solution $\psi_1(t), \psi_2(t)$, respectively, on R , and an exponential dichotomy on both R_+ and R_- , respectively. The constants of the exponential dichotomies are also K, α .

We let

$$M_1 = \int_{-\infty}^{+\infty} \psi_2^*(t) f_v(q_1(t), 0, 0) dt,$$

$$M_2 = \int_{-\infty}^{+\infty} \psi_2^*(t) f_v(q_2(t), 0, 0) dt.$$

The main result of this paper is

Theorem 1 We assume the conditions C2, C2 and C3 are satisfied, then when ε, ν sufficiently small Equation (2.1) admits a unique hyperbolic equilibrium $p_2(\varepsilon, \nu)$ satisfying $p_1(0, 0) = p_1$. If the 2×2 matrix

$$M = \begin{pmatrix} M_1 & \\ & M_2 \end{pmatrix}$$

is invertible, the for $\varepsilon > 0$ sufficiently small there exists a continuous function $\nu = \nu(\varepsilon)$ satisfying

$$\mu(0) = \mu_0 = -M^{-1} \left(\begin{array}{c} \int_{-\infty}^{+\infty} \psi_1^*(t) f_\nu(q_1(t), 0, 0) ds \\ \int_{-\infty}^{+\infty} \psi_2^*(t) f_\nu(q_2(t), 0, 0) ds \end{array} \right)$$

such that the equation

$$\dot{x} = f(t, \varepsilon \nu(\varepsilon), \varepsilon) \tag{2.6}$$

admits a homoclinic orbit connecting $p_1(\varepsilon, \varepsilon \nu(\varepsilon))$ in the neighbourhood of the heteroclinic cycle Γ .

Remark If the conditions C1, C2 and C3 are satisfied, using the standard method (refer to Zeng [19]), we can obtain the bifurcative equations of persistence of the two heteroclinic orbits $q_1(t)$ and $q_2(t)$

$$\tilde{H}_1(\mu, \varepsilon) = \int_{-\infty}^{+\infty} \psi_1^*(s) g_1(s, \tilde{z}_1(s, \mu, \varepsilon), \mu, \varepsilon) ds = 0 \tag{2.7}$$

$$\tilde{H}_2(\mu, \varepsilon) = \int_{-\infty}^{+\infty} \psi_2^*(s) g_2(s, \tilde{z}_2(s, \mu, \varepsilon), \mu, \varepsilon) ds = 0 \tag{2.8}$$

where $\tilde{z}_i(s, 0, 0) = 0, i = 1, 2$. If the matrix M is invertible then we can easily prove (refer to Zeng [19]) that for $\varepsilon \neq 0$ sufficiently small there exists a continuously differentiable function $\mu = \tilde{\mu}(\varepsilon)$ such that

$$\tilde{H}(\varepsilon \tilde{\mu}(\varepsilon), \varepsilon) = 0 \quad \tilde{H}_2(\varepsilon \tilde{\mu}(\varepsilon), \varepsilon) = 0$$

and

$$\dot{x} = f(x, \varepsilon \tilde{\mu}(\varepsilon), \varepsilon) \tag{2.9}$$

has two hyperbolic equilibriums $\tilde{p}_1(\varepsilon), \tilde{p}_2(\varepsilon)$, satisfying $\tilde{p}_1(0) = p_1$ and $\tilde{p}_2(0) = p_2$, and two heteroclinic orbits $\tilde{q}_1(t, \varepsilon), \tilde{p}_2(t, \varepsilon)$ satisfying

$$\lim_{t \rightarrow -\infty} \tilde{q}_1(t, \varepsilon) = \tilde{p}_1(\varepsilon) \quad \lim_{t \rightarrow -\infty} \tilde{q}_1(t, \varepsilon) = \tilde{p}_2(\varepsilon),$$

$$\lim_{t \rightarrow -\infty} \tilde{q}_2(t, \varepsilon) = \tilde{p}_2(\varepsilon) \quad \lim_{t \rightarrow -\infty} \tilde{q}_2(t, \varepsilon) = \tilde{p}_1(\varepsilon).$$

That is, the heteroclinic cycle Γ persists in the region of parameters

$$\{(\varepsilon \tilde{\mu}(\varepsilon), \varepsilon)\}$$

From Theorem 1 of this paper we see that in the region of parameters

$$\{(\varepsilon \mu(\varepsilon), \varepsilon)\}$$

a homoclinic orbit connecting $p_1(\varepsilon, \varepsilon \nu(\varepsilon))$ bifurcates from the heteroclinic cycle Γ .

Kokubu [5] proved that

$$\partial \{(\varepsilon \mu(\varepsilon), \varepsilon)\} = \{(\varepsilon \tilde{\mu}(\varepsilon), \varepsilon)\}$$

We can also prove that if the conditions C1, C2 and C3 are satisfied then for $\varepsilon \neq 0$ sufficiently small a homoclinic orbit connecting $p_2(\varepsilon), P_2(0) = p_2$, bifurcates from the heteroclinic cycle Γ , but the region of parameters of bifurcation is different from $\{(\varepsilon \mu(\varepsilon), \varepsilon)\}$.

3. The Proof of the Main Result

To prove the main result of this paper, we want to find the bounded solutions of Equation (2.1) $x_1(t)$ on $(-\infty, \omega)$ and $x_2(t)$ on $(-\omega, \infty)$ satisfying

$$x_1(\omega) = x_2(-\omega)$$

We make a change of variables for Equation (2.1)

$$x = z_1 + q_1(t), -\infty \leq t < \omega.$$

$$x = z_2 + q_2(t), -\omega < t \leq \infty.$$

respectively, and obtain the equations

$$z_1 = f(z_1 + q_1(t), \nu, \varepsilon) - f(q_1(t), 0, 0), -\infty < t \leq \omega.$$

$$z_2 = f(z_2 + q_2(t), \nu, \varepsilon) - f(q_2(t), 0, 0), -\omega \leq t < \infty.$$

We write the above equations in the following form

$$z_1 = A_1(t) z_1 + g_1(t, z_1, \nu, \varepsilon), -\infty < t \leq \omega. \tag{3.1}$$

$$z_2 = A_2(t) z_2 + g_2(t, z_2, \nu, \varepsilon), -\omega \leq t < \infty. \tag{3.2}$$

And the boundary value condition in the following form

$$z_1(\omega) - z_2(-\omega) = q_2(-\omega) - q_1(\omega) \tag{3.3}$$

where ω is sufficiently large.

$$A_i(t) = f_x(q_i(t), 0, 0) \tag{3.4}$$

$$g_i(t, z, \nu, \varepsilon) = f(z_1 + q_i(t), \nu, \varepsilon) - f(q_i(t), 0, 0) - A_i(t) z_i$$

$i = 1, 2$. $g_i(t, z_i, \nu, \varepsilon) = i = 1, 2$ satisfying:

$$|g_i(t, z_i, \nu, \varepsilon)| \leq C_1 (|z_i|^2 + |\nu| + |\varepsilon|) \quad i = 1, 2 \tag{3.5}$$

In order to find the bounded solutions of Equations (3.1), (3.2) and (3.3), we consider the following boundary value problem

$$z_1 = A_1(t) z_1 + g_1(t, z_1, \nu, \varepsilon), -\infty < t \leq \ln \varepsilon. \tag{3.6}$$

$$z_2 = A_2(t) z_2 + g_2(t, z_2, \nu, \varepsilon), \ln \varepsilon \leq t < \infty. \tag{3.7}$$

$$z_1(-\ln \varepsilon) - z_2(\ln \varepsilon) = q_2(\ln \varepsilon) - q_1(-\ln \varepsilon). \tag{3.8}$$

where $\varepsilon < 1$. For any $h_1(t), h_2(t) \in C_b^2(R, R^n)$, we first consider the following boundary value problems for $\varepsilon > 0$

$$z_1 = A_1(t)z_1 + h_1(t), -\infty < t \leq -\ln \varepsilon. \tag{3.9}$$

$$z_2 = A_2(t)z_2 + h_2(t), \ln \varepsilon \leq t < \infty \tag{3.10}$$

$$z_1(-\ln \varepsilon) - z_2(\ln \varepsilon) = q_2(\ln \varepsilon) - q_1(-\ln \varepsilon) \tag{3.11}$$

We let $b(\varepsilon) = q_2(\ln \varepsilon) - q_1(-\ln \varepsilon)$ and have the following lemma:

Lemma 1 Assume the conditions C1, C2 and C3 are satisfied.

Then there exists sufficiently small $\varepsilon_0 > 0$ such that for $0 < \varepsilon \leq \varepsilon_0$ Equations (3.9), (3.10) and (3.11) admit a unique continuous except at $t = 0$ bounded solution $z_i(t, \varepsilon)$ satisfying $\phi_i^*(0)z_i(0-, \varepsilon) = 0, i = 1, 2$ with

$$|z_i(t, \varepsilon)| \leq C_2(|b(\varepsilon)| + \|h_i(t)\|), i = 1, 2. \tag{3.12}$$

Moreover, $z_i(t, \varepsilon)$ is differentiable in ε and with

$$|z_{1\varepsilon}(t, \varepsilon)| + |z_{2\varepsilon}(t, \varepsilon)| \leq L.$$

where $z_i(0-)$ denotes the left limit of function $z_i(t, \varepsilon)$ at $t = 0$, C_1 is a constant independent of ε . Moreover, if

$$Z_1(\ln \varepsilon) - Z_2(\ln \varepsilon) = \{A_1(-\ln \varepsilon)z_1(-\ln \varepsilon, \varepsilon) + A_2(\ln \varepsilon)z_2(\ln \varepsilon, \varepsilon) + A_1(-\ln \varepsilon)q_1(-\ln \varepsilon) + A_2(\ln \varepsilon)q_2(\ln \varepsilon)\} / \varepsilon$$

Let

$$\begin{aligned} \rho_1(t) &= z_1(t, \varepsilon + h) - z_1(t, \varepsilon) - Z_1(t, \varepsilon)h, \\ \rho_2(t) &= z_2(t, \varepsilon + h) - z_2(t, \varepsilon) - Z_2(t, \varepsilon)h. \end{aligned}$$

then $\rho_1(t), \rho_2(t)$ are the solutions of equations

$$\begin{aligned} \rho_1 &= A_1(t)\rho_1, -\infty < t \leq -\ln \varepsilon. \\ \rho_2 &= A_2(t)\rho_2, \ln \varepsilon \leq t < \infty. \\ \rho_1(-\ln \varepsilon) - \rho_2(\ln \varepsilon) &= z_1(-\ln \varepsilon, \varepsilon + h) \\ &\quad - z_2(\ln \varepsilon, \varepsilon + h) - z_1(-\ln \varepsilon, \varepsilon) \\ &\quad + z_2(\ln \varepsilon, \varepsilon) - Z_1(-\ln \varepsilon, \varepsilon)h + Z_2(\ln \varepsilon, \varepsilon)h. \end{aligned}$$

In the same method as follows, we can show that

$$|\eta_1(t)| + |\eta_2(t)| = O(|h|)$$

$$|\eta_1(t, \varepsilon)| + |\eta_2(t, \varepsilon)| \leq 2C_1|z_1(-\ln \varepsilon, \varepsilon + h) - z_2(\ln \varepsilon, \varepsilon + h) - z_1(-\ln \varepsilon, \varepsilon) + z_2(\ln \varepsilon, \varepsilon)|$$

hence there exists a constant $L > 0$ such that

$$|z_{1\varepsilon}(t, \varepsilon)| + |z_{2\varepsilon}(t, \varepsilon)| \leq L.$$

This completes the proof of Lemma 2.

Now we consider Equations (3.1)-(3.3). We have the following lemma:

Lemma2 Assume conditions C1, C2 and C3 are satisfied. Then there exist sufficiently small $\varepsilon_0 > 0$ and the

$$\int_{-\infty}^{-\ln \varepsilon} \psi_1^*(s)h_1(s)ds - \psi_1^*(-\ln \varepsilon)z_1(-\ln \varepsilon) = 0. \tag{3.13}$$

$$\int_{\ln \varepsilon}^{-\ln \varepsilon} \psi_2^*(s)h_2(s)ds - \psi_2^*(-\ln \varepsilon)z_2(-\ln \varepsilon) = 0. \tag{3.14}$$

then $z_1(t), z_2(t)$ are continuous at $t = 0$.

Proof Lemma 2 is mainly due to Lin [10]. For the proof of the first part of existences of the solutions $z_i(t, \varepsilon)$ satisfying (3.12), (3.13) and (3.14), we refer to Lin [10] and omit the proof. We now want to prove the second part that $z_i(t, \varepsilon)$ is differentiable in ε . We let $z_1(t, \varepsilon), z_2(t, \varepsilon)$ be the bounded solutions, which are continuous except at $t = 0$ and satisfy (3.12), (3.13) and (3.14), of equations

$$\begin{aligned} z_1 &= A_1(t)z_1 + h_1(t), -\infty < t \leq -\ln \varepsilon. \\ z_2 &= A_2(t)z_2 + h_2(t), \ln \varepsilon \leq t < \infty \\ z_1(-\ln \varepsilon) - z_2(\ln \varepsilon) &= q_2(\ln \varepsilon) - q_1(-\ln \varepsilon) \end{aligned}$$

Let $Z_1(t, \varepsilon), Z_2(t, \varepsilon)$ be the bounded solutions of equations

$$\begin{aligned} Z_1 &= A_1(t)Z_1, -\infty < t \leq -\ln \varepsilon. \\ Z_2 &= A_2(t)Z_2, \ln \varepsilon \leq t < \infty. \end{aligned}$$

$$z_{i\varepsilon}(t, \varepsilon) = Z_i(t, \varepsilon), i = 1, 2.$$

Now we prove the boundness of $z_{i\varepsilon}$. Let

$$\begin{aligned} \eta_1(t, \varepsilon) &= z_1(t, \varepsilon + h) - z_1(t, \varepsilon), \\ \eta_2(t, \varepsilon) &= z_2(t, \varepsilon + h) - z_2(t, \varepsilon), \end{aligned}$$

then $\eta_1(t, \varepsilon), \eta_2(t, \varepsilon)$ are the solutions of equations

$$\begin{aligned} \eta_1 &= A_1(t)\eta_1, -\infty < t \leq -\ln \varepsilon. \\ \eta_2 &= A_2(t)\eta_2, -\infty < t \leq -\ln \varepsilon. \\ \eta_1(-\ln \varepsilon, \varepsilon) - \eta_2(\ln \varepsilon, \varepsilon) &= z_1(-\ln \varepsilon, \varepsilon + h) - z_2(\ln \varepsilon, \varepsilon + h) \\ &\quad - z_1(-\ln \varepsilon, \varepsilon) + z_2(\ln \varepsilon, \varepsilon) \end{aligned}$$

From (3.12) we obtain

constants $C_2, L > 0$ such that for $0 < \varepsilon \leq \varepsilon_0$ Equations (3.1)-(3.3) admit a unique continuous except at $t = 0$ bounded solution $z_i(t, v, \varepsilon)$ satisfying

$$\phi_i^*(0)z_i(0-, v, \varepsilon) = 0, i = 1, 2$$

with

$$\begin{aligned} |z_i(t, v, \varepsilon)| &\leq C_2(|b(\varepsilon)| + |v| + |\varepsilon|) \\ &\leq C_2(|2K\varepsilon^\alpha| + |v| + |\varepsilon|), i = 1, 2 \end{aligned}$$

$$|z_{iv}(t, v, \varepsilon)| + |z_{i\varepsilon}(t, v, \varepsilon)| \leq L, i = 1, 2. \tag{3.15}$$

Moreover, if

$$G_1(v, \varepsilon) = \int_{-\infty}^{-\ln \varepsilon} \psi_1^*(s) g_1(s, z_1(s, v, \varepsilon), v, \varepsilon) ds - \psi_1^*(-\ln \varepsilon) z_1(-\ln \varepsilon, v, \varepsilon) = 0 \tag{3.16}$$

$$G_1(v, \varepsilon) = \int_{-\infty}^{-\ln \varepsilon} \psi_1^*(s) g_2(s, z_2(s, v, \varepsilon), v, \varepsilon) ds - \psi_1^*(-\ln \varepsilon) z_2(-\ln \varepsilon, v, \varepsilon) = 0 \tag{3.17}$$

then $z_1(t, v, \varepsilon)$, $z_2(t, v, \varepsilon)$ are continuous at $t = 0$.

The proof of Lemma 2 can be proved by contract fixed point theorem and is similar to that of Lin [10].

From Lemma 2 we see that if we have proved that bifurcative Equations (3.16) and (3.17) can be can be solved then we find the continuously bounded solutions of Equations (3.1), (3.2) and (3.3)

$$z_1(t, v, \varepsilon) - \infty < t \leq -\ln \varepsilon$$

and

$$z_2(t, v, \varepsilon), \ln \varepsilon \leq t < \infty.$$

Now we mainly solve bifurcative Equations (3.16) and (3.17). We make a change of variable for Equations (3.16) and (3.17) $v \sim \varepsilon v$ and obtain the following bifurcative equation

$$B_1(v, \varepsilon) = G_1(v, \varepsilon) = \int_{-\infty}^{-\ln \varepsilon} \psi_1^*(s) g_1(s, z_1(s, v, \varepsilon), v, \varepsilon) ds - \psi_1^*(-\ln \varepsilon) z_1(-\ln \varepsilon, v, \varepsilon) = 0 \tag{3.18}$$

$$B_1(v, \varepsilon) = G_1(v, \varepsilon) = \int_{-\infty}^{-\ln \varepsilon} \psi_1^*(s) g_2(s, z_2(s, v, \varepsilon), v, \varepsilon) ds - \psi_1^*(-\ln \varepsilon) z_2(-\ln \varepsilon, v, \varepsilon) = 0 \tag{3.19}$$

From (3.15) we have

$$|z_i(t, v, \varepsilon)| \leq C_2 (2K |\varepsilon|^\alpha + |\varepsilon| |v| + |\varepsilon|) i = 1, 2 \tag{3.20}$$

Letting $\varepsilon \rightarrow 0$ in the above equation, we obtain

$$z_i(t, 0, 0) = 0 \quad i = 1, 2 \tag{3.21}$$

(Remark ACTUALLY, $z_i(t, \varepsilon v, \varepsilon)$ is defined only for $\varepsilon > 0$. but due to the existence of its limit, here we define the vaule of the limit to be the value at $\varepsilon = 0$. In the sequel, we make the same definition.)

From the property of $\psi_i(t)$ we have

$$|\psi_i(\ln \varepsilon)| \leq K e^{\alpha \ln \varepsilon} = K \varepsilon, i = 1, 2$$

hence

$$\lim_{t \rightarrow -\infty} \psi_i(-\ln \varepsilon) = 0, i = 1, 2 \tag{3.22}$$

From the representation of (3.18), (3.19), (3.21) and

(3.33) we obtain

$$B_1(v, 0) = \int_{-\infty}^{\infty} \psi_1^*(s) g_1(s, z_1(s, 0, 0), 0, 0) ds - \lim_{\varepsilon \rightarrow 0} \psi_1^*(-\ln \varepsilon) z_1(-\ln \varepsilon, \varepsilon v, \varepsilon) = \int_{-\infty}^{\infty} \psi_1^*(s) g_1(s, z_1(s, 0, 0, 0)) ds = 0 \tag{3.23}$$

In the same way, we can obtain

$$B_2(v, 0) = \int_{-\infty}^{\infty} \psi_2^*(s) g_2(s, z_1(s, 0, 0, 0)) ds = 0 \tag{3.24}$$

For convenience, we define a 2×1 matrix

$$B(v, \varepsilon) = \begin{pmatrix} B_1(v, \varepsilon) \\ B_2(v, \varepsilon) \end{pmatrix}$$

then we have

$$B(v, 0) = 0$$

We define

$$H(\varepsilon, v) = \begin{cases} \frac{B(v, \varepsilon)}{\varepsilon}, & \varepsilon \neq 0 \\ B_\varepsilon(v, \varepsilon), & \varepsilon = 0 \end{cases}$$

Obviously, for $\varepsilon \neq 0$ equation

$$B(\varepsilon, v) = 0 \tag{3.25}$$

And equation

$$H(\varepsilon, v) = 0 \tag{3.26}$$

equivalent. Now we want to find the solutions of Equation (3.26). We first compute $B_\varepsilon(v, 0)$. From (3.18) we have

$$B(\varepsilon, v) = -\psi_2(-\ln \varepsilon) \frac{1}{\varepsilon} g_1(-\ln \varepsilon, z_1(-\ln \varepsilon, \varepsilon v, \varepsilon), \varepsilon v, \varepsilon) + \int_{-\infty}^{\ln \varepsilon} \psi_1^*(s) \frac{d}{d\varepsilon} g_1(s, z_1(s, \varepsilon v, \varepsilon), \varepsilon v, \varepsilon) ds - \frac{d}{d\varepsilon} \{ \psi_1^*(-\ln \varepsilon) z_1(-\ln \varepsilon, \varepsilon v, \varepsilon) \} \tag{3.27}$$

Now we compute (3.27). Since

$$|g_i(t, z, \varepsilon v, \varepsilon)| \leq C_1 (|z_1|^2 + |\varepsilon| |v| + |\varepsilon|)$$

we have

$$\begin{aligned} & \left| \frac{1}{\varepsilon} g_1(-\ln \varepsilon, z_1(-\ln \varepsilon, \varepsilon v, \varepsilon), \varepsilon v, \varepsilon) \right| \\ & \leq \frac{1}{\varepsilon} C_1 (|z_1 - \ln \varepsilon, \varepsilon v, \varepsilon|^2 + |\varepsilon| |v| + |\varepsilon|) \\ & \leq \frac{1}{\varepsilon} C_2 \left[C_2 \left(2K e^{\alpha - \frac{1}{2}} + |\varepsilon| |v| + |\varepsilon| \right)^2 + |\varepsilon| |v| + |\varepsilon| \right] \\ & \leq C_2 \left[C_1 \left(2K e^{\alpha - \frac{1}{2}} + |\varepsilon|^{\frac{1}{2}} |v| + |\varepsilon|^{\frac{1}{2}} \right)^2 + |v| + 1 \right] \end{aligned}$$

and hence $\frac{1}{\varepsilon}g_1(-\ln \varepsilon, z_1(-\ln \varepsilon, \varepsilon v, \varepsilon), \varepsilon v, \varepsilon)$ is bounded for $\varepsilon \geq 0$.

Since

$$\lim_{\varepsilon \rightarrow 0} \psi_1^*(-\ln \varepsilon) + \psi_1^*(\infty) = 0,$$

we have

$$\lim_{\varepsilon \rightarrow 0} \psi_1^*(-\ln \varepsilon) \frac{1}{\varepsilon} g_1(-\ln \varepsilon), z_1(-\ln \varepsilon, \varepsilon v, \varepsilon), \varepsilon v, \varepsilon = 0 \tag{3.28}$$

Noting $z_1(t, 0, 0) = 0$, we can easily prove that

$$\lim_{\varepsilon \rightarrow 0} \frac{d}{d\varepsilon} g_1(s, z_1(-\ln \varepsilon, \varepsilon v, \varepsilon), \varepsilon v, \varepsilon) = f_v(q_1(s), 0, 0)v + f_\varepsilon(q_1(s), 0, 0),$$

hence

$$\lim_{\varepsilon \rightarrow 0} \int_{-\infty}^{-\ln \varepsilon} \psi_1^*(s) \frac{d}{d\varepsilon} g_1(s, z_1(-\ln \varepsilon, \varepsilon v, \varepsilon)) \varepsilon v, \varepsilon ds = \int_{-\infty}^{-\ln \varepsilon} \psi_1^*(s) f_v(q_1(s), 0, 0)v + f_\varepsilon(q_1(s), 0, 0) ds. \tag{3.29}$$

Last, since

$$\begin{aligned} & \frac{d}{d\varepsilon} \{ \psi_1^*(-\ln \varepsilon) z_1(-\ln \varepsilon, \varepsilon v, \varepsilon) \} \\ & \leq \left| \psi_1^*(-\ln \varepsilon) f_x(q_1(-\ln \varepsilon)) \frac{2}{\varepsilon} z_1(-\ln \varepsilon, \varepsilon v, \varepsilon) \right| \\ & + \left| \psi_1^*(-\ln \varepsilon) [f(z_1(-\ln \varepsilon, \varepsilon v, \varepsilon) + q_1(-\ln \varepsilon) \varepsilon v, \varepsilon) - f(q_1(-\ln \varepsilon), 0, 0)] \frac{2}{\varepsilon} \right| \\ & + \left| \varepsilon z_{1v}(-\ln \varepsilon, \varepsilon v, \varepsilon) \right| + \left| z_{1\varepsilon}(-\ln \varepsilon, \varepsilon v, \varepsilon) \right| \\ & \leq K \|z_1(t, \varepsilon v, \varepsilon)\| \varepsilon^{\alpha-1} + \left| \psi_1^*(-\ln \varepsilon) \right| \\ & \left[C_1 (|z_1(-\ln \varepsilon, \varepsilon v, \varepsilon)| + |\varepsilon| |v| + |\varepsilon|) \right] \frac{2}{\varepsilon} + L \\ & \leq K \|z_1(t, \varepsilon v, \varepsilon)\| \varepsilon^{\alpha-1} \\ & + \left| \psi_1^*(-\ln \varepsilon) \right| [C_1(C_1|v|+1) + 1 + |v|] + L \end{aligned} \tag{3.30}$$

we obtain

$$\lim_{\varepsilon \rightarrow 0} \frac{d}{d\varepsilon} \{ \psi_1^*(-\ln \varepsilon) z_1(-\ln \varepsilon, \varepsilon v, \varepsilon) \} = 0 \tag{3.31}$$

From (3.28), (3.29) and (3.31) we have

$$B_{1\varepsilon}(v, 0) = \int_{-\infty}^{+\infty} \psi_1^*(s) (f_v(q_1(s), 0, 0)v + f_\varepsilon(q_1(s), 0, 0)) ds \tag{3.32}$$

In the same way, we can prove

$$B_{2\varepsilon}(v, 0) = \int_{-\infty}^{+\infty} \psi_1^*(s) (f_v(q_2(s), 0, 0)v + f_\varepsilon(q_2(s), 0, 0)) ds \tag{3.33}$$

Hence we have

$$\begin{aligned} H(v, 0) &= B_\varepsilon(v, 0) = \begin{pmatrix} B_{1\varepsilon}(v, 0) \\ B_{2\varepsilon}(v, 0) \end{pmatrix} \\ &= \begin{pmatrix} \int_{-\infty}^{+\infty} \psi_1^*(s) (f_v(q_1(s), 0, 0)v + f_\varepsilon(q_1(s), 0, 0)) ds \\ \int_{-\infty}^{+\infty} \psi_1^*(s) (f_v(q_2(s), 0, 0)v + f_\varepsilon(q_2(s), 0, 0)) ds \end{pmatrix} \\ &= \begin{pmatrix} M_1 \\ M_2 \end{pmatrix} v + \begin{pmatrix} \int_{-\infty}^{+\infty} \psi_1^*(s) (f_v(q_1(s), 0, 0)) ds \\ \int_{-\infty}^{+\infty} \psi_2^*(s) (f_v(q_2(s), 0, 0)) ds \end{pmatrix} \end{aligned}$$

Let

$$v_0 = -M^{-1} \begin{pmatrix} \int_{-\infty}^{+\infty} \psi_1^*(s) (f_\varepsilon(q_1(s), 0, 0)) ds \\ \int_{-\infty}^{+\infty} \psi_2^*(s) (f_\varepsilon(q_2(s), 0, 0)) ds \end{pmatrix}$$

then we have

$$H_v(v_0, 0) = B_\varepsilon(v_0, 0) = 0 \tag{3.35}$$

From (3.34) we have

$$H_v(v_0, 0) = M \tag{3.36}$$

Since the matrix M is invertible, it follows from the implicit function theorem that for $\varepsilon > 0$ sufficiently small there exists a continuous function $v = v(\varepsilon)$, $v(0) = v_0$ satisfying

$$H(v(\varepsilon), \varepsilon) = 0$$

Hence for $\varepsilon > 0$ sufficiently small we have

$$B(v(\varepsilon), \varepsilon) = 0 \tag{3.37}$$

Hence for $\varepsilon > 0$ sufficiently small Equations (3.6), (3.7) and (3.8)

$$\begin{aligned} z_1(t, \varepsilon) &= z_1(t, \varepsilon v(\varepsilon), \varepsilon), -\infty < t \leq -\ln \varepsilon, \\ z_1(t, \varepsilon) &= z_1(t, \varepsilon v(\varepsilon), \varepsilon), \ln \varepsilon < t \leq \infty, \end{aligned}$$

So for $\varepsilon > 0$ sufficiently small the equation

$$\dot{x} = f(t, \varepsilon v, (\varepsilon), \varepsilon). \tag{3.38}$$

has two solutions

$$\begin{aligned} x_1(t, \varepsilon) &= z_1(t, \varepsilon) + q_1(t), -\infty < t \leq -\ln \varepsilon, \\ x_2(t, \varepsilon) &= z_2(t, \varepsilon) + q_2(t), \ln \varepsilon < t \leq \infty, \end{aligned}$$

satisfying

$$x_1(-\ln \varepsilon, \varepsilon) = x_2(-\ln \varepsilon, \varepsilon)$$

We construct a solution of Equation (3.38) by making use of $x_1(t, \varepsilon)$ and $x_2(t, \varepsilon)$

$$x(t, \varepsilon) = \begin{cases} x_1(t - \ln \varepsilon, \varepsilon), & -\infty < t \leq 0 \\ x_2(t - \ln \varepsilon, \varepsilon), & 0 < t \leq \infty \end{cases}$$

Since $x_1(-\ln \varepsilon, \varepsilon) = x_2(-\ln \varepsilon, \varepsilon)$, $x(t, \varepsilon)$ is a continuously bounded solution of Equation (3.38).

Now we show $x(t, \varepsilon)$ is a homoclinic orbit connecting the equilibrium $q_1(\varepsilon, \varepsilon v(\varepsilon))$. Since when $-\infty < t \leq 0$

$$\begin{aligned} & |x_1(t, \varepsilon) - q_1(\varepsilon, \varepsilon v(\varepsilon))| \\ &= |z_1(t - \ln \varepsilon, \varepsilon v(\varepsilon), \varepsilon) + q_1(t - \ln \varepsilon) - q_1(\varepsilon, \varepsilon v(\varepsilon))| \\ &\leq |z_1(t - \ln \varepsilon, \varepsilon v(\varepsilon), \varepsilon)| + |q_1(t - \ln \varepsilon) - q_1| \\ &+ |q_1(\varepsilon, \varepsilon v(\varepsilon)) - q_1| \end{aligned} \tag{3.40}$$

Hence for any $\delta > 0$, there exist $\varepsilon_0 > 0$ and $T > 0$ such that when $0 < \varepsilon \leq \varepsilon_0$ and $t < -T + \ln \varepsilon$, we have

$$|x_1(t, \varepsilon) - q_1(\varepsilon, \varepsilon v(\varepsilon))| \leq \delta$$

Since $q_1(\varepsilon, \varepsilon v(\varepsilon))$ is hyperbolic, we obtain (refer to [9]) for $\varepsilon \neq 0$ sufficiently small

$$\lim_{t \rightarrow -\infty} x(t, \varepsilon) = q_1(\varepsilon, \varepsilon v(\varepsilon))$$

In the same way, we can prove that

$$\lim_{t \rightarrow \infty} x(t, \varepsilon) = q_1(\varepsilon, \varepsilon v(\varepsilon))$$

Hence $x(t, \varepsilon)$ is a homoclinic orbit connecting $q_1(\varepsilon, \varepsilon v(\varepsilon))$ in the neighbourhood of the heteroclinic cycle Γ .

Theorem 1 discussed the second case of bifurcations of Kokubu [6]. Actually, we also investigate the first case of bifurcation as in **Figure 2** in the same way and have the following result. We assume

B1 for $v = 0$, $\varepsilon = 0$, unperturbed equation

$$\dot{x} = f(x, 0, 0) \tag{3.41}$$

Admits three hyperbolic equilibriums p_1, p_2, p_3 and two heteroclinic orbits $q_1(t), q_2(t)$ connecting p_1 to p_2, p_2 to p_3 , respectively, that

$$\lim_{t \rightarrow -\infty} q_1(t) = p_1, \lim_{t \rightarrow \infty} q_1(t) = p_2$$

$$\lim_{t \rightarrow -\infty} q_2(t) = p_2, \lim_{t \rightarrow \infty} q_2(t) = p_3$$

We denote by $\Pi = q_1(t) \cup q_2(t) \cup q_1 \cup q_2 \cup p_3$.

Theorem 2 We assume the conditions B1, C2 and C3 are satisfied, then when ε, ρ sufficiently small Equation (1.1) admits two hyperbolic equilibrium $p_1(\varepsilon, \rho), p_2(\varepsilon, \rho)$ satisfying $p_1(0, 0) = p_1, p_2(0, 0) = p_2$. If the 2×2 matrix

$$M = \begin{pmatrix} M_1 \\ M_2 \end{pmatrix}$$

Is invertible, then for $\varepsilon > 0$ sufficiently small there exists a continuous function $\rho = \rho(\varepsilon)$ satisfying

$$\rho(0) = -M^{-1} \left(\begin{aligned} & \int_{-\infty}^{+\infty} \psi_1^*(s) (f_\varepsilon(q_1(s), 0, 0) ds) \\ & \int_{-\infty}^{+\infty} \psi_2^*(s) (f_\varepsilon(q_2(s), 0, 0) ds) \end{aligned} \right)$$

Such that the equation

$$\dot{x} = f(t, \varepsilon \rho(\varepsilon), \varepsilon) \tag{3.42}$$

Admits a heteroclinic orbit connecting $p_1(\varepsilon, \varepsilon \rho(\varepsilon))$ to $p_3(\varepsilon, \varepsilon \rho(\varepsilon))$ in the neighbourhood of the heteroclinic cycle Π .

REFERENCES

- [1] S. Wiggins, "Global Bifurcations and Chaos," Springer-Verlag, New York, 1988.
- [2] K. J. Palmer, "Transversal Heteroclinic Orbits and Cherry's Example of a Nonintegrable Hamiltonian System," *Journal of Differential Equations*, Vol. 65, No. 3, 1986, pp. 321-360. [doi:10.1016/0022-0396\(86\)90023-9](https://doi.org/10.1016/0022-0396(86)90023-9)
- [3] K. J. Palmer, "Exponential Dichotomies and Transversal Homoclinic Points," *Journal of Differential Equations*, Vol. 55, No. 2, 1984, pp. 225-256. [doi:10.1016/0022-0396\(84\)90082-2](https://doi.org/10.1016/0022-0396(84)90082-2)
- [4] S. Campbell and P. Holmes, "Bifurcation from O(2) Symmetric Heteroclinic Cycles with Three Interacting Modes," *Nonlinearity*, Vol. 4, 1991, pp. 697-726.
- [5] K. R. Meyer and G. R. Sell, "Melnikov Transforms, Bernoulli Bundle and Almost Periodic Perturbations," *Transactions of the American Mathematical Society*, Vol. 314, No. 1, 1989, pp. 63-105.
- [6] H. Kokubu, "Homoclinic and Heteroclinic Bifurcations of Vector Fields," *Japan Journal of Industrial and Applied Mathematics*, Vol. 5, No. 3, 1988, pp. 455-501. [doi:10.1007/BF03167912](https://doi.org/10.1007/BF03167912)
- [7] S. N. Chow, B. Deng and D. Terman, "The Bifurcations of a Homoclinic and a Periodic Orbit from Two Heteroclinic Orbits," *SIAM Journal on Mathematical Analysis*, Vol. 21, No. 1, 2000, pp. 179-204. [doi:10.1137/0521010](https://doi.org/10.1137/0521010)
- [8] J. M. Gambaudo, P. Glendinning and C. Tresser, "Collages de Cycles et Suites de Farey," *Comptes Rendus de l'Académie des Sciences*, Vol. 299, 1984, pp. 711-714.
- [9] S. N. Chow, B. Deng and D. Terman, "The Bifurcation of a Homoclinic Orbit from Two Heteroclinic Orbits—A Topological Approach," *Applicable Analysis: An International Journal*, Vol. 42, No. 1-4, 1991, pp. 1057-1080. [doi:10.1080/00036819108840047](https://doi.org/10.1080/00036819108840047)
- [10] X. B. Lin, "Using Melnikov's Method to Solve Silnikov Problems," *Proceedings of the Royal Society of Edinburgh: Section A Mathematics*, Vol. 116, No. 3-4, 1990, pp. 295-325. [doi:10.1017/S0308210500031528](https://doi.org/10.1017/S0308210500031528)
- [11] B. Sandstede and A. Scheel, "Forced Symmetry Breaking of Homoclinic Cycles," *Nonlinearity*, Vol. 8, No. 3, 2009, pp. 333-365. [doi:10.1088/0951-7715/8/3/003](https://doi.org/10.1088/0951-7715/8/3/003)
- [12] J. Guckenheimer and P. Holmes, "Structurally Stable Pulse Heteroclinic Cycles," *Mathematical Proceedings of the Cambridge Philosophical Society*, Vol. 103, No. 1, 2008,

- pp. 189-192. [doi:10.1017/S0305004100064732](https://doi.org/10.1017/S0305004100064732)
- [13] M. Krupa and I. Melbourne, "Asymptotic Stability of Heteroclinic Cycles in Systems with Symmetry," *Proceedings of the Royal Society of Edinburgh: Section A Mathematics*, Vol. 134, No. 6, 2004, pp. 1177-1197. [doi:10.1017/S0308210500003693](https://doi.org/10.1017/S0308210500003693)
- [14] M. Krupa, "Robust Heteroclinic Cycles," *Journal of Nonlinear Science*, Vol. 7, No. 2, 2011, pp. 129-176. [doi:10.1007/BF02677976](https://doi.org/10.1007/BF02677976)
- [15] W. A. Coppel, "Dichotomies in Stability Theory, Lecture Notes in Mathematics," Springer-Verlag, New York, 1978.
- [16] R. J. Sacker and G. R. Sell, "A Spectral Theory for Linear Differential Systems," *Journal of Differential Equations*, Vol. 27, No. 3, 1978, pp. 320-385. [doi:10.1016/0022-0396\(78\)90057-8](https://doi.org/10.1016/0022-0396(78)90057-8)
- [17] P. Chossat, M. Krupa, I. Melbourne and A. Scheel, "Transverse Bifurcations of Homoclinic Cycles," *Physica D: Nonlinear Phenomena*, Vol. 100, No. 1-2, 2011, pp. 85-100. [doi:10.1016/S0167-2789\(96\)00186-8](https://doi.org/10.1016/S0167-2789(96)00186-8)
- [18] V. Naudot, "Hyperbolic Dynamice in the Unfolding of a Degenerate Homoclinic Orbit," Preprint.
- [19] W. Y. Zeng, "Exponential Dichotomies and Transversal Homoclinic Orbits in Degenerate Cases," *Journal of Dynamics and Differential Equations*, Vol. 7, No. 4, 1995, pp. 521-548. [doi:10.1007/BF02218723](https://doi.org/10.1007/BF02218723)
- [20] G. R. Sell, "Bifurcation of Higher Dimensional Tori," *Archive for Rational Mechanics and Analysis*, Vol. 69, No. 3, 1979, pp. 199-230. [doi:10.1007/BF00248134](https://doi.org/10.1007/BF00248134)