

# Combinatorial Proof of the Inversion Formula on the Kazhdan-Lusztig $R$ -Polynomials

William Y.C. Chen<sup>1</sup>, Neil J.Y. Fan<sup>2</sup>, Alan J.X. Guo<sup>3</sup>, Peter L. Guo<sup>4</sup>  
Harry H.Y. Huang<sup>5</sup>, Michael X.X. Zhong<sup>6</sup>

<sup>1,3,4,5,6</sup>Center for Combinatorics, LPMC-TJKLC  
Nankai University, Tianjin 300071, P.R. China

<sup>2</sup>Department of Mathematics  
Sichuan University, Chengdu, Sichuan 610064, P.R. China

<sup>1</sup>chen@nankai.edu.cn, <sup>2</sup>fan@scu.edu.cn, <sup>3</sup>aalen@mail.nankai.edu.cn  
<sup>4</sup>lguo@nankai.edu.cn, <sup>5</sup>hhuang@cfc.nankai.edu.cn  
<sup>6</sup>michaelzhong@mail.nankai.edu.cn

## Abstract

In this paper, we present a combinatorial proof of the inversion formula on the Kazhdan-Lusztig  $R$ -polynomials. This problem was raised by Brenti. As a consequence, we obtain a combinatorial interpretation of the equi-distribution property due to Verma stating that any nontrivial interval of a Coxeter group in the Bruhat order has as many elements of even length as elements of odd length. The same argument leads to a combinatorial proof of an extension of Verma's equi-distribution to the parabolic quotients of a Coxeter group obtained by Deodhar. As another application, we derive a refinement of the inversion formula for the symmetric group by restricting the summation to permutations ending with a given element.

**Keywords:** Kazhdan-Lusztig  $R$ -polynomial, inversion formula, Bruhat order

**AMS Classification:** 05A19, 05E15, 20F55

## 1 Introduction

Let  $(W, S)$  be a Coxeter system. For  $u, v \in W$ , let  $R_{u,v}(q)$  be the Kazhdan-Lusztig  $R$ -polynomial indexed by  $u$  and  $v$ . The following inversion formula was obtained by Kazhdan and Lusztig [8]:

$$\sum_{u \leq w \leq v} (-1)^{\ell(w) - \ell(u)} R_{u,w}(q) R_{w,v}(q) = \delta_{u,v}, \quad (1.1)$$

where  $\leq$  is the Bruhat order and  $\ell$  is the length function, see also Humphreys [6]. The aim of this paper is to present a combinatorial interpretation of this formula. This problem was raised by Brenti [3].

To give a combinatorial proof of (1.1), we start with Dyer's combinatorial description of the  $R$ -polynomials in terms of increasing Bruhat paths [5]. Then we reformulate the inversion formula in terms of  $V$ -paths. For  $u \leq w \leq v$ , by a  $V$ -path from  $u$  to  $v$  with bottom  $w$  we mean a pair  $(\Delta_1, \Delta_2)$  of Bruhat paths such that  $\Delta_1$  is a decreasing path from  $u$  to  $w$  and  $\Delta_2$  is an increasing path from  $w$  to  $v$ . We construct an involution on  $V$ -paths. This leads to a combinatorial proof of (1.1).

We give two applications of the involution. First, we restrict the involution to  $V$ -paths from  $u$  to  $v$  with maximal length. This induces an involution on the interval  $[u, v]$  with  $u < v$ , which leads to a combinatorial proof of the equi-distribution property that any nontrivial interval  $[u, v]$  has as many elements of even length as elements of odd length. This property was proved inductively by Verma [12], which was used to deduce the Möbius function of the Bruhat order. Other proofs of the Möbius function formula for the Bruhat order can be found in [2, 4, 9, 11]. Recently, Jones [7] found a combinatorial proof for the equi-distribution property by constructing an involution on the intervals of a Coxeter group  $W$ . When  $W$  is finite, Jones [7] showed that this involution agrees with the construction of Rietsch and Williams [10] in their study of discrete Morse theory and totally nonnegative flag varieties.

The idea that we have used to prove Verma's equi-distribution can also be applied to Deodhar's [4] extension to parabolic quotients. For  $J \subseteq S$ , let  $W_J$  be the parabolic subgroup of  $W$  generated by  $J$ , and let  $W^J$  be the quotient of  $W$  consisting of minimal representatives of the left cosets of  $W_J$  in  $W$ , that is,

$$W^J = \{w \in W \mid \ell(ws) > \ell(w) \text{ for any } s \in J\}.$$

The quotient  $W^J$  forms a subposet of  $W$  in the Bruhat order. For  $u \leq v \in W^J$ , let

$$[u, v]^J = [u, v] \cap W^J$$

and let

$$K_J(u, v) = \{w \in [u, v]^J \mid [w, v]^J = [w, v]\}.$$

When  $u < v$ , Deodhar [4] showed that  $K_J(u, v)$  contains as many elements of even length as elements of odd length, from which the Möbius function of the Bruhat order on  $W^J$  can be easily deduced. When  $J = \emptyset$ , Deodhar's assertion reduces to Verma's equi-distribution. The Möbius function on  $W^J$  was rederived by Björner and Wachs [2] with the aid of topological techniques, and by Stembridge [11] by an algebraic approach. We construct an involution on  $K_J(u, v)$  that leads to a simple combinatorial interpretation of Deodhar's equi-distribution.

As a second application, we find a refinement of the inversion formula when  $W$  is the symmetric group  $S_n$ . For a permutation  $w \in S_n$ , we write  $w = w(1)w(2) \cdots w(n)$ ,

where  $w(i)$  denotes the element in the  $i$ -th position. Let  $u$  and  $v$  be two permutations in  $S_n$  such that  $u < v$  in the Bruhat order. For  $1 \leq k \leq n$ , let  $[u, v]_k$  denote the set of permutations in the interval  $[u, v]$  that end with  $k$ , that is,

$$[u, v]_k = \{w \in [u, v] \mid w(n) = k\}.$$

By using a variation of the involution, we show that the summation

$$\sum_{w \in [u, v]_k} (-1)^{\ell(w) - \ell(u)} R_{u, w}(q) R_{w, v}(q)$$

equals zero or a power of  $q$  up to a sign.

## 2 An involution on $V$ -paths

Our combinatorial proof of the inversion formula is based on an equivalent formulation of (1.1) in terms of the  $\tilde{R}$ -polynomials. Let  $(W, S)$  be a Coxeter system. For  $u, v \in W$  with  $u \leq v$ , the  $\tilde{R}$ -polynomials  $\tilde{R}_{u, v}(q)$  were introduced by Dyer [5], which are connected to the  $R$ -polynomials via the following relation

$$R_{u, v}(q) = q^{\frac{\ell(v) - \ell(u)}{2}} \tilde{R}_{u, v}(q^{\frac{1}{2}} - q^{-\frac{1}{2}}),$$

see also Björner and Brenti [1]. Thus the inversion formula (1.1) can be restated as

$$\sum_{u \leq w \leq v} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u, w}(q) \tilde{R}_{w, v}(q) = \delta_{u, v}. \quad (2.1)$$

To give a bijective proof of (2.1), we need a combinatorial interpretation of the  $\tilde{R}$ -polynomials due to Dyer [5] in terms of increasing Bruhat paths of a Coxeter group. For a Coxeter system  $(W, S)$ , let

$$T = \{wsw^{-1} \mid s \in S, w \in W\}$$

be the set of reflections. The Bruhat graph  $BG(W)$  of  $W$  is a directed graph whose nodes are the elements of  $W$  such that there is an arc from  $u$  to  $v$  if  $v = ut$  for some  $t \in T$  and  $\ell(u) < \ell(v)$ . We use  $u \xrightarrow{t} v$  to denote the arc from  $u$  to  $v$ , where  $t$  is the reflection such that  $v = ut$ . An increasing path in the Bruhat graph is defined based on the reflection ordering on the positive roots of  $W$ . Let  $\Phi$  be the root system of  $W$ , and  $\Phi^+$  be the positive root system. A total ordering  $\prec$  on  $\Phi^+$  is called a reflection ordering if for any  $\alpha \prec \beta \in \Phi^+$  and two nonnegative real numbers  $\lambda, \mu$  such that  $\lambda\alpha + \mu\beta \in \Phi^+$ , then we have  $\alpha \prec \lambda\alpha + \mu\beta \prec \beta$ . Since positive roots in  $\Phi^+$  are in one-to-one correspondence with reflections, a reflection ordering induces a total ordering on the reflection set  $T$ .

Let  $\Delta = u_0 \xrightarrow{t_1} u_1 \xrightarrow{t_2} \cdots \xrightarrow{t_r} u_r$  be a path from  $u$  to  $v$ , where  $u_0 = u$  and  $u_r = v$ . We say that  $\Delta$  is increasing if  $t_1 \prec t_2 \prec \cdots \prec t_r$ , and  $\Delta$  is decreasing if  $t_1 \succ t_2 \succ \cdots \succ t_r$ .

Let  $\ell(\Delta)$  denote the length of  $\Delta$ , that is, the number of arcs in  $\Delta$ . Dyer [5] showed that for any fixed reflection ordering  $\prec$  on  $T$ , we have

$$\tilde{R}_{u,v}(q) = \sum_{\Delta} q^{\ell(\Delta)}, \quad (2.2)$$

where the sum ranges over increasing Bruhat paths from  $u$  to  $v$  with respect to  $\prec$ , see also Björner and Brenti [1]. By definition, the reverse of a reflection ordering is also a reflection ordering. So (2.2) can be restated as

$$\tilde{R}_{u,v}(q) = \sum_{\Delta'} q^{\ell(\Delta')},$$

where the sum ranges over decreasing Bruhat paths from  $u$  to  $v$  with respect to  $\prec$ .

By a  $V$ -path from  $u$  to  $v$  with bottom  $w$ , we mean a pair  $(\Delta_1, \Delta_2)$  of Bruhat paths such that  $\Delta_1$  is a decreasing path from  $u$  to  $w$  and  $\Delta_2$  is an increasing path from  $w$  to  $v$ . The sign of a  $V$ -path  $(\Delta_1, \Delta_2)$  is defined as

$$\text{sgn}(\Delta_1, \Delta_2) = (-1)^{\ell(\Delta_1)}.$$

The length of a Bruhat path from  $u$  to  $w$  has the same parity as  $\ell(w) - \ell(u)$ , see, e.g., Björner and Brenti [1]. It follows that

$$\text{sgn}(\Delta_1, \Delta_2) = (-1)^{\ell(w) - \ell(u)},$$

and so (2.1) can be rewritten as

$$\sum_{u \leq w \leq v} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u,w}(q) \tilde{R}_{w,v}(q) = \sum_{(\Delta_1, \Delta_2)} \text{sgn}(\Delta_1, \Delta_2) q^{\ell(\Delta_1) + \ell(\Delta_2)} = \delta_{u,v}, \quad (2.3)$$

where the second sum ranges over  $V$ -paths from  $u$  to  $v$ .

We now define an involution  $\Phi$  on  $V$ -paths, which preserves the length, but reverses the sign of a  $V$ -path. This leads to a combinatorial proof of (2.3).

**An Involution  $\Phi$  on  $V$ -Paths:** For  $u < v$ , let  $(\Delta_1, \Delta_2)$  be a  $V$ -path from  $u$  to  $v$  with bottom  $w$ . Write

$$\Delta_1 = u_0 \xrightarrow{t_1} u_1 \xrightarrow{t_2} \cdots \xrightarrow{t_i} u_i \quad \text{and} \quad \Delta_2 = v_0 \xrightarrow{t'_1} v_1 \xrightarrow{t'_2} \cdots \xrightarrow{t'_j} v_j,$$

where  $u_0 = u$ ,  $u_i = v_0 = w$  and  $v_j = v$ . The  $V$ -path  $\Phi(\Delta_1, \Delta_2) = (\Delta'_1, \Delta'_2)$  is constructed according to the following two cases.

Case 1:  $u = w$  or  $t_i \succ t'_1$ . Set

$$\Delta'_1 = u_0 \xrightarrow{t_1} u_1 \xrightarrow{t_2} \cdots \xrightarrow{t_i} u_i \xrightarrow{t'_1} v_1 \quad \text{and} \quad \Delta'_2 = v_1 \xrightarrow{t'_2} \cdots \xrightarrow{t'_j} v_j.$$

Case 2:  $v = w$  or  $t_i \prec t'_1$ . Set

$$\Delta'_1 = u_0 \xrightarrow{t_i} u_1 \xrightarrow{t_2} \cdots \xrightarrow{t_{i-1}} u_{i-1} \quad \text{and} \quad \Delta'_2 = u_{i-1} \xrightarrow{t_i} v_0 \xrightarrow{t'_1} v_1 \xrightarrow{t'_2} \cdots \xrightarrow{t'_j} v_j.$$

It turns out that the involution  $\Phi$  yields a simple combinatorial interpretation of the following parity property of Verma [12].

**Theorem 2.1** (Verma [12]) *Let  $(W, S)$  be a Coxeter system and  $u < v \in W$ . Then the interval  $[u, v]$  has the same number of elements of odd length as elements of even length.*

Indeed, for  $u < v \in W$ , there exists a unique maximal increasing (or, decreasing) Bruhat path from  $u$  to  $v$  [5]. Thus, for any  $w \in [u, v]$  there is a unique maximal  $V$ -path from  $u$  to  $v$  with bottom  $w$ . So the maximal  $V$ -paths from  $u$  to  $v$  are in one-to-one correspondence with elements in the interval  $[u, v]$ . Restricting the involution  $\Phi$  to the maximal  $V$ -paths from  $u$  to  $v$  induces an involution on the interval  $[u, v]$ , which reverses the parity of the length of each element in  $[u, v]$ . This proves Theorem 2.1.

The above argument also serves as a combinatorial interpretation of the following equi-distribution due to Deodhar [4]. Let us recall the common notation as mentioned in Introduction. For  $J \subseteq S$ , let

$$W^J = \{w \in W \mid \ell(ws) > \ell(w) \text{ for any } s \in J\}.$$

For  $u \leq v \in W^J$ , let

$$[u, v]^J = [u, v] \cap W^J$$

and let

$$K_J(u, v) = \{w \in [u, v]^J \mid [w, v]^J = [w, v]\}.$$

**Theorem 2.2** (Deodhar [4]) *Let  $(W, S)$  be a Coxeter system, and  $J \subseteq S$ . Then, for  $u < v \in W$ , the set  $K_J(u, v)$  has the same number of elements of odd length as elements of even length.*

To construct an involution on  $K_J(u, v)$ , we recall a labeling on the edges of the poset  $[u, v]^J$  introduced by Björner and Wachs [2], see also Björner and Brenti [1]. Let  $v = s_1 s_2 \cdots s_q$  be a given reduced expression of  $v$ . We read a maximal chain in  $[u, v]^J$  from top to bottom. Let  $v = w_0 \rightarrow w_1 \rightarrow \cdots \rightarrow w_r = u$  be a maximal chain in  $[u, v]^J$ , where  $r = \ell(v) - \ell(u)$ . Then there is a unique sequence  $(i_1, i_2, \dots, i_r)$  of distinct integers such that for  $1 \leq k \leq r$ ,  $w_k$  has a reduced expression obtained from  $s_1 s_2 \cdots s_q$  by deleting simple reflections indexed by  $i_1, i_2, \dots, i_k$ . Label the edge from  $w_{k-1}$  to  $w_k$  by  $i_k$ . We denote the maximal chain with such a labeling by  $v = w_0 \xrightarrow{i_1} w_1 \xrightarrow{i_2} \cdots \xrightarrow{i_r} w_r = u$ , and say that the chain  $v = w_0 \xrightarrow{i_1} w_1 \xrightarrow{i_2} \cdots \xrightarrow{i_r} w_r = u$  is increasing if  $i_1 < i_2 < \cdots < i_r$ , and it is decreasing if  $i_1 > i_2 > \cdots > i_r$ . The following theorem is due to Björner and Wachs [2], see also Björner and Brenti [1].

**Theorem 2.3** (Björner and Wachs [2]) *Let  $u < v \in W^J$ , and let  $v = s_1 s_2 \cdots s_q$  be a given reduced expression of  $v$ . Then there is a unique increasing maximal chain from  $v$  to  $u$  in  $[u, v]^J$ .*

We remark that when  $J = \emptyset$ , the proof of Theorem 2.3 can be employed to show that for any given reduced expression of  $v$ , there is a unique decreasing maximal chain from  $v$  to  $u$  in  $[u, v]^\emptyset = [u, v]$ .

We are now ready to present an involution  $\Psi$  on  $K_J(u, v)$ , which reverses the parity of the length. This leads to a combinatorial proof of Theorem 2.2.

**An Involution  $\Psi$  on  $K_J(u, v)$ :** Let  $w \in K_J(u, v)$ , and let  $v = s_1 s_2 \cdots s_q$  be a fixed reduced expression of  $v$ . Since  $[w, v]^J = [w, v]$ , by the above remark, there exists a unique decreasing maximal chain  $v = v_0 \xrightarrow{i_1} v_1 \xrightarrow{i_2} \cdots \xrightarrow{i_m} v_m = w$  from  $v$  to  $w$  in  $[u, v]^J$ . Let  $w = s_{k_1} s_{k_2} \cdots s_{k_p}$  be the reduced expression of  $w$  obtained from  $s_1 s_2 \cdots s_q$  by deleting the generators indexed by  $i_1, i_2, \dots, i_m$ , that is,  $1 \leq k_1 < k_2 < \cdots < k_p \leq q$  and  $\{k_1, k_2, \dots, k_p\} = \{1, 2, \dots, q\} \setminus \{i_1, i_2, \dots, i_m\}$ . Assume that  $w = w_0 \xrightarrow{k_{j_1}} w_1 \xrightarrow{k_{j_2}} \cdots \xrightarrow{k_{j_t}} w_t = u$  is the unique increasing maximal chain in  $[u, w]^J$  with respect to the reduced expression  $w = s_{k_1} s_{k_2} \cdots s_{k_p}$ . Note that  $1 \leq j_1 < \cdots < j_t \leq p$ . Then  $\Psi(w)$  is defined according to the following two cases:

Case 1:  $u = w$  or  $i_m < k_{j_1}$ . Set  $\Psi(w) = v_{m-1}$ ;

Case 2:  $v = w$  or  $i_m > k_{j_1}$ . Set  $\Psi(w) = w_1$ .

The following theorem shows that  $\Psi$  is an involution on  $K_J(u, v)$ . The proof relies on the following properties of the Bruhat order, see, for example, Björner and Brenti [1].

**The Subword Property:** Let  $u, v \in W$ . Then  $u \leq v$  in the Bruhat order if and only if every reduced expression of  $v$  has a subword that is a reduced expression of  $u$ .

**The Lifting Property:** Suppose that  $u < v \in W$ , and  $s \in S$  is a simple reflection. If  $\ell(sv) < \ell(v)$  and  $\ell(su) > \ell(u)$ , then  $u \leq sv$  and  $su \leq v$ . Similarly, if  $\ell(vs) < \ell(v)$  and  $\ell(us) > \ell(u)$ , then  $u \leq vs$  and  $us \leq v$ .

**Theorem 2.4** *The map  $\Psi$  is an involution on  $K_J(u, v)$ .*

*Proof.* By the construction of  $\Psi$ , it suffices to show that for  $w \in K_J(u, v)$ ,  $\Psi(w)$  also belongs to  $K_J(u, v)$ . This is trivial when  $u = w$  or  $i_m < k_{j_1}$ . Now we consider the case when  $v = w$  or  $i_m > k_{j_1}$ . Let  $w' = \Psi(w)$ . Assume that  $w = s_1 \cdots \hat{s}_{i_m} \cdots \hat{s}_{i_2} \cdots \hat{s}_{i_1} \cdots s_q$  and  $w' = s_1 \cdots \hat{s}_{k_{j_1}} \cdots \hat{s}_{i_m} \cdots \hat{s}_{i_1} \cdots s_q$ , where for a simple reflection  $s \in S$ ,  $\hat{s}$  means that  $s$  is missing. We aim to prove that  $w' \in K_J(u, v)$ .

Suppose to the contrary that  $w' \notin K_J(u, v)$ . Then there exists an element  $w'' \in [w', v]$  such that  $w'' \notin [w', v]^J$ . By definition, there exists  $s \in J$  such that  $\ell(w''s) < \ell(w'')$ . Since  $\ell(w's) > \ell(w')$ , the lifting property implies that  $w's \leq w''$ . Thus we have  $w's \leq v$ . Since  $\ell(vs) > \ell(v)$ , we see that  $w's \neq v$ . It follows that  $w's < v$ , that is,

$$s_1 \cdots \hat{s}_{k_{j_1}} \cdots \hat{s}_{i_m} \cdots \hat{s}_{i_1} \cdots s_q s < s_1 s_2 \cdots s_q.$$

It is easily checked that  $\hat{s}_{k_{j_1}} \cdots \hat{s}_{i_m} \cdots \hat{s}_{i_1} \cdots s_q s < s_{k_{j_1}} \cdots s_q$ . By the lifting property,

we deduce that  $s_{k_{j_1}} \cdots \hat{s}_{i_m} \cdots \hat{s}_{i_1} \cdots s_q s \leq s_{k_{j_1}} \cdots s_q$ . Thus we have

$$ws = s_1 \cdots \hat{s}_{i_m} \cdots \hat{s}_{i_1} \cdots s_q s \leq s_1 \cdots s_q = v,$$

which implies that  $ws \in [w, v]$ . On the other hand, it is obvious that  $ws \notin [w, v]^J$ . So we conclude that  $w \notin K_J(u, v)$ , contradicting the assumption that  $w \in K_J(u, v)$ . This completes the proof.  $\blacksquare$

From the proof of Theorem 2.4, we see that for  $w \in [u, v]^J$ ,  $w \in K_J(u, v)$  if and only if there does not exist any  $s \in J$  such that  $ws \in [u, v]$ . Notice that this characterization has been observed by Deodhar [4, Lemma 3].

### 3 A refinement of the inversion formula for $S_n$

In this section, we use a variation of the involution  $\Phi$  to give a refinement of the inversion formula for the symmetric group  $S_n$ . We introduce the notion of an  $S$ -interval. Let  $u, v$  be two permutations in  $S_n$  with  $u < v$ . Let

$$D(u, v) = \{1 \leq i \leq n \mid u(i) \neq v(i)\}.$$

Suppose that  $D(u, v) = \{i_1, i_2, \dots, i_j\}_<$ , that is,  $D(u, v) = \{i_1, i_2, \dots, i_j\}$  and  $i_1 < i_2 < \dots < i_j$ . Let  $b_1 < b_2 < \dots < b_j$  be the values of  $u(i_1), u(i_2), \dots, u(i_j)$  listed in increasing order. We say that  $[u, v]$  is an  $S$ -interval if it satisfies the following conditions:

- (1)  $i_j = n$  and  $u(i_j) = b_j$ ;
- (2) The values in  $\{b_1, b_2, \dots, b_j\}$  that are greater than  $u(i_1)$  appear in increasing order in  $u$ , whereas the values in  $\{b_1, b_2, \dots, b_j\}$  that are less than  $u(i_1)$  appear in decreasing order in  $u$ ;
- (3) In the cycle notation,  $v = (b_1, b_2, \dots, b_j)u$ , that is,  $v$  is obtained from  $u$  by rotating the elements  $b_1, b_2, \dots, b_j$  in  $u$ .

Recall that for  $u < v \in S_n$ ,  $[u, v]_k$  denotes the set of permutations in  $[u, v]$  that end with  $k$ . The following theorem gives a refinement of the inversion formula for  $S_n$ .

**Theorem 3.1** *Assume that  $u < v \in S_n$ . Let  $m$  be the smallest index such that  $u(m) \neq v(m)$ . If  $[u, v]$  is an  $S$ -interval, and  $k = u(m)$  or  $k = v(m)$ , then we have*

$$\sum_{w \in [u, v]_k} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u, w}(q) \tilde{R}_{w, v}(q) = (-1)^r q^{s-1},$$

where  $s = |D(u, v)|$  and

$$r = |\{j \in D(u, v) \mid u(j) > k\}|;$$

Otherwise, we have

$$\sum_{w \in [u, v]_k} (-1)^{\ell(w) - \ell(u)} \tilde{R}_{u, w}(q) \tilde{R}_{w, v}(q) = 0.$$

For  $1 \leq k \leq n$ , let  $P_k(u, v)$  denote the set of  $V$ -paths from  $u$  to  $v$  with bottoms contained in  $[u, v]_k$ . To prove Theorem 3.1, we shall construct an involution  $\Omega$  on  $P_k(u, v)$ . The reflection set  $T$  of  $S_n$  consists of transpositions of  $S_n$ , that is,

$$T = \{(i, j) \mid 1 \leq i < j \leq n\}.$$

For two permutations  $u, v$  in  $S_n$ , it is known that there is an arc from  $u$  to  $v$  in the Bruhat graph of  $S_n$  if  $v = u(i, j)$  and  $u(i) < u(j)$ , see Björner and Brenti [1].

From now on, we choose the reflection ordering  $\prec$  on  $T$  to be the lexicographic ordering:

$$(1, 2) \prec (1, 3) \prec \cdots \prec (1, n) \prec (2, 3) \prec \cdots \prec (n-1, n). \quad (3.1)$$

For a Bruhat path  $\Delta = u_0 \xrightarrow{t_1} u_1 \xrightarrow{t_2} \cdots \xrightarrow{t_r} u_r$ , let

$$L(\Delta) = (t_1, t_2, \dots, t_r).$$

**An Involution  $\Omega$  on  $P_k(u, v)$ :** Let  $(\Delta_1, \Delta_2)$  be a  $V$ -path in  $P_k(u, v)$  with bottom  $w$ .

Write  $\Delta_1 = u_0 \xrightarrow{t_1} u_1 \xrightarrow{t_2} \cdots \xrightarrow{t_i} u_i$  and  $\Delta_2 = v_0 \xrightarrow{t'_1} v_1 \xrightarrow{t'_2} \cdots \xrightarrow{t'_j} v_j$ , where  $u_0 = u$ ,  $u_i = v_0 = w$  and  $v_j = v$ . Let  $t = \min\{t_i, t'_1\}$ . Then the  $V$ -path  $\Omega(\Delta_1, \Delta_2) = (\Delta'_1, \Delta'_2)$  is defined as follows. We consider three cases.

Case 1:  $t$  is an internal transposition, that is,  $t = (a, b)$  and  $1 \leq a < b < n$ . In this case, set  $(\Delta'_1, \Delta'_2) = \Phi(\Delta_1, \Delta_2)$ .

Case 2:  $t$  is a boundary transposition, that is,  $t = (a, n)$  for some  $a < n$ , and there is an internal transposition among the transpositions  $t_1, \dots, t_i, t'_1, \dots, t'_j$ . Let  $\tilde{t}$  be the smallest internal transposition among  $t_1, \dots, t_i, t'_1, \dots, t'_j$ . By the choice of the reflection ordering in (3.1), it is easy to check that  $\tilde{t}$  belongs to either  $\{t_1, \dots, t_i\}$  or  $\{t'_1, \dots, t'_j\}$ , but not both. So we have the following two subcases.

Subcase 1:  $\tilde{t}$  belongs to  $\{t_1, \dots, t_i\}$ . Assume that  $t_{i_0} = \tilde{t}$ , where  $1 \leq i_0 \leq i$ . Let  $\Delta'_1$  be the path such that  $L(\Delta'_1)$  is the sequence obtained from  $L(\Delta_1)$  by deleting  $t_{i_0}$ , and let  $\Delta'_2$  be the path such that  $L(\Delta'_2)$  is the sequence obtained from  $L(\Delta_2)$  by inserting  $t_{i_0}$  such that  $L(\Delta'_2)$  remains increasing.

Subcase 2:  $\tilde{t}$  belongs to  $\{t'_1, \dots, t'_j\}$ . Assume that  $t'_{j_0} = \tilde{t}$ , where  $1 \leq j_0 \leq j$ . Let  $\Delta'_2$  be the path such that  $L(\Delta'_2)$  is the sequence obtained from  $L(\Delta_2)$  by deleting  $t'_{j_0}$ , and let  $\Delta'_1$  be the path such that  $L(\Delta'_1)$  is the sequence obtained from  $L(\Delta_1)$  by inserting  $t_{j_0}$  such that  $L(\Delta'_1)$  remains decreasing.

Case 3: The transpositions  $t_1, \dots, t_i, t'_1, \dots, t'_j$  are all boundary transpositions. In this case, set  $(\Delta'_1, \Delta'_2) = (\Delta_1, \Delta_2)$ .



It is easy to verify that  $\Omega$  is a length preserving involution on  $P_k(u, v)$ , and it is clear that  $\Omega$  reverses the sign of  $(\Delta_1, \Delta_2)$  unless  $(\Delta_1, \Delta_2)$  is a fixed point. To prove Theorem 3.1, we also need the following property.

**Proposition 3.2** *Assume that  $u < v \in S_n$  and  $1 \leq k \leq n$ . Then the involution  $\Omega$  on  $P_k(u, v)$  has at most one fixed point. Moreover,  $\Omega$  has a fixed point if and only if  $[u, v]$  is an  $S$ -interval and  $k = u(m)$  or  $k = v(m)$ , where  $m$  is the smallest integer such that  $u(m) \neq v(m)$ .*

*Proof.* To prove that  $\Omega$  has at most one fixed point, assume that  $(\Delta_1, \Delta_2) \in P_k(u, v)$  is a  $V$ -path that is fixed by  $\Omega$ . We proceed to show that  $(\Delta_1, \Delta_2)$  is uniquely determined.

Let  $\Delta_1 = u_0 \xrightarrow{t_1} u_1 \xrightarrow{t_2} \dots \xrightarrow{t_i} u_i$  and  $\Delta_2 = v_0 \xrightarrow{t'_1} v_1 \xrightarrow{t'_2} \dots \xrightarrow{t'_j} v_j$ . By the construction of  $\Omega$ , we see that  $t_1, \dots, t_i$  and  $t'_1, \dots, t'_j$  are all boundary transpositions. Assume that  $t_1 = (p_1, n), \dots, t_i = (p_i, n)$  and  $t'_1 = (p'_1, n), \dots, t'_j = (p'_j, n)$ . Since  $\Delta_1$  and  $\Delta_2$  are Bruhat paths, we see that

$$u(n) > u(p_1) > \dots > u(p_i) = k = w(n) > w(p'_1) > \dots > w(p'_j). \quad (3.2)$$

Noting that  $t_1 \succ t_2 \succ \dots \succ t_i$  and  $t'_1 \prec t'_2 \prec \dots \prec t'_j$ , we find that  $n > p_1 > \dots > p_i$  and  $p'_1 < \dots < p'_j < n$ .

By (3.2) together with the relation  $w = u(p_1, n) \dots (p_i, n)$ , it is easily seen that

$$\{p_1, \dots, p_i\} \cap \{p'_1, \dots, p'_j\} = \emptyset.$$

This yields that  $w(p'_1) = u(p'_1), \dots, w(p'_j) = u(p'_j)$ , and so (3.2) becomes

$$u(n) > u(p_1) > \dots > u(p_i) = k = w(n) > u(p'_1) > \dots > u(p'_j). \quad (3.3)$$

Observe that

$$\{p_1, \dots, p_i\} \cup \{p'_1, \dots, p'_j\} \cup \{n\} = D(u, v).$$

In view of (3.3), we deduce that given  $u, v$  and  $k$ , the values of  $i, j$  as well as the elements  $p_1, \dots, p_i, p'_1, \dots, p'_j$  are uniquely determined. In other words, the  $V$ -path  $(\Delta_1, \Delta_2)$  is uniquely determined.

It remains to prove that  $\Omega$  has a fixed point if and only if  $[u, v]$  is an  $S$ -interval and  $k = u(m)$  or  $k = v(m)$ . By the above argument, we see that if  $\Omega$  has a fixed point, then  $[u, v]$  is an  $S$ -interval and  $k = u(p_i) = v(p'_1)$ . Since  $m = \min\{p_i, p'_1\}$ , we obtain that  $k = u(m)$  if  $p_i < p'_1$  and  $k = v(m)$  if  $p_i > p'_1$ . Conversely, if  $[u, v]$  is an  $S$ -interval, it is easy to construct a  $V$ -path in  $P_k(u, v)$  fixed by  $\Omega$ , where  $k = u(m)$  or  $k = v(m)$ . This completes the proof.  $\blacksquare$

We are now ready to complete the proof of Theorem 3.1.

*Proof of Theorem 3.1.* By Proposition 3.2, we only need to consider the case when  $[u, v]$  is an  $S$ -interval and  $k = u(m)$  or  $k = v(m)$ . In this case, we have

$$\sum_{w \in [u, v]_k} (-1)^{\ell(w) - \ell(u)} \widetilde{R}_{u, w}(q) \widetilde{R}_{w, v}(q) = (-1)^{\ell(\Delta_1)} q^{\ell(\Delta_1) + \ell(\Delta_2)},$$

where  $(\Delta_1, \Delta_2)$  is the unique  $V$ -path in  $P_k(u, v)$  that is fixed by  $\Omega$ . Evidently,

$$\ell(\Delta_1) + \ell(\Delta_2) = |D(u, v)| - 1.$$

It is also clear that

$$\ell(\Delta_1) = |\{j \in D(u, v) \mid u(j) > k\}|.$$

Hence the proof is complete. ■

**Acknowledgments.** We wish to thank the referee for valuable suggestions. This work was supported by the 973 Project and the National Science Foundation of China.

## References

- [1] A. Björner and F. Brenti, *Combinatorics of Coxeter Groups*, Grad. Texts in Math., Vol. 231, Springer, New York, 2005.
- [2] A. Björner and M. Wachs, Bruhat order of Coxeter groups and shellability, *Adv. Math.* 43 (1982), 87–100.
- [3] F. Brenti, Kazhdan-Lusztig and  $R$ -polynomials from a combinatorial point of view, *Discrete Math.* 193 (1998), 93–116.
- [4] V.V. Deodhar, Some characterizations of Bruhat ordering on a Coxeter group and determination of the relative Möbius function, *Invent. Math.* 39 (1977), 187–198.
- [5] M.J. Dyer, Hecke algebras and shellings of Bruhat intervals, *Compos. Math.* 89 (1993), 91–115.
- [6] J.E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics, No. 29, Cambridge Univ. Press, Cambridge, 1990.
- [7] B.C. Jones, An explicit derivation of the Möbius function for Bruhat order, *Order* 26 (2009), 319–330.
- [8] D. Kazhdan and G. Lusztig, Representations of Coxeter groups and Hecke algebras, *Invent. Math.* 53 (1979), 165–184.
- [9] M. Marietti, Algebraic and combinatorial properties of zircons, *J. Algebraic Combin.* 26 (2007), 363–382.
- [10] K. Rietsch and L. Williams, Discrete Morse theory for totally non-negative flag varieties, *Adv. Math.* 223 (2010), 1855–1884.
- [11] J.R. Stembridge, A short derivation of the Möbius function for the Bruhat order, *J. Algebraic Combin.* 25 (2007), 141–148.
- [12] D.-N. Verma, Möbius inversion for the Bruhat ordering on a Weyl group, *Ann. Sci. École Norm. Sup.* 4 (1971), 393–398.