

to reach the desired audience. However, the methods used to assess their diversity or distribution are both unusual and, at present, not well established. Recognition of these facts by editors and reviewers of these journals and the appointment of microbial ecologists to their editorial boards would be great first steps.

All conservation efforts are ultimately funded by the public and here some education is needed, not only to counteract negative perceptions ('bad bacteria' or 'poisonous fungi'), but also to highlight the beauty and biotechnological utility of microbes, as well as their fundamental importance to ecosystem function. As Tom Curtis [12] has stated: 'if the last blue whale choked to death on the last panda, it would be disastrous but not the end of the world. But if we accidentally poisoned the last two species of ammonia-oxidizers, that would be another matter. It could be happening now and we wouldn't even know'.

Acknowledgements

Thanks are owed to the following for invaluable comments: Gareth Morris FRS, Dylan Gwynn-Jones, Elinor Gwynn, John Warren, Arwyn Edwards, Arthur Chater and Andy Mitchell.

References

- Whitman, W.B. *et al.* (1998) Prokaryotes: the unseen majority. *Proc. Natl. Acad. Sci. U.S.A.* 95, 6578–6583
- Secretariat of the Convention on Biological Diversity (2010) *Report of the Fourteenth Meeting of the Subsidiary Body on Scientific, Technical and Technological Advice*, UNEP/CBD/COP/10/3
- Baas Becking, L. (1934) *Geobiologie of inleiding tot de milieukunde*, W.P. van Stockum and Zoon [in Dutch]
- Redford, A.J. *et al.* (2010) The ecology of the phyllosphere: geographic and phylogenetic variability in the distribution of bacteria on tree leaves. *Environ. Microbiol.* 12, 2885–2893
- Gerstein, A.C. and Moore, J-S. (2011) Small is the new big: assessing the population structure of microorganisms. *Mol. Ecol.* 20, 4385–4387
- Caporaso, J.G. *et al.* (2011) Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc. Natl. Acad. Sci. U.S.A.* 108, 4516–4522
- Liggenstoffer, A.S. *et al.* (2010) Phylogenetic diversity and community structure of anaerobic gut fungi (phylum Neocallimastigomycota) in ruminant and non-ruminant herbivores. *ISME J.* 4, 1225–1235
- McCormick, M.K. *et al.* (2006) Orchid-fungus fidelity: a marriage meant to last? *Ecology* 87, 903–911
- Blaser, M.J. and Falkow, S. (2009) What are the consequences of the disappearing human microbiota? *Nat. Rev. Microbiol.* 7, 887–894
- Schiermeier, Q. (2011) Race against time for raiders of the lost lake. *Nature* 469, 275
- Parker, S.S. (2010) Buried treasure: soil biodiversity and conservation. *Biodivers. Conserv.* 19, 3743–3756
- Curtis, T.P. (2006) Microbial ecologists: it's time to 'go large'. *Nat. Rev. Microbiol.* 4, 488

0169-5347/\$ – see front matter © 2011 Elsevier Ltd. All rights reserved.

doi:10.1016/j.tree.2011.10.002 Trends in Ecology and Evolution, January 2012, Vol. 27, No. 1

Megadiverse developing countries face huge risks from invasives

Gábor L. Lövei^{1,2} and Thomas M. Lewinsohn³, the Biological Invasions in Megadiverse Regions Network*

¹Department of Agroecology, Aarhus University, Flakkebjerg Research Centre, DK-4200 Slagelse, Denmark

²Department of Biological Invasions, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, 100081 Beijing, China

³Department of Animal Biology, Institute of Biology, University of Campinas, Brazil

Recently, Davis *et al.* [1] claimed that “non-native” species have been vilified for... generally polluting “natural” environments’. They further assert that ‘a pervasive bias

against alien species... has been embraced by the public, conservationists, land managers and policy-makers, as well as by scientists’. They postulate that eradication

Corresponding author: Lövei, G.L. (gabor.lovei@agrsci.dk)

* Members of the Biological Invasions in Megadiverse Regions Network contributing to this paper were: Gabor L. Lövei, Thomas M. Lewinsohn, Rodolfo Dirzo (Stanford University, Department of Biology, Stanford, CA 94305, USA), Esmat Faki Mohammed Elhassan (Wildlife Research Center, Sudanese Wildlife Society, PO Box 321/18, University of Khartoum, Sudan), Exequiel Ezcurra (UC Institute for Mexico and the United States, University of California, Riverside, CA 92521-0147, USA), Carolina A. de Oliveira Freire (Department of Physiology, Federal University of Paraná, 81531-970, Curitiba, Paraná, Brazil), Fu-Rong Gui (Plant Protection College, Yunnan Agricultural University, Kunming 650201, China), John M. Halley (Department of Biological Applications and Technology, University of Ioannina, 451 10 Ioannina, Greece), Flora Ismail Tibazarwa (Botany Department, University of Dar es Salaam, PO Box 35060, Dar es Salaam, Tanzania), Ming-Xing Jiang (Institute of Insect Sciences, Zhejiang University, Hangzhou 310029, China), Raymond Katebeka (African Union of Conservationists, Botanical Garden, Makerere University, Box 28973, Kampala, Uganda), Jenesis Kinyamario (School of Biological Sciences, University of Nairobi, Chiromo Campus, PO Box 21553, GPO-00100 Nairobi, Kenya), Samuel Kymanywa (Department of Crop Science, Makerere University, Box 7062, Kampala, Uganda), Feng-Quan Liu (Department of Plant Pathology, Nanjing Agricultural University, Nanjing 210095, China), Shu-Sheng Liu (Institute of Insect Sciences, Zhejiang University, Hangzhou 310029, China), Wan-Xue Liu (Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100081, China), Ying-Quan Liu (Institute of Applied Entomology, Zhejiang University, Hangzhou 310029, China), Bao-Rong Lu (Institute of Biodiversity Science, Fudan University, Shanghai 200433, China), Ed O. Minot (Department of Ecology, Massey University, Palmerston North, New Zealand), Sheng Qiang (Weed Research Laboratory, Nanjing Agricultural University, Nanjing 210095, China), Bao-Li Qiu (Department of Entomology, Southern China Agricultural University, Guangzhou 510650, China), Hao Shen (Southern China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China), Jorge Soberon (Biodiversity Institute, The University of Kansas, Lawrence, KS 66045, USA), Edison Ryoiti Sujii (EMBRAPA Recursos Genéticos e Biotecnologia – Cenargen Parque Estação Biológica – PqEB Final W5 Norte 70770.900, Brasília, DF, Brazil), Jian-Wen Tang (Southern China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China), Ahmet Uludag (Iğdir University, Faculty of Agriculture, Iğdir, 76100, Turkey), Jean R.S. Vitule (Department of Environmental Engineering, Federal University of Paraná, 81531-970, Curitiba, Brazil), Fang-Hao Wan (Department of Biological Invasions, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100081, China), Yuan-Chao Wang (Department of Plant Protection, College of Plant Protection, Nanjing Agricultural University, Nanjing 210095, China), Guo-Qing Yang (Yangzhou University, Yangzhou, Jiangsu 225009, China), Xing-Yao Zhang (Chinese Academy of Forestry, Waishou Shan, Beijing 100091, China), Min-Sheng You (Institute of Applied Ecology, Fujian Agriculture and Forestry University, Fuzhou, Fujian 350002, China).

attempts are mostly a waste of time and money, that many introductions are ecologically beneficial and that the alarm raised about invasive organisms has been exacerbated by unsubstantiated nativism. Reactions so far [2,3] have lacked the voice of developing-country scientists. As a group of scientists from developing countries and/or with extensive field experience in those countries, we seek to change this.

Davis *et al.*'s [1] views are not only imprecise, but in the case of megadiverse developing countries, are also potentially damaging. They maintain that only 'some' species caused extinctions and 'many' of the claims of harm caused by invasive organisms are not backed by data. These qualifiers should be reversed. Many species caused profound, and well-documented, negative effects on native organisms and ecosystems [4], whereas, in some cases, the expected effects were not as drastic as originally predicted.

International transportation and trade have intensified the movement of non-native species [5]. Although all countries face a growing threat from biological invasions, the challenge for many developing countries in blocking the advance of these invaders can be especially formidable, particularly when these countries are undergoing intensive economic development [6]. This often leads to a pernicious false choice of 'economy or ecology'. Additionally, effective management strategies for non-native species in developing countries would be more beneficial, because they have the high biodiversity that is key to future benefits world-wide [7].

Although the effect of both non-native and native species can vary with time, the two should not be conflated. Non-natives turn problematic more frequently than do natives: 49% of the alien insects established in Europe have a negative economic or health impact, whereas <5% of native insects reach pest status in temperate countries [8]. Invasions often lead to decreased biodiversity and faunal homogenization. Thus, they are rightly recognized as an important component of global change [9], and constitute a serious threat to biodiversity, especially in megadiverse (often also developing) tropical countries [7]. The 'invasion debt' [5] could exacerbate the problem, as human-mediated non-native species introduction and native species extinction processes act on different temporal scales.

In some cases, attempts to eradicate introduced species will be useless, but, apart from a steady development in eradication methodology [10], this is an insufficient and inadequate ground for a general change of worldwide policy. In developing countries, the 'embrace the invasives' policy could be used to justify hasty introductions, for instance, in aquaculture. There are examples of introduced fish causing limnological perturbations, introduction of diseases and parasites, or hybridization with native spe-

cies, with negative consequences for fisheries and the local economy [11].

The ethical dimension is also significant. Humans have profound links to, and a psychological need for, nature, subsumed in the 'cultural services' category of ecosystem services. It matters deeply to humans what kind of environment they live in, and it does (and should) matter to them what kind of species populate that environment. This motivated many early attempts to translocate species, and this motive remains valid when considering the desirability, or not, of non-native species. Even if a cow were the full ecological equivalent of an elephant, to an African they are both integral parts of his/her environment, although there is a clearly perceived and well-justified difference. Recently, Didham [12] raised the alarm about the creeping change in mentality of conservation goals, calling attention to the importance of the ethical dimension of biodiversity conservation.

If the real danger of species introductions is minimized because of insufficient and unreliable data, or misguided ecological pragmatism, *carte blanche* is in fact provided to decision-makers or organizations who think only about immediate profit that introductions can bring, without considering longer term losses. Those losses will be tremendous and experienced by all future generations, especially those who live in megadiverse tropical countries.

References

- 1 Davis, M.A. *et al.* (2011) Do not judge species on their origin. *Nature* 474, 153–154
- 2 Simberloff, D. *et al.* (2011) Non-natives: 141 scientists object. *Nature* 475, 36
- 3 Lambertini, M. *et al.* (2011) Invasives: a major conservation threat. *Science* 333, 404–405
- 4 Powell, K.I. *et al.* (2011) A synthesis of plant invasion effects on biodiversity across spatial scales. *Am. J. Bot.* 98, 539–548
- 5 Essl, F. *et al.* (2011) Socioeconomic legacy yields an invasion debt. *Proc. Natl. Acad. Sci. U.S.A.* 108, 203–207
- 6 Lin, W. *et al.* (2007) Fast economic development accelerates biological invasions in China. *PLoS ONE* 2, e1208
- 7 Nuñez, M.A. and Pauchard, A. (2010) Biological invasions in developing and developed countries: does one model fit all? *Biol. Invasions* 12, 707–714
- 8 Kenis, M. *et al.* (2007) How can alien species inventories and interception data help us prevent insect invasions? *Bull. Entomol. Res.* 97, 489–502
- 9 Vitousek, P.M. *et al.* (1997) Introduced species: a significant component of human-caused global change. *N. Z. J. Ecol.* 21, 1–16
- 10 Howald, G. *et al.* (2007) Invasive rodent eradication on islands. *Conserv. Biol.* 21, 1258–1268
- 11 Vitule, J.R.S. *et al.* (2009) Introduction of non-native freshwater fish can certainly be bad. *Fish Fisheries* 10, 98–108
- 12 Didham, R. (2011) Life after logging: strategic withdrawal from the Garden of Eden or tactical error for wilderness conservation? *Biotropica* 43, 393–395