

# Nontrivial solutions for a a class of singular problems

Marcelo Montenegro  
Universidade Estadual de Campinas

Elves A. B. Silva  
Universidade de Brasília

# Introduction

$$\begin{cases} -\Delta u = \left(-u^{-\beta} + \lambda u^p\right) \chi_{\{u>0\}} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (1)$$

$$\begin{cases} -\Delta u = \left(-u^{-\beta} + \lambda u^p\right) \chi_{\{u>0\}} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (1)$$

- ▶  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^n$ ,  $n \geq 1$

$$\begin{cases} -\Delta u = \left(-u^{-\beta} + \lambda u^p\right) \chi_{\{u>0\}} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (1)$$

- ▶  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^n$ ,  $n \geq 1$
- ▶  $0 < \beta < 1$ ,  $0 < p < 1$ ,  $\lambda > 0$

$$\begin{cases} -\Delta u = \left(-u^{-\beta} + \lambda u^p\right) \chi_{\{u>0\}} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (1)$$

- ▶  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^n$ ,  $n \geq 1$
- ▶  $0 < \beta < 1$ ,  $0 < p < 1$ ,  $\lambda > 0$
- ▶ This problem has been studied in several articles: Choi- Lazer - McKenna, Cîrstea - Ghergu - Radulescu, Dávila, Dávila - Montenegro, Diaz - Morel - Oswald

$$\begin{cases} -\Delta u = \left(-u^{-\beta} + \lambda u^p\right) \chi_{\{u>0\}} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (1)$$

- ▶  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^n$ ,  $n \geq 1$
- ▶  $0 < \beta < 1$ ,  $0 < p < 1$ ,  $\lambda > 0$
- ▶ This problem has been studied in several articles: Choi- Lazer - McKenna, Cîrstea - Ghergu - Radulescu, Dávila, Dávila - Montenegro, Diaz - Morel - Oswald
- ▶ By a solution we mean a function  $u \in H_0^1(\Omega)$  satisfying (1) in the weak sense, that is,

$$\begin{cases} -\Delta u = (-u^{-\beta} + \lambda u^p) \chi_{\{u>0\}} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (1)$$

- ▶  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^n$ ,  $n \geq 1$
- ▶  $0 < \beta < 1$ ,  $0 < p < 1$ ,  $\lambda > 0$
- ▶ This problem has been studied in several articles: Choi- Lazer - McKenna, Cîrstea - Ghergu - Radulescu, Dávila, Dávila - Montenegro, Diaz - Morel - Oswald
- ▶ By a solution we mean a function  $u \in H_0^1(\Omega)$  satisfying (1) in the weak sense, that is,

$$\int_{\Omega} \nabla u \nabla \varphi = \int_{\{u>0\}} \left( -\frac{1}{u^{\beta}} + \lambda u^p \right) \varphi$$

for every  $\varphi \in C_c^1(\Omega)$ .



- ▶ There are a few recent papers using variational methods for studying an equation with a positive singular nonlinearity on the right hand side

- ▶ There are a few recent papers using variational methods for studying an equation with a positive singular nonlinearity on the right hand side

$$-\Delta u = \frac{1}{u^\beta} + \lambda u^p,$$

- ▶ There are a few recent papers using variational methods for studying an equation with a positive singular nonlinearity on the right hand side

$$-\Delta u = \frac{1}{u^\beta} + \lambda u^p,$$

- ▶ Canino, Canino - Degiovanni, Hirano - Saccon - Shioji, Perera - —, Long - Sun - Wu, Gonçalves - Santos

- ▶ There are a few recent papers using variational methods for studying an equation with a positive singular nonlinearity on the right hand side

$$-\Delta u = \frac{1}{u^\beta} + \lambda u^p,$$

- ▶ Canino, Canino - Degiovanni, Hirano - Saccon - Shioji, Perera - —, Long - Sun - Wu, Gonçalves - Santos
- ▶ Boccardo for related results

- ▶ There are a few recent papers using variational methods for studying an equation with a positive singular nonlinearity on the right hand side

$$-\Delta u = \frac{1}{u^\beta} + \lambda u^p,$$

- ▶ Canino, Canino - Degiovanni, Hirano - Saccon - Shioji, Perera - —, Long - Sun - Wu, Gonçalves - Santos
- ▶ Boccardo for related results
- ▶ Crandall - Rabinowitz - Tartar: the existence of one solution via bifurcation theory

- ▶ There are a few recent papers using variational methods for studying an equation with a positive singular nonlinearity on the right hand side

$$-\Delta u = \frac{1}{u^\beta} + \lambda u^p,$$

- ▶ Canino, Canino - Degiovanni, Hirano - Saccon - Shioji, Perera - —, Long - Sun - Wu, Gonçalves - Santos
- ▶ Boccardo for related results
- ▶ Crandall - Rabinowitz - Tartar: the existence of one solution via bifurcation theory

# Existence of two solutions

## Theorem 1

*Problem (1) has two distinct nontrivial solutions for  $\lambda > 0$  large.*

# Perturbed Problem



# Perturbed Problem

Consider the perturbation

$$g_\varepsilon(t) = \begin{cases} \frac{t^q}{(t + \varepsilon)^{q+\beta}} & \text{for } t \geq 0 \\ 0 & \text{for } t < 0, \end{cases} \quad (2)$$

# Perturbed Problem

Consider the perturbation

$$g_\varepsilon(t) = \begin{cases} \frac{t^q}{(t + \varepsilon)^{q+\beta}} & \text{for } t \geq 0 \\ 0 & \text{for } t < 0, \end{cases} \quad (2)$$

where  $0 < q < p < 1$

# Perturbed Problem

Consider the perturbation

$$g_\varepsilon(t) = \begin{cases} \frac{t^q}{(t + \varepsilon)^{q+\beta}} & \text{for } t \geq 0 \\ 0 & \text{for } t < 0, \end{cases} \quad (2)$$

where  $0 < q < p < 1$  and the corresponding perturbed problem

$$\begin{cases} -\Delta u + g_\varepsilon(u) = \lambda u^p & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (3)$$

# Perturbed Problem

Consider the perturbation

$$g_\varepsilon(t) = \begin{cases} \frac{t^q}{(t + \varepsilon)^{q+\beta}} & \text{for } t \geq 0 \\ 0 & \text{for } t < 0, \end{cases} \quad (2)$$

where  $0 < q < p < 1$  and the corresponding perturbed problem

$$\begin{cases} -\Delta u + g_\varepsilon(u) = \lambda u^p & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (3)$$

The associated functional  $I_\varepsilon \in C^1(H_0^1(\Omega), \mathbb{R})$  is given by

# Perturbed Problem

Consider the perturbation

$$g_\varepsilon(t) = \begin{cases} \frac{t^q}{(t + \varepsilon)^{q+\beta}} & \text{for } t \geq 0 \\ 0 & \text{for } t < 0, \end{cases} \quad (2)$$

where  $0 < q < p < 1$  and the corresponding perturbed problem

$$\begin{cases} -\Delta u + g_\varepsilon(u) = \lambda u^p & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (3)$$

The associated functional  $I_\varepsilon \in C^1(H_0^1(\Omega), \mathbb{R})$  is given by

$$I_\varepsilon(u) = \frac{1}{2} \int_\Omega |\nabla u|^2 + \int_\Omega G_\varepsilon(u) - \frac{\lambda}{p+1} \int_\Omega (u^+)^{p+1}$$

where  $G_\varepsilon(u) = \int_0^t g_\varepsilon(s) ds \geq 0$ .

# Two solutions for the perturbed problem

# Two solutions for the perturbed problem

## Lemma 1

*For every  $\lambda > 0$ , there is  $\rho > 0$  such that,  $I_\varepsilon(u) \geq \frac{1}{4}\rho^2$  whenever  $\|u\|_{H_0^1} = \rho$  and  $0 < \varepsilon < 1$ .*

# Two solutions for the perturbed problem

## Lemma 1

*For every  $\lambda > 0$ , there is  $\rho > 0$  such that,  $I_\varepsilon(u) \geq \frac{1}{4}\rho^2$  whenever  $\|u\|_{H_0^1} = \rho$  and  $0 < \varepsilon < 1$ .*

## Lemma 2

*For  $\lambda > 0$  large enough, we have  $I_\varepsilon(\varphi_1) < b < 0$ .*



# Two solutions for the perturbed problem

## Lemma 1

*For every  $\lambda > 0$ , there is  $\rho > 0$  such that,  $I_\varepsilon(u) \geq \frac{1}{4}\rho^2$  whenever  $\|u\|_{H_0^1} = \rho$  and  $0 < \varepsilon < 1$ .*

## Lemma 2

*For  $\lambda > 0$  large enough, we have  $I_\varepsilon(\varphi_1) < b < 0$ .*

Here,  $\varphi_1 > 0$  is the first eigenfunction of  $-\Delta$  in  $H_0^1(\Omega)$ .

# Two solutions for the perturbed problem

## Lemma 1

*For every  $\lambda > 0$ , there is  $\rho > 0$  such that,  $I_\varepsilon(u) \geq \frac{1}{4}\rho^2$  whenever  $\|u\|_{H_0^1} = \rho$  and  $0 < \varepsilon < 1$ .*

## Lemma 2

*For  $\lambda > 0$  large enough, we have  $I_\varepsilon(\varphi_1) < b < 0$ .*

Here,  $\varphi_1 > 0$  is the first eigenfunction of  $-\Delta$  in  $H_0^1(\Omega)$ .

## Proposition 1

*For  $\lambda > 0$  large enough, and  $0 < \varepsilon < 1$ ; there is  $b < 0$  and a global minimizer  $u_\varepsilon^1 \in H_0^1(\Omega)$  with  $I_\varepsilon(u_\varepsilon^1) < b$ .*

## Proposition 1

*For  $\lambda > 0$  large enough, and  $0 < \varepsilon < 1$ ; there is  $b < 0$  and a global minimizer  $u_\varepsilon^1 \in H_0^1(\Omega)$  with  $I_\varepsilon(u_\varepsilon^1) < b$ .*

## Proposition 2

*For  $\lambda > 0$  large enough, and  $0 < \varepsilon < 1$ ; there is  $a > 0$  and a critical point  $u_\varepsilon^2 \in H_0^1(\Omega)$  of mountain pass type such that  $I_\varepsilon(u_\varepsilon^2) > a$ .*

# Concluding Remarks for the Perturbed Problem

Fix  $\lambda > 0$  sufficiently large.

# Concluding Remarks for the Perturbed Problem

Fix  $\lambda > 0$  sufficiently large.

The critical values  $c_\varepsilon^1 = I_\varepsilon(u_\varepsilon^1)$  and  $c_\varepsilon^2 = I_\varepsilon(u_\varepsilon^2)$  satisfy

$$-\infty < \beta \leq c_1^\varepsilon \leq b < 0 < a \leq c_2^\varepsilon \leq \alpha < \infty \quad (4)$$

## Concluding Remarks for the Perturbed Problem

Fix  $\lambda > 0$  sufficiently large.

The critical values  $c_\varepsilon^1 = I_\varepsilon(u_\varepsilon^1)$  and  $c_\varepsilon^2 = I_\varepsilon(u_\varepsilon^2)$  satisfy

$$-\infty < \beta \leq c_1^\varepsilon \leq b < 0 < a \leq c_2^\varepsilon \leq \alpha < \infty \quad (4)$$

where the constants  $\beta$ ,  $b$ ,  $a$  do not depend on  $0 < \varepsilon < 1$ .

## Concluding Remarks for the Perturbed Problem

Fix  $\lambda > 0$  sufficiently large.

The critical values  $c_\varepsilon^1 = I_\varepsilon(u_\varepsilon^1)$  and  $c_\varepsilon^2 = I_\varepsilon(u_\varepsilon^2)$  satisfy

$$-\infty < \beta \leq c_1^\varepsilon \leq b < 0 < a \leq c_2^\varepsilon \leq \alpha < \infty \quad (4)$$

where the constants  $\beta$ ,  $b$ ,  $a$  do not depend on  $0 < \varepsilon < 1$ .

Claim: the upper bound  $\alpha$  is independent of  $\varepsilon > 0$ .



# Concluding Remarks for the Perturbed Problem

Fix  $\lambda > 0$  sufficiently large.

The critical values  $c_\varepsilon^1 = I_\varepsilon(u_\varepsilon^1)$  and  $c_\varepsilon^2 = I_\varepsilon(u_\varepsilon^2)$  satisfy

$$-\infty < \beta \leq c_1^\varepsilon \leq b < 0 < a \leq c_2^\varepsilon \leq \alpha < \infty \quad (4)$$

where the constants  $\beta$ ,  $b$ ,  $a$  do not depend on  $0 < \varepsilon < 1$ .

Claim: the upper bound  $\alpha$  is independent of  $\varepsilon > 0$ .

Observe that the solutions  $u_\varepsilon$  of (3) are a priori bounded:

# Concluding Remarks for the Perturbed Problem

Fix  $\lambda > 0$  sufficiently large.

The critical values  $c_\varepsilon^1 = I_\varepsilon(u_\varepsilon^1)$  and  $c_\varepsilon^2 = I_\varepsilon(u_\varepsilon^2)$  satisfy

$$-\infty < \beta \leq c_1^\varepsilon \leq b < 0 < a \leq c_2^\varepsilon \leq \alpha < \infty \quad (4)$$

where the constants  $\beta$ ,  $b$ ,  $a$  do not depend on  $0 < \varepsilon < 1$ .

Claim: the upper bound  $\alpha$  is independent of  $\varepsilon > 0$ .

Observe that the solutions  $u_\varepsilon$  of (3) are a priori bounded:  
multiply (3) by  $u_\varepsilon$ , integrate, discard the term involving  $g_\varepsilon$  and use the Sobolev imbedding, to obtain

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ .

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ . The claim is proved.

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ . The claim is proved.

Now, let  $\varepsilon \rightarrow 0$ .

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ . The claim is proved.

Now, let  $\varepsilon \rightarrow 0$ .

Then  $c_{\varepsilon}^1 \rightarrow c_1$ ,  $c_{\varepsilon}^2 \rightarrow c_2$  with

$$\beta \leq c_1 \leq b < 0 < a \leq c_2 \leq \alpha,$$



$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ . The claim is proved.

Now, let  $\varepsilon \rightarrow 0$ .

Then  $c_{\varepsilon}^1 \rightarrow c_1$ ,  $c_{\varepsilon}^2 \rightarrow c_2$  with

$$\beta \leq c_1 \leq b < 0 < a \leq c_2 \leq \alpha,$$

Moreover,

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ . The claim is proved.

Now, let  $\varepsilon \rightarrow 0$ .

Then  $c_{\varepsilon}^1 \rightarrow c_1$ ,  $c_{\varepsilon}^2 \rightarrow c_2$  with

$$\beta \leq c_1 \leq b < 0 < a \leq c_2 \leq \alpha,$$

Moreover,  $u_{\varepsilon}^1 \rightarrow u^1$  and  $u_{\varepsilon}^2 \rightarrow u^2$  a.e., since solutions of (3) are a priori bounded.

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ . The claim is proved.

Now, let  $\varepsilon \rightarrow 0$ .

Then  $c_{\varepsilon}^1 \rightarrow c_1$ ,  $c_{\varepsilon}^2 \rightarrow c_2$  with

$$\beta \leq c_1 \leq b < 0 < a \leq c_2 \leq \alpha,$$

Moreover,  $u_{\varepsilon}^1 \rightarrow u^1$  and  $u_{\varepsilon}^2 \rightarrow u^2$  a.e., since solutions of (3) are a priori bounded. We also have,  $u^1 \neq u^2$  a.e. and they are nontrivial and nonnegative.

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ . The claim is proved.

Now, let  $\varepsilon \rightarrow 0$ .

Then  $c_{\varepsilon}^1 \rightarrow c_1$ ,  $c_{\varepsilon}^2 \rightarrow c_2$  with

$$\beta \leq c_1 \leq b < 0 < a \leq c_2 \leq \alpha,$$

Moreover,  $u_{\varepsilon}^1 \rightarrow u^1$  and  $u_{\varepsilon}^2 \rightarrow u^2$  a.e., since solutions of (3) are a priori bounded. We also have,  $u^1 \neq u^2$  a.e. and they are nontrivial and nonnegative.

Our objective is to obtain gradient estimates for solutions of (3).

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ . The claim is proved.

Now, let  $\varepsilon \rightarrow 0$ .

Then  $c_{\varepsilon}^1 \rightarrow c_1$ ,  $c_{\varepsilon}^2 \rightarrow c_2$  with

$$\beta \leq c_1 \leq b < 0 < a \leq c_2 \leq \alpha,$$

Moreover,  $u_{\varepsilon}^1 \rightarrow u^1$  and  $u_{\varepsilon}^2 \rightarrow u^2$  a.e., since solutions of (3) are a priori bounded. We also have,  $u^1 \neq u^2$  a.e. and they are nontrivial and nonnegative.

Our objective is to obtain gradient estimates for solutions of (3).

Then, taking  $\varepsilon \rightarrow 0$ ,

$$c(\Omega) \left( \int_{\Omega} u_{\varepsilon}^{p+1} \right)^{\frac{2}{p+1}} \leq \int_{\Omega} |\nabla u_{\varepsilon}|^2 \leq \lambda \int_{\Omega} u_{\varepsilon}^{p+1}.$$

Since  $0 < p < 1$ , a bootstrap argument implies that the norms  $\|u_{\varepsilon}\|_{H_0^1(\Omega)}$  and  $\|u_{\varepsilon}\|_{L^{\infty}(\Omega)}$  are bounded independent of  $\varepsilon$ .

From this,  $c_2^{\varepsilon} = I_{\varepsilon}(u_{\varepsilon}^2) \leq \alpha$ , with  $\alpha > 0$  independent of  $\varepsilon > 0$ . The claim is proved.

Now, let  $\varepsilon \rightarrow 0$ .

Then  $c_{\varepsilon}^1 \rightarrow c_1$ ,  $c_{\varepsilon}^2 \rightarrow c_2$  with

$$\beta \leq c_1 \leq b < 0 < a \leq c_2 \leq \alpha,$$

Moreover,  $u_{\varepsilon}^1 \rightarrow u^1$  and  $u_{\varepsilon}^2 \rightarrow u^2$  a.e., since solutions of (3) are a priori bounded. We also have,  $u^1 \neq u^2$  a.e. and they are nontrivial and nonnegative.

Our objective is to obtain gradient estimates for solutions of (3).

Then, taking  $\varepsilon \rightarrow 0$ , we show that the functions  $u_1$  and  $u_2$  are solutions of (1).

# Gradient Estimates

# Gradient Estimates

Let the weight  $\psi$  be such that

$\psi \in C^2(\bar{\Omega})$ ,  $\psi > 0$  in  $\Omega$ ,  $\psi = 0$  on  $\partial\Omega$  and  $\frac{|\nabla\psi|^2}{\psi}$  is bounded in  $\Omega$ .



# Gradient Estimates

Let the weight  $\psi$  be such that

$\psi \in C^2(\bar{\Omega})$ ,  $\psi > 0$  in  $\Omega$ ,  $\psi = 0$  on  $\partial\Omega$  and  $\frac{|\nabla\psi|^2}{\psi}$  is bounded in  $\Omega$ .

Observe that  $\psi = \varphi_1^2$  is a possible example.

Let the weight  $\psi$  be such that

$\psi \in C^2(\bar{\Omega})$ ,  $\psi > 0$  in  $\Omega$ ,  $\psi = 0$  on  $\partial\Omega$  and  $\frac{|\nabla\psi|^2}{\psi}$  is bounded in  $\Omega$ .

Observe that  $\psi = \varphi_1^2$  is a possible example.

## Lemma 3

*If  $u_\varepsilon$  is a solution of (3), then there is a constant  $M > 0$  independent of  $\varepsilon$  such that*

$$\psi(x)|\nabla u_\varepsilon(x)|^2 \leq M(u_\varepsilon(x)^{1-\beta} + u_\varepsilon(x)) \quad \forall x \in \Omega,$$

*where  $M$  depends only on  $\Omega$ ,  $N$ ,  $\beta$ ,  $\psi$  and  $\|u_\varepsilon\|_{L^\infty(\Omega)}$ .*

- ▶ The proof of this lemma is based on an argument by Dávila - Montenegro.

- ▶ The proof of this lemma is based on an argument by Dávila - Montenegro.
- ▶ Remark that a nontrivial solution  $u_\varepsilon$  of (3) is nonnegative and belongs to  $C^2(\overline{\Omega})$ . However, we cannot use the maximum principle to ensure that  $u_\varepsilon$  is positive or identically zero, since  $u^{q-1}/(u + \varepsilon)^{q+\beta}$  is singular when  $u \sim 0$ .

- ▶ The proof of this lemma is based on an argument by Dávila - Montenegro.
- ▶ Remark that a nontrivial solution  $u_\varepsilon$  of (3) is nonnegative and belongs to  $C^2(\overline{\Omega})$ . However, we cannot use the maximum principle to ensure that  $u_\varepsilon$  is positive or identically zero, since  $u^{q-1}/(u+\varepsilon)^{q+\beta}$  is singular when  $u \sim 0$ .
- ▶ Consider the functions

$$w = \frac{|\nabla u|^2}{Z(u)}, \quad v = w\psi,$$

where  $Z(u_\varepsilon) = u_\varepsilon^{1-\beta} + u_\varepsilon + \delta$ , with  $\delta > 0$ , in order to have  $Z > 0$ .

- ▶ The proof of this lemma is based on an argument by Dávila - Montenegro.
- ▶ Remark that a nontrivial solution  $u_\varepsilon$  of (3) is nonnegative and belongs to  $C^2(\overline{\Omega})$ . However, we cannot use the maximum principle to ensure that  $u_\varepsilon$  is positive or identically zero, since  $u^{q-1}/(u+\varepsilon)^{q+\beta}$  is singular when  $u \sim 0$ .
- ▶ Consider the functions

$$w = \frac{|\nabla u|^2}{Z(u)}, \quad v = w\psi,$$

where  $Z(u_\varepsilon) = u_\varepsilon^{1-\beta} + u_\varepsilon + \delta$ , with  $\delta > 0$ , in order to have  $Z > 0$ . In the end of the proof, we let  $\delta \rightarrow 0$ .

- ▶ The proof of this lemma is based on an argument by Dávila - Montenegro.
- ▶ Remark that a nontrivial solution  $u_\varepsilon$  of (3) is nonnegative and belongs to  $C^2(\overline{\Omega})$ . However, we cannot use the maximum principle to ensure that  $u_\varepsilon$  is positive or identically zero, since  $u^{q-1}/(u + \varepsilon)^{q+\beta}$  is singular when  $u \sim 0$ .
- ▶ Consider the functions

$$w = \frac{|\nabla u|^2}{Z(u)}, \quad v = w\psi,$$

where  $Z(u_\varepsilon) = u_\varepsilon^{1-\beta} + u_\varepsilon + \delta$ , with  $\delta > 0$ , in order to have  $Z > 0$ . In the end of the proof, we let  $\delta \rightarrow 0$ .

- ▶ We also use the fact that a nontrivial solution  $u_\varepsilon$  of (3) belongs to  $C^3$  on a neighborhood of every point where it is positive

Next result shows that  $u_\varepsilon$  converges in  $C_{loc}^1$  to some  $u$  which is in  $C_{loc}^{\frac{1-\beta}{1+\beta}}$ .



Next result shows that  $u_\varepsilon$  converges in  $C_{loc}^1$  to some  $u$  which is in  $C_{loc}^{\frac{1-\beta}{1+\beta}}$ .

#### Lemma 4

For any  $\Omega' \subset \Omega$  there exists  $C$  such that

$$|\nabla u_\varepsilon(x) - \nabla u_\varepsilon(y)| \leq C|x - y|^{\frac{1-\beta}{1+\beta}} \quad \forall x, y \in \Omega'.$$

The constant  $C$  depends only on  $\Omega$ ,  $N$ ,  $\beta$ ,  $p$ ,  $\|u_\varepsilon\|_{L^\infty(\Omega)}$ , but not on  $\varepsilon$ .

Considering  $u$ , a weak limit of solutions  $u_\varepsilon$  of (3),

### Lemma 5

$\frac{1}{u^\beta} \chi_{\Omega_+} \in L^1_{loc}(\Omega)$ , where  $\Omega_+ = \{x \in \Omega : u(x) > 0\}$ .

Considering  $u$ , a weak limit of solutions  $u_\varepsilon$  of (3),

### Lemma 5

$\frac{1}{u^\beta} \chi_{\Omega_+} \in L^1_{loc}(\Omega)$ , where  $\Omega_+ = \{x \in \Omega : u(x) > 0\}$ .

- ▶ The proof is done by choosing appropriate test functions for the perturbed problem.

# Proof of Theorem 1

# Proof of Theorem 1

Let  $\eta \in C^\infty(\mathbb{R})$ ,  $0 \leq \eta \leq 1$ ,  $\eta(s) = 0$  for  $s \leq 1/2$ ,  $\eta(s) = 1$  for  $s \geq 1$ .

# Proof of Theorem 1

Let  $\eta \in C^\infty(\mathbb{R})$ ,  $0 \leq \eta \leq 1$ ,  $\eta(s) = 0$  for  $s \leq 1/2$ ,  $\eta(s) = 1$  for  $s \geq 1$ .

Given  $\varphi \in C_c^1(\Omega)$ ,

# Proof of Theorem 1

Let  $\eta \in C^\infty(\mathbb{R})$ ,  $0 \leq \eta \leq 1$ ,  $\eta(s) = 0$  for  $s \leq 1/2$ ,  $\eta(s) = 1$  for  $s \geq 1$ .

Given  $\varphi \in C_c^1(\Omega)$ , for  $m > 0$ , we have

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) = \int_{\hat{\Omega}} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m), \quad (5)$$

# Proof of Theorem 1

Let  $\eta \in C^\infty(\mathbb{R})$ ,  $0 \leq \eta \leq 1$ ,  $\eta(s) = 0$  for  $s \leq 1/2$ ,  $\eta(s) = 1$  for  $s \geq 1$ .

Given  $\varphi \in C_c^1(\Omega)$ , for  $m > 0$ , we have

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) = \int_{\hat{\Omega}} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m), \quad (5)$$

where  $\hat{\Omega}$  is an open set such that  $\overline{\hat{\Omega}} \subset \Omega$  and  $\text{support}(\varphi) \subset \hat{\Omega}$ .



# Proof of Theorem 1

Let  $\eta \in C^\infty(\mathbb{R})$ ,  $0 \leq \eta \leq 1$ ,  $\eta(s) = 0$  for  $s \leq 1/2$ ,  $\eta(s) = 1$  for  $s \geq 1$ .

Given  $\varphi \in C_c^1(\Omega)$ , for  $m > 0$ , we have

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) = \int_{\hat{\Omega}} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m), \quad (5)$$

where  $\hat{\Omega}$  is an open set such that  $\overline{\hat{\Omega}} \subset \Omega$  and  $\text{support}(\varphi) \subset \hat{\Omega}$ .  
Set  $\Omega_0 = \Omega_+ \cap \hat{\Omega}$ .

# Proof of Theorem 1

Let  $\eta \in C^\infty(\mathbb{R})$ ,  $0 \leq \eta \leq 1$ ,  $\eta(s) = 0$  for  $s \leq 1/2$ ,  $\eta(s) = 1$  for  $s \geq 1$ .

Given  $\varphi \in C_c^1(\Omega)$ , for  $m > 0$ , we have

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) = \int_{\hat{\Omega}} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m), \quad (5)$$

where  $\hat{\Omega}$  is an open set such that  $\overline{\hat{\Omega}} \subset \Omega$  and  $\text{support}(\varphi) \subset \hat{\Omega}$ .

Set  $\Omega_0 = \Omega_+ \cap \hat{\Omega}$ .

Since  $u_\varepsilon \rightarrow u$  in  $C_{loc}^1(\Omega)$ ,

# Proof of Theorem 1

Let  $\eta \in C^\infty(\mathbb{R})$ ,  $0 \leq \eta \leq 1$ ,  $\eta(s) = 0$  for  $s \leq 1/2$ ,  $\eta(s) = 1$  for  $s \geq 1$ .

Given  $\varphi \in C_c^1(\Omega)$ , for  $m > 0$ , we have

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) = \int_{\hat{\Omega}} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m), \quad (5)$$

where  $\hat{\Omega}$  is an open set such that  $\overline{\hat{\Omega}} \subset \Omega$  and  $\text{support}(\varphi) \subset \hat{\Omega}$ .

Set  $\Omega_0 = \Omega_+ \cap \hat{\Omega}$ .

Since  $u_\varepsilon \rightarrow u$  in  $C_{loc}^1(\Omega)$ , for every given  $m > 0$ , there is an  $\varepsilon_0 > 0$  such that

$$u_\varepsilon(x) \leq m/2, \quad \forall x \in \Omega_0 \setminus \Omega_+ \text{ and } 0 < \varepsilon \leq \varepsilon_0. \quad (6)$$

# Proof of Theorem 1

Let  $\eta \in C^\infty(\mathbb{R})$ ,  $0 \leq \eta \leq 1$ ,  $\eta(s) = 0$  for  $s \leq 1/2$ ,  $\eta(s) = 1$  for  $s \geq 1$ .

Given  $\varphi \in C_c^1(\Omega)$ , for  $m > 0$ , we have

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) = \int_{\hat{\Omega}} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m), \quad (5)$$

where  $\hat{\Omega}$  is an open set such that  $\overline{\hat{\Omega}} \subset \Omega$  and  $\text{support}(\varphi) \subset \hat{\Omega}$ .

Set  $\Omega_0 = \Omega_+ \cap \hat{\Omega}$ .

Since  $u_\varepsilon \rightarrow u$  in  $C_{loc}^1(\Omega)$ , for every given  $m > 0$ , there is an  $\varepsilon_0 > 0$  such that

$$u_\varepsilon(x) \leq m/2, \quad \forall x \in \Omega_0 \setminus \Omega_+ \text{ and } 0 < \varepsilon \leq \varepsilon_0. \quad (6)$$

Taking  $0 < \varepsilon < \varepsilon_0$ ,

Taking  $0 < \varepsilon < \varepsilon_0$ , We split the previous integral as

$$A_\varepsilon := \int_{\Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m)$$

and

$$B_\varepsilon := \int_{\hat{\Omega} \setminus \Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m).$$

Taking  $0 < \varepsilon < \varepsilon_0$ , We split the previous integral as

$$A_\varepsilon := \int_{\Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m)$$

and

$$B_\varepsilon := \int_{\hat{\Omega} \setminus \Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m).$$

Clearly,  $B_\varepsilon = 0$  by (6) and the definition of  $\eta$ .

Taking  $0 < \varepsilon < \varepsilon_0$ , We split the previous integral as

$$A_\varepsilon := \int_{\Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m)$$

and

$$B_\varepsilon := \int_{\hat{\Omega} \setminus \Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m).$$

Clearly,  $B_\varepsilon = 0$  by (6) and the definition of  $\eta$ . Moreover,

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \eta(u/m) \quad \text{as } \varepsilon \rightarrow 0.$$



Taking  $0 < \varepsilon < \varepsilon_0$ , We split the previous integral as

$$A_\varepsilon := \int_{\Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m)$$

and

$$B_\varepsilon := \int_{\hat{\Omega} \setminus \Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m).$$

Clearly,  $B_\varepsilon = 0$  by (6) and the definition of  $\eta$ . Moreover,

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \eta(u/m) \quad \text{as } \varepsilon \rightarrow 0.$$

If  $u \leq m/4$ , for a sufficiently small  $\varepsilon > 0$ , we have  $u_\varepsilon \leq m/2$ . The integral  $A_\varepsilon$  restricted to this set is zero.

Taking  $0 < \varepsilon < \varepsilon_0$ , We split the previous integral as

$$A_\varepsilon := \int_{\Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m)$$

and

$$B_\varepsilon := \int_{\hat{\Omega} \setminus \Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m).$$

Clearly,  $B_\varepsilon = 0$  by (6) and the definition of  $\eta$ . Moreover,

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \eta(u/m) \quad \text{as } \varepsilon \rightarrow 0.$$

If  $u \leq m/4$ , for a sufficiently small  $\varepsilon > 0$ , we have  $u_\varepsilon \leq m/2$ . The integral  $A_\varepsilon$  restricted to this set is zero.

If  $u > m/4$ , we have  $u_\varepsilon \geq m/8$  for  $\varepsilon > 0$  small enough.

Taking  $0 < \varepsilon < \varepsilon_0$ , We split the previous integral as

$$A_\varepsilon := \int_{\Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m)$$

and

$$B_\varepsilon := \int_{\hat{\Omega} \setminus \Omega_0} (-g_\varepsilon(u_\varepsilon) + \lambda u_\varepsilon^p) \varphi \eta(u_\varepsilon/m).$$

Clearly,  $B_\varepsilon = 0$  by (6) and the definition of  $\eta$ . Moreover,

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \eta(u/m) \quad \text{as } \varepsilon \rightarrow 0.$$

If  $u \leq m/4$ , for a sufficiently small  $\varepsilon > 0$ , we have  $u_\varepsilon \leq m/2$ . The integral  $A_\varepsilon$  restricted to this set is zero.

If  $u > m/4$ , we have  $u_\varepsilon \geq m/8$  for  $\varepsilon > 0$  small enough. Then, we apply the Dominated Convergence Theorem.

Now we take a second limit as  $m \rightarrow 0$  to conclude that

Now we take a second limit as  $m \rightarrow 0$  to conclude that

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0)$$

Now we take a second limit as  $m \rightarrow 0$  to conclude that

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0)$$

Note that  $\eta(u/m) \leq 1$  and  $-u^{-\beta} + u^p \in L^1(\Omega_0)$ .

Now we take a second limit as  $m \rightarrow 0$  to conclude that

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0)$$

Note that  $\eta(u/m) \leq 1$  and  $-u^{-\beta} + u^p \in L^1(\Omega_0)$ .

Now,

Now we take a second limit as  $m \rightarrow 0$  to conclude that

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0)$$

Note that  $\eta(u/m) \leq 1$  and  $-u^{-\beta} + u^p \in L^1(\Omega_0)$ .

Now, considering the first integral in (5),



Now we take a second limit as  $m \rightarrow 0$  to conclude that

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0)$$

Note that  $\eta(u/m) \leq 1$  and  $-u^{-\beta} + u^p \in L^1(\Omega_0)$ .

Now, considering the first integral in (5), we set

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) := H_\varepsilon + J_\varepsilon.$$

Now we take a second limit as  $m \rightarrow 0$  to conclude that

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0)$$

Note that  $\eta(u/m) \leq 1$  and  $-u^{-\beta} + u^p \in L^1(\Omega_0)$ .

Now, considering the first integral in (5), we set

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) := H_\varepsilon + J_\varepsilon.$$

Clearly,

$$H_\varepsilon := \int_{\Omega} (\nabla u_\varepsilon \nabla \varphi) \eta(u_\varepsilon/m) \rightarrow \int_{\Omega_0} (\nabla u \nabla \varphi) \eta(u/m) \quad \text{as } \varepsilon \rightarrow 0.$$

and

Now we take a second limit as  $m \rightarrow 0$  to conclude that

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0)$$

Note that  $\eta(u/m) \leq 1$  and  $-u^{-\beta} + u^p \in L^1(\Omega_0)$ .

Now, considering the first integral in (5), we set

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) := H_\varepsilon + J_\varepsilon.$$

Clearly,

$$H_\varepsilon := \int_{\Omega} (\nabla u_\varepsilon \nabla \varphi) \eta(u_\varepsilon/m) \rightarrow \int_{\Omega_0} (\nabla u \nabla \varphi) \eta(u/m) \quad \text{as } \varepsilon \rightarrow 0.$$

and

$$\int_{\Omega_0} (\nabla u \nabla \varphi) \eta(u/m) \rightarrow \int_{\Omega_0} \nabla u \nabla \varphi \quad \text{as } m \rightarrow 0,$$

Now we take a second limit as  $m \rightarrow 0$  to conclude that

$$A_\varepsilon \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0)$$

Note that  $\eta(u/m) \leq 1$  and  $-u^{-\beta} + u^p \in L^1(\Omega_0)$ .

Now, considering the first integral in (5), we set

$$\int_{\Omega} \nabla u_\varepsilon \nabla (\varphi \eta(u_\varepsilon/m)) := H_\varepsilon + J_\varepsilon.$$

Clearly,

$$H_\varepsilon := \int_{\Omega} (\nabla u_\varepsilon \nabla \varphi) \eta(u_\varepsilon/m) \rightarrow \int_{\Omega_0} (\nabla u \nabla \varphi) \eta(u/m) \quad \text{as } \varepsilon \rightarrow 0.$$

and

$$\int_{\Omega_0} (\nabla u \nabla \varphi) \eta(u/m) \rightarrow \int_{\Omega_0} \nabla u \nabla \varphi \quad \text{as } m \rightarrow 0,$$

by the Dominated Convergence Theorem.

We assert that

$$J_\varepsilon := \int_{\Omega_0} \frac{|\nabla u_\varepsilon|^2}{m} \eta'(u_\varepsilon/m) \varphi \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0).$$

We assert that

$$J_\varepsilon := \int_{\Omega_0} \frac{|\nabla u_\varepsilon|^2}{m} \eta'(u_\varepsilon/m) \varphi \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0 \quad (\text{and then as } m \rightarrow 0).$$

By the estimate  $|\nabla u_\varepsilon|^2 \leq M(u_\varepsilon^{1-\beta} + u_\varepsilon)$  in  $\Omega_0$  (provided by Lemma 3), we obtain

$$\begin{aligned} |J_\varepsilon| &\leq M \int_{\Omega_0 \cap \{\frac{m}{2} \leq u_\varepsilon \leq m\}} \frac{(u_\varepsilon^{1-\beta} + u_\varepsilon)}{m} \eta'(u_\varepsilon/m) \varphi \rightarrow \\ &\rightarrow M \int_{\Omega_0 \cap \{\frac{m}{2} \leq u \leq m\}} \frac{(u^{1-\beta} + u)}{m} \eta'(u/m) \varphi \quad \text{as } \varepsilon \rightarrow 0, \end{aligned}$$

but this last integral goes to 0 as  $m \rightarrow 0$ .

We have shown that,

We have shown that, as  $\varepsilon \rightarrow 0$  and  $m \rightarrow 0$ ,



We have shown that, as  $\varepsilon \rightarrow 0$  and  $m \rightarrow 0$ ,

$$\int_{\Omega} (-g_{\varepsilon}(u_{\varepsilon}) + \lambda u_{\varepsilon}^p) \varphi \eta(u_{\varepsilon}/m) \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi$$

and

We have shown that, as  $\varepsilon \rightarrow 0$  and  $m \rightarrow 0$ ,

$$\int_{\Omega} (-g_{\varepsilon}(u_{\varepsilon}) + \lambda u_{\varepsilon}^p) \varphi \eta(u_{\varepsilon}/m) \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi$$

and

$$\int_{\Omega} \nabla u_{\varepsilon} \nabla(\varphi \eta(u_{\varepsilon}/m)) \rightarrow \int_{\Omega} \nabla u \nabla \varphi$$

We have shown that, as  $\varepsilon \rightarrow 0$  and  $m \rightarrow 0$ ,

$$\int_{\Omega} (-g_{\varepsilon}(u_{\varepsilon}) + \lambda u_{\varepsilon}^p) \varphi \eta(u_{\varepsilon}/m) \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi$$

and

$$\int_{\Omega} \nabla u_{\varepsilon} \nabla(\varphi \eta(u_{\varepsilon}/m)) \rightarrow \int_{\Omega} \nabla u \nabla \varphi$$

Combining these facts with (5), we obtain

$$\int_{\Omega} \nabla u \nabla \varphi = \int_{\{u>0\}} \left( -\frac{1}{u^{\beta}} + \lambda u^p \right) \varphi$$

for every  $\varphi \in C_c^1(\Omega)$ .

We have shown that, as  $\varepsilon \rightarrow 0$  and  $m \rightarrow 0$ ,

$$\int_{\Omega} (-g_{\varepsilon}(u_{\varepsilon}) + \lambda u_{\varepsilon}^p) \varphi \eta(u_{\varepsilon}/m) \rightarrow \int_{\Omega_0} (-u^{-\beta} + \lambda u^p) \varphi$$

and

$$\int_{\Omega} \nabla u_{\varepsilon} \nabla(\varphi \eta(u_{\varepsilon}/m)) \rightarrow \int_{\Omega} \nabla u \nabla \varphi$$

Combining these facts with (5), we obtain

$$\int_{\Omega} \nabla u \nabla \varphi = \int_{\{u>0\}} \left(-\frac{1}{u^{\beta}} + \lambda u^p\right) \varphi$$

for every  $\varphi \in C_c^1(\Omega)$ .

This concludes the proof of Theorem 1.

# The existence of a positive solution

# The existence of a positive solution

Our second result reads as follows.

# The existence of a positive solution

Our second result reads as follows.

## Theorem 2

*Problem (1) has a positive solution for  $\lambda > 0$  large.*

# The existence of a positive solution

Our second result reads as follows.

## Theorem 2

*Problem (1) has a positive solution for  $\lambda > 0$  large.*

- ▶ We are unable to prove that one of the solutions of Theorem 1 is positive.



# The existence of a positive solution

Our second result reads as follows.

## Theorem 2

*Problem (1) has a positive solution for  $\lambda > 0$  large.*

- ▶ We are unable to prove that one of the solutions of Theorem 1 is positive. We believe that one of them is positive and the other one vanishes somewhere in  $\Omega$ .

# The existence of a positive solution

Our second result reads as follows.

## Theorem 2

*Problem (1) has a positive solution for  $\lambda > 0$  large.*

- ▶ We are unable to prove that one of the solutions of Theorem 1 is positive. We believe that one of them is positive and the other one vanishes somewhere in  $\Omega$ . This would be in agreement with the result for the radial problem proved by Ouyang - Shi - Yao.

- ▶ Theorem 2 is related to a result by Dávila:

- ▶ Theorem 2 is related to a result by Dávila: for  $\lambda$  greater than a precise constant, the maximal solution  $u_\lambda$  is a strict local minimizer  $I$  in the convex subset of  $H_0^1(\Omega)$  of nonnegative functions in  $\Omega$ .

# Proof of Theorem 2

## Proof of Theorem 2

Associated with problem (1) we have the functional  $I : H_0^1(\Omega) \rightarrow \mathbb{R}$  given by

$$I(u) = \int_{\Omega} \frac{1}{2} |\nabla u|^2 - F(u^+),$$

where  $f(u) = -\frac{1}{u^\beta} + \lambda u^p$  and  $F(u) = \int_0^u f(s) ds$ .

It is known ( Dávila-Montenegro) that  $\underline{u} = c\varphi_1^{\frac{2}{1+\beta}}$  is a subsolution (if  $\lambda$  is large) for the problem (1), which in our new notation is

$$\begin{cases} -\Delta u = f(u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases} \quad (7)$$

Take a sequence of smooth domains

$$\emptyset \neq \Omega_1 \subset\subset \Omega_2 \dots \subset\subset \Omega$$

such that  $\Omega = \bigcup_{k=1}^{\infty} \Omega_k$ .

Define the truncated function

$$\hat{f}(u) = \begin{cases} f(\underline{u}(x)) & \text{for } s \leq \underline{u}(x) \\ f(s) & \text{for } s \geq \underline{u}(x) \end{cases} \quad (8)$$

Consider the truncated problems on each domain  $\Omega_k$ ,

$$\begin{cases} -\Delta u_k = \hat{f}(u_k) & \text{in } \Omega_k \\ u_k = \underline{u}(x) & \text{on } \partial\Omega_k. \end{cases} \quad (9)$$

In order to find a solution to (9) we consider the translated problem for  $v_k = u_k - \underline{u}$  with homogeneous boundary conditions

$$\begin{cases} -\Delta v_k = \hat{f}(v_k + \underline{u}) - \Delta \underline{u} & \text{in } \Omega_k \\ v_k = 0 & \text{on } \partial\Omega_k. \end{cases} \quad (10)$$



Define the functional  $\tilde{I}_k : H_0^1(\Omega_k) \rightarrow \mathbb{R}$  by

$$\tilde{I}_k(v) = \int_{\Omega_k} \frac{1}{2} |\nabla v|^2 - \tilde{F}(v) + \nabla \underline{u} \nabla v,$$

here

$$\tilde{F}(v) = \int_0^v \hat{f}(t^+ + \underline{u}) dt.$$

Notice that

$$\tilde{F}(v) = \begin{cases} f(\underline{u}(x))v & \text{for } v \leq 0 \\ \hat{F}(v + \underline{u}) - \hat{F}(\underline{u}) & \text{for } v > 0 \end{cases} \quad (11)$$

where  $\hat{F}(s) = \int_0^s \hat{f}(t) dt$ .

- ▶  $\tilde{I}_k$  is coercive and satisfies the Palais-Smale condition.

- ▶  $\tilde{I}_k$  is coercive and satisfies the Palais-Smale condition.
- ▶ There is  $v_k \in H_0^1(\Omega_k)$  such that

$$\tilde{I}_k(v_k) = \inf_{v \in H_0^1(\Omega_k)} \tilde{I}_k(v).$$

- ▶  $\tilde{I}_k$  is coercive and satisfies the Palais-Smale condition.
- ▶ There is  $v_k \in H_0^1(\Omega_k)$  such that

$$\tilde{I}_k(v_k) = \inf_{v \in H_0^1(\Omega_k)} \tilde{I}_k(v).$$

- ▶  $u_k = v_k + \underline{u}$  is a solution of (9).
- ▶  $v_k \geq 0$  on  $\Omega_k$  ( by the maximum principle since  $\underline{u}$  is a subsolution).
- ▶ Given  $k_0$ ,  $\|v_k\|_{H_0^1(\Omega_{k_0})}$  is bounded for every  $k \geq k_0$ .

Taking a subsequence, we obtain

- ▶  $u_k \rightharpoonup u$  in  $H_0^1(\Omega)$ ,
- ▶  $u_k \rightarrow u$  in  $L^\sigma$  for  $1 \leq \sigma < 2N/(N-2)$ ,
- ▶  $u_k \rightarrow u$  a.e in  $\Omega$ .
- ▶ Hence  $\underline{u} \leq u$  in  $\Omega$ .

Let  $\varphi$  be a test function in  $C_0^\infty(\Omega)$ . There is a  $k' > 0$  and a bounded domain  $\Omega'$  such that  $\text{support}(\varphi) \subset\subset \Omega' \subset\subset \Omega_k$  for every  $k \geq k'$ . Thus,

$$\int_{\Omega'} \nabla u_k \nabla \varphi = \int_{\Omega'} f(u_k) \varphi \quad \text{for every } k \geq k'.$$

Letting  $k \rightarrow \infty$  we obtain

$$\int_{\Omega'} \nabla u \nabla \varphi = \int_{\Omega'} f(u) \varphi.$$

This last integral also holds in  $\Omega$ , so  $u$  is a weak solution.

Let  $\varphi$  be a test function in  $C_0^\infty(\Omega)$ . There is a  $k' > 0$  and a bounded domain  $\Omega'$  such that  $\text{support}(\varphi) \subset\subset \Omega' \subset\subset \Omega_k$  for every  $k \geq k'$ . Thus,

$$\int_{\Omega'} \nabla u_k \nabla \varphi = \int_{\Omega'} f(u_k) \varphi \quad \text{for every } k \geq k'.$$

Letting  $k \rightarrow \infty$  we obtain

$$\int_{\Omega'} \nabla u \nabla \varphi = \int_{\Omega'} f(u) \varphi.$$

This last integral also holds in  $\Omega$ , so  $u$  is a weak solution.