

Joint effects of *CSN3* and *LGB* genes on milk quality and coagulation properties in Czech Fleckvieh

A. MATĚJÍČEK¹, J. MATĚJÍČKOVÁ¹, M. ŠTÍPKOVÁ¹, O. HANUŠ², V. GENČUROVÁ², J. KYSELOVÁ¹, E. NĚMCOVÁ¹, T. KOTT¹, J. ŠEFROVÁ¹, M. KREJČOVÁ¹, S. MELČOVÁ¹, I. HÖLZELOVÁ¹, J. BOUŠKA¹, J. FRELICH³

¹Institute of Animal Science, Prague-Uhřetěves, Czech Republic

²Research Institute for Cattle Breeding, Rapotín, Czech Republic

³University of South Bohemia, České Budějovice, Czech Republic

ABSTRACT: The aim of this study was to determine the joint effects of *CSN3* and *LGB* genotypes on parameters of production, quality and coagulation of milk in Czech Fleckvieh cows. Three hundred and twenty-eight Czech Fleckvieh cows were determined for *CSN3* (kappa-casein) and *LGB* (beta-lactoglobulin) genotypes using the PCR-RFLP method, milk quality parameters and coagulation properties. Milk production parameters were obtained from the Official Database of Progeny Testing. Fifteen genotype combinations were detected, with *ABAB* (21.0%) and *AAAB* (18.3%) occurring as the most frequent. The observed genes significantly affected the contents of milk protein (crude protein, true protein, casein and whey protein) as well as solid non-fat in milk, casein number and curd quality. *BBAA* was found to be the genotype with the highest positive impact on most of the milk characteristics evaluated. Whereas *ABBB*, *BBBB*, *BBAB* and *ABAB* had a positive influence on milk quality and milk coagulation properties, genotypes containing *CSN3* allele *E* had a negative effect. Results presented in this study are applicable in the selection of Czech Fleckvieh cattle.

Keywords: cattle; milk protein genes; kappa-casein; beta-lactoglobulin; milk production parameters; coagulation properties

Milk protein genes, especially kappa-casein (*CSN3*) and beta-lactoglobulin (*LGB*), are important determinants of milk quality and milk coagulation properties. According to Strzalkowska et al. (2002) there are significant interactions between *CSN3* and *LGB* genes, and their influence on milk quality should be evaluated as a joint effect.

Kappa-casein protein, the only casein fraction that contains the sulphur amino acids cysteine and methionine, constitutes approximately 13% of milk casein (Farrell et al., 2004). The kappa-casein gene

(*CSN3*) is situated on bovine chromosome 6, and its polymorphism has been known since 1964 (Neelin, 1964). Eleven alleles have been identified (Farrell et al., 2004) with *A*, *B* and *E* as the most common (Neubauerová, 2001). Whereas the amino acids threonine and aspartic acid are encoded by allele *A* at positions 136 and 148, the corresponding amino acids for allele *B* are isoleucine and alanine (Eigel et al., 1984). At position 155, allele *E* encodes the amino acid glycine, while the amino acid serine commonly occurs at this position (Erhardt, 1989).

Supported by the Ministry of Agriculture of the Czech Republic (Project No. 1G46086) and the Ministry of Education, Youth and Sports of the Czech Republic (Project No. MSM 6007665806).

Allele *A* has been reported as the most frequent, with the effect of increasing milk yield but decreasing protein content (Neubauerová, 2001; Kučerová et al., 2005). Allele *B* is often referred to as a “key allele”, increasing milk protein quality and coagulation properties (Hanuš et al., 1995; Amigo et al., 2001; Comin et al., 2006), whereas a negative effect of allele *E* on milk coagulation properties was reported by Ikonen et al. (1999a) and Comin et al. (2006).

Beta-lactoglobulin is the main protein of whey. The gene for beta-lactoglobulin (*LGB*) is situated on bovine chromosome 11, with its polymorphism discovered in 1955 (Aschaffenburg and Drewry, 1955). A total of 11 alleles are known (Farrell et al., 2004), with alleles *A* and *B* being found the most frequent (Panicke et al., 1996) and alleles *C* and *D* rarely occurring in European cattle breeds. Allele *A* encodes the amino acids aspartic acid and valine at positions 64 and 118. At corresponding positions, allele *B* encodes glycine and alanine (Eigel et al., 1984). Allele *B* was reported to increase milk protein and milk fat contents, while allele *A* increases milk and protein yields (Neubauerová, 2001; Kaminski et al., 2002; Kučerová et al., 2006). In Czech Fleckvieh cows, Hanuš et al. (1995) found a positive effect of the *BB* genotype on milk coagulation properties, whereas higher curd production due to allele *B* was reported by Choi and Ng-Kwai-Hang (2002).

Studies evaluating milk protein genes usually focus either on phenotypic characteristics of cows (Ikonen et al., 2001; Choi and Ng-Kwai-Hang, 2002; Strzalkowska et al., 2002; Comin et al., 2006) or on breeding values of sires (Ron et al., 1994; Sabour et al., 1996; Kaminski et al., 2002; Kučerová et al., 2006). In a previous study (Matějčíček et al., 2007) the joint effects of *CSN3* and *LGB* genotypes on breeding values of sires were evaluated. The aim of the present study was to investigate the joint effects of *CSN3* and *LGB* genotypes on parameters of milk production, milk quality and milk coagulation properties of Czech Fleckvieh cows.

MATERIAL AND METHODS

Kappa-casein (*CSN3*) and beta-lactoglobulin (*LGB*) genotypes were identified in 328 Czech Fleckvieh cows using a Polymerase Chain Reaction and Restriction Fragment Length Polymorphism method (PCR-RFLP) and electrophoresis on agarose gel. DNA was obtained from blood samples of

cows (Kawasaki, 1990). The identification of *CSN3* alleles *A*, *B* and *E* was done according to Lien and Rogne (1993) and of *LGB* alleles *A* and *B* according to Agrawala et al. (1992). The PCR conditions for *CSN3* were 1 min at 95°C, followed by 35 cycles of 30 s at 95°C, 25 s at 62°C, 30 s at 72°C. The protocol for *LGB* was 3 min. at 95°C, followed by 40 cycles of 60 s at 95°C, 30 s at 60°C, and 30 s at 72°C.

Milk samples were collected between 50 and 140 days after first calving and analysed for contents of:

- dry matter (%);
- solid non-fat (%);
- crude protein (%; Kjeldahl total N × 6.38);
- true protein (%; Kjeldahl protein N × 6.38);
- casein (%; Kjeldahl casein N × 6.38);
- whey protein (%) = true protein – casein;
- non-protein nitrogen substances (%) = crude protein – true protein;
- casein number = % of casein from crude protein.

The parameters were analysed according to standardized methods defined by ČSN 57 0530 and ČSN 57 0536 and standard operation procedures of the accredited laboratory in the Research Institute for Cattle Breeding, Rapotín (National Reference Laboratory for Raw Milk; No. 1 340, EN ISO 17025). The fractions such as crude protein, true protein and casein contents were determined on the instrument 2200 Kjeltac Auto Distillation (Foss-Tecator AB, Sweden), contents of dry matter and solid non-fat were measured on the instrument MilkoScan 133B (Foss Electric, Denmark).

Evaluated cheese-making parameters were as follows:

- coagulation time (s) = means the interval (s) from adding Renilase (microbial enzyme) into the milk sample (50 ml, 35°C) until the first visible milk protein coagulation;
- curd quality (graded from 1 = excellent to 4 = poor) = evaluated by examination and palpation of curd cake obtained from 50 ml of milk;
- curd firmness (mm; lower mm = firmer curd) = investigated after 60 min of curd incubation (i.e. 60 min after Renilase addition) by measurement of the penetration depth of cylindrical corpuscle which falls into the curd cake under standard repeated conditions (weight and fall height of the corpuscle);
- whey amount (ml) = amount of whey obtained from coagulation of 50 ml of milk (coagulated under previously described conditions) and measured with a volumetric glass cylinder.

Milk production parameters (milk yield, protein and fat yields, protein and fat contents) were obtained from the Official Database of Progeny Testing.

Frequencies of *CSN3* + *LGB* genotypes were calculated, and the statistical analysis (GLM – general linear model – with fixed effects) was performed by SAS program. The following model was used to evaluate the relation between genotypes and milk production characteristics:

$$y_{ijklmn} = \mu + HYS_i + G_j + C_k + S_l + bA_m + e_{ijklmn} \quad (\text{Model 1})$$

where:

y = observed characteristic

μ = average of the characteristic

HYS = effect of herd, year and season of calving

G = joint effect of *CSN3* and *LGB* genotypes

C = effect of the genetic part of Czech Fleckvieh breed in cows

S = effect of the breeding value of sire

bA_m = effect of the age at first calving of cow

e_{ijklmn} = residual effect

The model used to investigate the relation between genotypes and milk quality parameters and coagulation properties was as follows:

$$y_{ijkl} = \mu + HYS_i + G_j + bA_k + e_{ijkl} \quad (\text{Model 2})$$

where the symbols correspond to the symbols explained in Model 1.

RESULTS AND DISCUSSION

Fifteen genotype combinations of *CSN3* and *LGB* genes were found in Czech Fleckvieh cows (Table 1). Genotype *ABAB* occurred as the most frequent, which is consistent with previous results (Matějček et al., 2007). Whereas a high frequency of the genotype combination *AAAB* was found, a low rate (less than 2%) of genotypes containing the *CSN3* allele *E* occurred. Genotype *BEAA* was identified only in a single cow. The low frequency of allele *E* is beneficial because this allele has a negative effect on milk coagulation properties (Ikonen et al., 1999a; Comin et al., 2006).

The influence of the joint effects of *CSN3* + *LGB* genotypes on milk production parameters during the first 100 and 200 days of lactation is presented in Table 2. A significant gene effect on protein content was found in both lactation periods. The genotype combinations *AEAA* and *AEBB* resulted

Table 1. Genotype frequencies found in cows

<i>CSN3</i> + <i>LGB</i> genotypes	n	Frequency (%)
<i>AAAA</i>	43	13.1
<i>AAAB</i>	60	18.3
<i>AABB</i>	35	10.7
<i>ABAA</i>	33	10.1
<i>ABAB</i>	69	21.0
<i>ABBB</i>	33	10.1
<i>AEAA</i>	2	0.6
<i>AEAB</i>	6	1.8
<i>AEBB</i>	6	1.8
<i>BBAA</i>	5	1.5
<i>BBAB</i>	12	3.7
<i>BBBB</i>	12	3.7
<i>BEAA</i>	1	0.3
<i>BEAB</i>	6	1.8
<i>BEBB</i>	5	1.5

in the highest milk and protein yields. However, it caused a low protein content. With genotype *AEAA* detected in two cows only no conclusions on this genotype are justified. High yield characteristics were also associated with genotypes *BBAA*, *AABB* and *AAAB*. This is in agreement with Ikonen et al. (1999b), who found that *CSN3* alleles *E* and *A* increase milk yield but decrease protein content. The highest protein content was found in the genotype combinations *BBAA*, *ABBB* and *ABAB*, which illustrates the effect of *CSN3* allele *B* on increasing protein content (Ikonen et al., 1999b; Kučerová et al., 2006). Similarly, Kaminski et al. (2002) reported the highest protein content as well as fat content associated with genotype combinations *ABBB* and *ABAB*.

A significant effect of *CSN3* + *LGB* genotypes was found in almost all milk quality parameters (Table 3), except for the contents of dry matter (DM) and non-protein nitrogen substances (NPNS). This effect was highly significant ($P < 0.001$) for the contents of true protein (TP), casein (C) and whey protein (WP), and for casein number (CN). The most favourable results were associated with genotype combinations *BBAA*, *BBBB* and *ABBB*. Genotype *BBAA* excelled in all parameters except for whey protein content and casein number. Genotypes

Table 2. Joint effects of *CSN3* + *LGB* genotypes on milk production parameters during the first 100 and 200 days of lactation ($n = 328$)

Genotype	n	Parameter for the first 100 and 200 days of lactation									
		milk yield (kg)		protein yield (kg)		protein* content(%)		fat yield (kg)		fat content (%)	
		100/200	100/200	100/200	100/200	100/200	100/200	100/200	100/200	100/200	100/200
AAAA	43	2 434	4 739	80	159	3.29	3.37	95	182	3.91	3.87
AAAB	60	2 531	4 926	82	164	3.26	3.35	99	190	3.91	3.88
AABB	35	2 545	4 918	84	167	3.33	3.41	106	197	4.15	4.02
ABAA	33	2 467	4 888	82	164	3.34	3.37	100	190	4.05	3.91
ABAB	69	2 459	4 826	82	165	3.37	3.43	98	188	4.00	3.92
ABBB	33	2 498	4 703	84	163	3.38	3.47	102	183	4.08	3.92
AEAA	2	3 163	6 277	103	202	3.22	3.19	117	242	3.66	3.84
AEAB	6	2 486	4 552	82	154	3.32	3.38	97	177	3.91	3.89
AEBB	6	2 738	5 520	87	181	3.18	3.29	111	218	4.09	3.95
BBAA	5	2 418	4 699	84	169	3.46	3.59	99	189	4.08	4.02
BBAB	12	2 527	5 018	83	170	3.30	3.40	99	189	3.92	3.79
BBBB	12	2 440	4 562	81	155	3.32	3.41	95	170	3.92	3.77
BEAA	1	2 548	4 873	75	152	3.00	3.16	68	147	2.83	3.12
BEAB	6	2 301	4 456	75	148	3.26	3.34	90	169	3.95	3.85
BEBB	5	2 224	4 274	72	145	3.25	3.40	87	170	3.98	4.02

*significance at $P < 0.05$ Table 3. Joint effects of *CSN3* + *LGB* genotypes on milk quality parameters ($n = 328$)

Genotype	n	Milk quality parameter	DM (%)	SNF** (%)	CP** (%)	TP*** (%)	NPNS (%)	C*** (%)	WP*** (%)	CN*** (%)
AAAA	43	13.03	9.05	3.38	3.17	0.21	2.64	0.53	78.05	
AAAB	60	13.02	9.04	3.36	3.17	0.19	2.69	0.48	79.87	
AABB	35	13.14	9.06	3.39	3.20	0.19	2.73	0.49	80.51	
ABAA	33	12.98	8.96	3.32	3.13	0.20	2.59	0.54	77.89	
ABAB	69	13.13	9.11	3.44	3.25	0.20	2.76	0.49	80.23	
ABBB	33	13.35	9.19	3.54	3.34	0.20	2.90	0.43	82.08	
AEAA	2	12.37	9.20	3.32	3.14	0.17	2.58	0.57	77.67	
AEAB	6	12.98	9.08	3.35	3.16	0.19	2.68	0.48	80.05	
AEBB	6	12.96	8.98	3.29	3.10	0.19	2.72	0.37	82.82	
BBAA	5	13.49	9.54	3.78	3.62	0.16	3.07	0.55	81.19	
BBAB	12	13.25	9.07	3.44	3.24	0.20	2.77	0.47	80.40	
BBBB	12	13.01	9.09	3.47	3.27	0.20	2.82	0.45	81.41	
BEAA	1	12.83	9.16	3.19	2.98	0.21	2.53	0.45	79.31	
BEAB	6	13.41	8.95	3.33	3.10	0.24	2.65	0.45	79.32	
BEBB	5	13.19	9.23	3.49	3.24	0.24	2.83	0.41	81.13	

**.*significance at $P < 0.01$ and $P < 0.001$

DM = dry matter; SNF = solid non-fat; CP = crude protein; TP = true protein; NPNS = non-protein nitrogen substances; C = casein; WP = whey protein; CN = casein number

Table 4. Joint effects of *CSN3* + *LGB* genotypes on cheese-making parameters (n = 328)

Genotype	n	Cheese-making parameter			
		coagulation time (s)	curd quality* (grade)	curd firmness (mm)	whey amount (ml)
AAAA	43	126	2.70	1.82	33
AAAB	60	127	2.40	1.78	33
AABB	35	121	2.24	1.82	33
ABAA	33	105	2.24	1.82	34
ABAB	69	110	2.16	1.80	34
ABBB	33	123	1.98	1.77	33
AEAA	2	95	2.16	1.66	35
AEAB	6	181	2.83	1.72	31
AEBB	6	131	2.65	1.88	34
BBAA	5	90	1.55	1.79	32
BBAB	12	100	1.61	1.72	34
BBBB	12	135	2.17	1.68	33
BEAA	1	131	1.06	1.83	35
BEAB	6	100	2.62	1.83	35
BEBB	5	110	2.77	1.88	30

*significance at $P < 0.05$

BBBB and *ABBB* resulted in a high casein content and low whey protein content. Genotypes *AAAA*, *ABAA*, *AEAA*, *AAAB* and *BEAB* were linked with the least favourable results. The results of genotype *BEAA* are low informative due to its presence in only one cow. Hanuš et al. (2000) reported that genotypes *BB* of *CSN3* and *LGB* genes significantly increased casein content and casein number, and genotype *AA* of *LGB* gene increased whey protein content, in milk from Czech Fleckvieh cows.

Regarding cheese-making characteristics, the joint effects of *CSN3* and *LGB* genes were significant only on curd quality (Table 4). As for milk quality and milk production characteristics, the genotype combination *BBAA* had the most favourable influence. Genotype *BBAA* was associated with the shortest coagulation time, and relatively high curd quality and firmness. However, it resulted in a lower amount of eliminated whey relative to other genotypes. Also genotypes *BBAB* and *ABBB* had a positive impact on cheese-making characteristics. Genotype *BBBB* was associated with relatively high curd firmness and good curd quality, but with a long coagulation time. A negative influence on cheese-making characteristics was caused by genotypes *AEAB*, *AEBB*, *BEBB*, *BEAB* and *AAAA*. Although genotype *BEAA* was associated with a relatively

high curd quality, it was identified in one cow only. Results are in agreement with Choi and Ng-Kwai Hang (2002), who found that the genotype combinations *BBBB*, *BBAB*, *BBAA*, *ABBB* and *AABB* positively affected cheese-making parameters, whereas *AAAA* and *AAAB* had a negative influence. Unfortunately, they did not identify *CSN3* allele *E* in their study. On the contrary, Comin et al. (2006) reported that the genotypes containing *CSN3* allele *E* were associated with poor coagulating milk, and genotypes with *CSN3* allele *B* resulted in the best coagulation properties of milk.

CONCLUSION

A strong influence of the *CSN3* gene in combination with the *LGB* gene on milk protein composition and milk coagulation properties was found in the present study. Whereas *CSN3* allele *B* had a positive influence on most characteristics evaluated, allele *E* had a negative impact. The genotype combination *BBAA* was found to be the most favourable. Genotypes *ABBB*, *BBBB*, *BBAB* and *ABAB* had a positive effect on milk quality and coagulation properties, while genotypes containing *CSN3* allele *E* caused a negative response. Findings

presented in this study are applicable in the selection of Czech Fleckvieh cattle.

Acknowledgements

The authors thank the Czech Fleckvieh farms for providing blood samples of cows and also the Czech-Moravian Breeders' Corporation, Ltd. for providing the milk production parameters of cows for the first 100 and 200 days of lactation.

REFERENCES

- Agrawala P.L., Wagner V.A., Geldermann H. (1992): Sex determination and milk protein genotyping of preimplantation stage bovine embryos using multiplex PCR. *Theriogenology*, 38, 969–978.
- Amigo L., Martin-Alvarez P.J., Garcia-Muro E., Zarazaga I. (2001): Effect of milk protein haplotypes on the composition and technological properties of Fleckvieh bovine milk. *Milchwissenschaft*, 56, 488–491.
- Aschaffenburg R., Drewry J. (1955): Occurrence of different beta-lactoglobulins in cow's milk. *Nature*, 176, 218 pp.
- Comin A., Cassandro M., Ojala M., Bittante G. (2006): Effect of β - and κ -casein genotypes on milk coagulation properties, milk production and content, and milk quality traits in Italian Holstein cows. In: 57th Annual Meeting of the EAAP, 17.–20.9. 2006, Antalya, Turkey.
- ČSN 57 0530 (1974): Methods for testing of milk and milk products. Czech Standards Institute, Prague, CR.
- ČSN 57 0536 (1999): Determination of milk composition by mid-infrared analyser. Czech Standards Institute, Prague, CR.
- Eigel W.N., Butler J.E., Ernstrom C.A., Farrell H.M., Harwalkar V.R., Jennes R., Whitney R.McL. (1984): Nomenclature of proteins of cow's milk. *J. Dairy Sci.*, 67, 1599–1631.
- Erhardt G. (1989): κ -caseins in bovine milk. Evidence of a further allele (κ -Cn E) in different breeds. *J. Anim. Breed. Genet.*, 106, 225–231.
- Farrell H.M., Jimenez-Flores R., Bleck G.T., Brown E.M., Butler J.E., Creamer L.K., Hicks C.L., Hollar C.M., Ng-Kwai-Hang K.F., Swaisgood H.E. (2004): Nomenclature of the proteins of cows' milk – sixth revision. *J. Dairy Sci.*, 87, 1641–1674.
- Hanuš O., Gajdůšek S., Gabriel B., Kopecký J., Jedelská R. (1995): Cheese-making properties of raw and pasteurized milk with respect to milk protein polymorphism. *Czech J. Anim. Sci.*, 40, 523–528.
- Hanuš O., Beber K., Kopecký J. (2000): Milk protein variants and characteristics of cows' milk. In: *Breeding, Nutritional and Technological Aspects of Milk Production and Quality*. Rapotín, Czech Republic, 47–49.
- Choi J.W., Ng-Kwai-Hang K.F. (2002): Effects of genetic variants of κ -casein and β -lactoglobulin and heat treatment of milk on cheese and whey compositions. *Asian-Aust. J. Anim. Sci.*, 5, 732–739.
- Ikonen T., Ahlfors K., Kempe R., Ojala M., Ruttinen O. (1999a): Genetic parameters for milk coagulation properties and prevalence of noncoagulating milk in Finnish dairy cows. *J. Dairy Sci.*, 82, 205–214.
- Ikonen T., Ojala M., Ruottinen O. (1999b): Associations between milk protein polymorphism and first lactation milk production traits in Finnish Ayrshire cows. *J. Dairy Sci.*, 82, 1026–1033.
- Ikonen T., Bovenhuis H., Ojala M., Ruottinen O., Georges M. (2001): Associations between casein haplotypes and first lactation milk production traits in Finnish Ayrshire cows. *J. Dairy Sci.*, 84, 507–514.
- Kaminski S., Rymkiewicz-Schymczyk J., Wojcik E., Rusc A. (2002): Associations between bovine milk protein genotypes and haplotypes and the breeding value of Polish Black-and-White bulls. *J. Anim. Feed Sci.*, 11, 205–221.
- Kawasaki E.S. (1990): Sample preparation from blood, cells and another fluids. In: *PCR Protocols: A Guide to Methods and Applications*. Academic Press, New York, USA, 146–152.
- Kučerová J., Němcová E., Štípková M., Jandurová O., Matějčíček A., Bouška J. (2005): The association between CSN3 genotypes and milk production parameters in Czech Pied Cattle. In: 56th Ann. Meet. EAAP, 5.–8.6. 2005, Uppsala, Sweden. Available at http://www.eaap.org/uppsala/Papers/added/G6.15_Jitka.pdf
- Kučerová J., Matějčíček A., Jandurová O. M., Sorensen P., Němcová E., Štípková M., Kott T., Bouška J., Frelich J. (2006): Milk protein genes *CSN1S1*, *CSN2*, *CSN3*, *LGB* and their relation to genetic values of milk production parameters in Czech Fleckvieh. *Czech J. Anim. Sci.*, 51, 241–247.
- Lien S., Rogne S. (1993): Bovine casein haplotypes: number, frequencies and applicability as genetic markers. *Anim. Genet.*, 24, 373–376.
- Matějčíček A., Matějčíčková J., Němcová E., Jandurová O.M., Štípková M., Bouška J., Frelich J. (2007): Joint effects of *CSN3* and *LGB* genotypes and their relation to breeding values of milk production parameters in Czech Fleckvieh. *Czech J. Anim. Sci.*, 52, 83–87.
- Neelin J. M. (1964): Variants of κ -casein revealed by improved starch gel electrophoresis. *J. Dairy Sci.*, 47, 506 pp.

- Neubauerová V. (2001): Detection of genetic markers and possibilities of their use in cattle and other subungulates. Thesis University of South Bohemia, Agricultural Faculty, České Budějovice, CR, 211 pp.
- Panicke L., Freyer G., Erhardt G. (1996): Effekte der Milchproteinpolymorphismen auf die Leistung. In: Kolloquium Milchprotein und Proteinansatz, Graal-Müritz, FBN Dummerstorf, Universität Rostock, Germany, 20 pp.
- Ron M., Yoffe O., Ezra E., Medrano J.F., Weller J.I. (1994): Determination of effects of milk protein genotype on production traits of Israeli Holsteins. *J. Dairy Sci.*, 77, 1106–1113.
- Sabour M.P., Lin C.Y., Lee A.J., McAllister A.J. (1996): Association between milk protein variants and genetic values of Canadian Holstein bulls for milk yield traits. *J. Dairy Sci.*, 79, 1050–1056.
- Strzalkowska N., Krzyzewski J., Zwierzchowski L., Ryniewicz Z. (2002): Effects of κ -casein and β -lactoglobulin loci polymorphism, cows' age, stage of lactation and somatic cell count on daily milk yield and milk composition in Polish Black-and-White cattle. Institute of Genetics and Animal Breeding, Jastrzebiec, Poland. *Anim. Sci. Pap. Rep.*, 20, 21–35.

Received: 2007–10–09

Accepted after corrections: 2008–03–16

Corresponding Author

Ing. Jitka Matějčková, Ph.D., Institute of Animal Science, Přátelství 815, 104 01 Prague-Uhřetěves, Czech Republic
Tel. +420 267 009 500, e-mail: jitka.k@seznam.cz
