

Effects of different levels of dietary supplemental caprylic acid and vitamin E on performance, breast muscle vitamin E and A, and oxidative stability in broilers

M. SKŘIVAN, G. DLOUHÁ, M. ENGLMAIEROVÁ, K. ČERVINKOVÁ

Institute of Animal Science, Prague-Uhřetěves, Czech Republic

ABSTRACT: The paper reports data from four experiments (EXP). Broiler cockerels Ross 308 were allocated to 3 or 5 (EXP 4) dietary treatments comprising 300 (3 replicated pens of 100 chicks per pen), 50, 50 or 100 chickens. The EXP lasted 42, 38, 42 and 38 days. All basal diets contained adequate ingredients. The inclusion of 0.5% caprylic acid (CA) in diets reduced ($P < 0.05$) body weight (EXP 1). The results of EXP 3 and 4 indicated that the diet supplemented with 0.25% CA and 30 mg vitamin E (total vitamin E concentration 50 mg) resulted in similar BW like the basal diet, but the increased supplemental vitamin E to 150 mg (EXP 3) or to 100 mg (EXP 4) decreased ($P < 0.05$ and $P < 0.001$) BW and increased mortality. No differences were found between the fat and crude protein contents in dry matter of breast meat. Higher vitamin E doses in feed mixture significantly ($P < 0.001$) increased the content of vitamins soluble in fats in breast meat – vitamin E (from 28.54 in the control to 80.28 mg/kg of dry matter) and vitamin A (from 0.34 to 0.44 mg/kg of dry matter). The addition of caprylic acid significantly decreased the speed of lipid oxidation measured after 3 ($P < 0.02$) and 5 ($P < 0.05$) days of storage in refrigerator at a temperature from 2.5 to 4°C. On the contrary, higher vitamin E doses and basal diet without CA supplement increased the oxidation of lipids.

Keywords: caprylic acid; vitamin E; meat quality; performance; male broiler

A reduction of pathogenic microorganisms in the digestive tract decreases the contamination of poultry products. It applies e.g. to strains of *Salmonella* or *Campylobacter*, which are one of the main causes of diseases of alimentary origin. Epidemiological studies demonstrate that poultry and poultry products are frequent sources of pathogenic agents. Especially at present when feeding antibiotics is banned, other substances become increasingly important. Fatty acids (FA), above all medium-chain fatty acids (MCFA), also show the antibacterial activity (Bergsson et al., 1998; Van Immerseel et al., 2004). Caprylic acid (CA) is a natural MCFA with eight carbons and occurs in

human and bovine milk (Jensen, 2002) and in coconut oil (Jensen et al., 1990; Sprong et al., 2001). The bacteriostatic and bactericidal effects of FA and their monoglycerides against a wide range of microorganisms have been known for a long time (Hassinen et al., 1951; Nieman, 1954; Kabara et al., 1972). Caprylic acid is active against coliform bacteria and may improve the resistance of weaned rabbits to enterocolitis (Skřivanová et al., 2008). The antimicrobial activity expressed as minimum inhibitory concentration (MIC) varied from 1 to 3 mg/ml (Skřivanová et al., 2006). Its bacterial activity is reduced at pH > 6 (Skřivanová et al., 2005). Both *in vitro* and fattening experiments demon-

Supported by the Ministry of Agriculture of the Czech Republic (Project No. Mze 0002701404).

strated that CA favourably influenced the digestive tract. Dierick et al. (2002) observed that MCEFA (C₆–C₁₂) significantly suppressed the intestinal flora (total anaerobic count, lactobacilli, *E. coli*) and improve the mucosal health and growth performance of piglets. The addition of caprylic acid (5 g/kg) to a diet for piglets increased ($P < 0.05$) weight gain compared to basal diet (Marounek et al., 2004). The supplementation of CA (0.35, 0.525, 0.7 and 0.875%) to feed significantly ($P < 0.05$) reduced the caecal *Campylobacter jejuni* colonization in young chickens. At higher doses, CA reduced feed consumption and body weight, but it did not affect feed conversion (Solis de los Santos et al., 2008). The supplementation of vitamin E tended to improve the growth and feed utilization of birds and it substantially improved the meat quality stability against oxidative deterioration (Guo et al., 2001). Whereas Attia et al. (2001) noted statistical differences among treatments only for body weight at six weeks of age, which increased significantly ($P < 0.05$) by elevating vitamin E levels, while body weight gain, feed consumption and feed conversion were not influenced by dietary treatments.

The utilization of caprylic acid as antibiotics replacement supposes to gather enough data on the influence of CA addition on poultry performance. These data for poultry have been missing until now. The aim of the present study was to determine CA dosage in feed mixture for broiler chickens and the effect of the combination of CA and vitamin E on performance, quality and oxidative stability of meat.

MATERIAL AND METHODS

Four experiments (EXP) were performed with Ross 308 cockerels. Experiment 1: nine hundred chickens (0 day old) were randomly assigned to 9 pens containing 100 chicks each. Each pen was equipped with nipple drinkers and pan feeders. Three replications of 3 dietary treatments were used (Table 1). The basal diet supplemented with 30 mg of α -tocopherol acetate per kg of feed was a control (total analyzed vitamin E 52.7 mg/kg). Chicks in dietary treatments 2 and 3 were fed a basal diet supplemented with caprylic acid (Sigma) 5 g/kg until the 21st day or during the whole EXP. The chicks were kept under a 24-h constant lighting schedule and allowed *ad libitum* access to feed and water. Broiler chickens were slaughtered at 42 days

of age. Compared to EXP 1, in EXP 2 50 cockerels were used in each treatment. Three treatments were used: basal diet, basal diet supplemented with 0.5% of caprylic acid or 130 mg of vitamin E (the total concentration of vitamin E was 150 mg/kg) and 0.5% of caprylic acid. The experiment was terminated at 38 days of age. EXP 3 was realized similarly like EXP 2 but with lower caprylic acid supplement (0.25%) and EXP 3 lasted for 42 days. In EXP 4, one hundred male broiler chicks were in each treatment. The basal diet with 50 mg of vitamin E without supplementation of caprylic acid was

Table 1. Composition of the basal diet

Ingredients	(g/kg)
Wheat	299.2
Maize	300
Soybean meal	320
Fish meal	10
Rapeseed oil ^a	40
Limestone	12
Dicalcium phosphate	10
Sodium chloride	2
Vitamin-mineral premix ^b	5
DL-methionine	1.8
Calculated nutrient composition ^c	
Dry matter	883
Crude protein	218
Crude fat	61
Calcium	8.9
Phosphorus	6.2
Caprylic acid (mg/kg) (analyzed)	0.009
Vitamin E (mg/kg) (analyzed)	52.7
AME _N (MJ/kg)	12.59

^acaprylic acid was mixed with rapeseed oil

^bpremix provided per kg of diet: retinyl acetate 3.6 mg; cholecalciferol 13 μ g; α -tocopherol acetate 30 mg; menadione 3 mg; thiamine 3 mg; riboflavin 5 mg; pyridoxine 4 mg; cyanocobalamin 40 μ g; calcium pantothenate 12 mg; biotin 0.15 mg; folic acid 1.5 mg; choline chloride 250 mg; ethoxyquin 100 g; copper 12 mg; iron 50 mg; iodine 1 mg; manganese 80 mg; zinc 60 mg; selenium 0.3 mg

^cZelenka et al. (2007)

a control treatment. Other treatments had 0.25% of caprylic acid and various total vitamin E contents (50, 100, 200 and 300 mg/kg) in diet. The experiment was terminated at 38 days of age and carcass analysis followed.

Analyses

The breast meat (EXP 4) for dry matter, fat and crude protein determination ($n = 10$ per treatment) was stored in plastic bags at -20°C and meat for MDA and vitamin determination was stored at -70°C . Fat content in basal diet and breast meat was determined by extraction with petroleum ether in a Soxtec 1045 apparatus (Tecator Comp., Sweden) and crude protein was analyzed using a Kjeltex Auto 1030 (Tecator Comp., Sweden). Lipid oxidation in breast meat samples ($n = 8$ per treatment) was measured by the method of Piette and

Raymond (1999), and results were expressed as thiobarbituric acid-reactive substances (TBARS) in mg of malondialdehyde (MDA) per kg of muscle. Before analysis, the breast meat was thawed and stored in a refrigerator at a temperature range from 2.5 to 4°C for 0, 3 or 5 days. The α -tocopherol and retinol contents in breast meat ($n = 10$ per treatment) were determined in accordance with the EN 12822 European standard (2000), by HPLC (Shimadzu, VP series) equipped with a diode-array detector. Each method was used for determination of vitamin E in diet. The concentration of caprylic acid in basal diet was determined by gas chromatographic analysis using a Hewlett-Packard 5890 gas chromatograph equipped with a programmed HP-Innova capillary column (180°C to 240°C) and FI detector.

Resultant values were statistically analyzed by analysis of variance (ANOVA) using the GLM procedure of SAS (2003).

Table 2. Growth traits (EXP 1)

Caprylic acid (%)	BW (g)	F:G	Mortality (%)
0	2 117.3 \pm 399.12 ^a	1.74	4
0.5 until 21 st day	1 899.9 \pm 408.95 ^b	1.76	0
0.5 during whole fattening	1 871.5 \pm 484.29 ^b	1.57	1

^{a,b}treatment means (\pm SE) with different superscripts differ ($P < 0.05$)

Table 3. Growth traits (EXP 2)

Caprylic acid (%)	Vitamin E (mg/kg)	BW (g)	F:G	Mortality (%)
0	50	2 197.6 \pm 316.28 ^a	1.66	2
0.5	50	2 134.1 \pm 254.61 ^{ab}	1.71	2
0.5	150	2 028.4 \pm 260.59 ^b	1.84	0

^{a,b}treatment means with different superscripts differ ($P < 0.05$)

Table 4. Growth traits (EXP 3)

Caprylic acid (%)	Vitamin E (mg/kg)	BW (g)	F:G	Mortality (%)
0	50	2 614.4 \pm 247.96 ^a	1.61	0
0.25	50	2 622.8 \pm 261.77 ^a	1.63	0
0.25	150	2 429.2 \pm 323.18 ^b	1.84	0

^{a,b}treatment means with different superscripts differ ($P < 0.05$)

Table 5. Growth traits (EXP 4)

Caprylic acid (%)	Vitamin E (mg/kg)	BW (g)			F:G	Mortality (%)
		1 st day	21 st day	38 th day		
0	50	47	845.8 ± 149.71 ^a	2 608.3 ± 2 74.49 ^{ab}	1.61	1
0.25	50	47	807.2 ± 134.59 ^b	2 555.9 ± 234.50 ^{bc}	1.62	2
0.25	100	47	764.6 ± 117.84 ^c	2 339.0 ± 211.95 ^d	1.52	4
0.25	200	46	830.7 ± 145.55 ^{ab}	2 633.2 ± 253.86 ^a	1.51	7
0.25	300	46	759.9 ± 141.97 ^c	2 538.1 ± 263.16 ^c	1.67	2

^{a,b,c,d}treatment means with different superscripts differ ($P < 0.001$)

RESULTS

As shown in Tables 2 and 3, the supplementation of higher doses of CA (0.5%) to the diet decreased ($P < 0.05$) body weight. The results of EXP 3 and 4 (Table 4 and 5) showed that the diet supplemented with 0.25% CA and 30 mg vitamin E (total vitamin E concentration 50 mg) resulted in similar BW like the control diet. At the higher dietary vitamin E concentration of 150 mg (EXP 3, Table 4) and 100 mg

(EXP 4, Table 5), the BW significantly ($P < 0.05$ and $P < 0.001$) decreased. Raising the vitamin E concentration in diet caused an increase in mortality with the exception of 300 mg. The highest supplement of vitamin E (total dietary concentration 300 mg) increased ($P < 0.02$) the breast percentage share in carcass weight, but no effect of vitamin E on thigh was observed (EXP 4, Table 6). No effect of treatments on fat and protein contents was recorded. The increasing vitamin E level caused a significant

Table 6. Carcass analysis – percentage share in carcass weight (EXP 4)

Characteristics	Basal diet and supplement					Significance
	basal	50 mg/kg	100 mg/kg	200 mg/kg	300 mg/kg	
BW (g)	2 704.0 ^a	2 542.3 ^b	2 338.1 ^c	2 651.2 ^a	2 561.5 ^b	0.001
Carcass weight (g)	2 001.6 ^a	1 831.4 ^b	1 647.6 ^c	1 831.1 ^b	1 892.1 ^b	0.001
Breast (%)	21.0 ^b	22.0 ^{ab}	21.1 ^b	21.1 ^b	23.3 ^a	0.017
Skin from breast (%)	1.9	2.0	1.9	1.9	2.0	NS
Thigh (%)	27.5	26.6	26.8	27.2	26.6	NS
Thigh without bone (%)	20.9 ^a	19.0 ^c	19.6 ^{bc}	20.5 ^{ab}	20.2 ^{ab}	0.016
Skin from thigh (%)	5.6	5.6	5.5	5.3	5.2	NS
Liver (%)	2.5	2.5	2.7	2.6	2.5	NS
Heart (%)	0.7	0.7	0.7	0.7	0.7	NS
Gizzard (%)	1.3 ^b	1.3 ^b	1.6 ^a	1.6 ^a	1.5 ^{ab}	0.028
Abdominal fat (%)	1.0	1.1	1.0	0.9	0.9	NS
Chemical analysis of breast meat (g/kg DM)						
Fat content	32.6	32.2	29.8	30.3	32.1	NS
Crude protein content	888.5	883.8	882.8	884.6	892.7	NS

^{a,b,c}values with different superscripts differ significantly

NS = not significant; $n = 10$ per treatment

Table 7. Vitamin E and A (mg/kg DM) and MDA (mg/kg) in breast meat (EXP 4)

Characteristics	Basal diet and supplement					Significance
	basal	50 mg/kg	100 mg/kg	200 mg/kg	300 mg/kg	
Vitamin E ¹	28.54 ^{cd}	23.54 ^d	35.22 ^c	80.28 ^a	65.87 ^b	0.001
Vitamin A ¹	0.341 ^b	0.337 ^b	0.333 ^b	0.440 ^a	0.433 ^a	0.001
MDA 0 day ²	0.593	0.490	0.523	0.480	0.490	NS
MDA after 3 days ²	0.678 ^a	0.563 ^c	0.613 ^{abc}	0.583 ^{bc}	0.643 ^{ab}	0.019
MDA after 5 days ²	0.713 ^a	0.593 ^b	0.698 ^a	0.643 ^{ab}	0.708 ^a	0.05

^{a,b,c,d} values with different superscripts differ significantly

NS = not significant; ¹*n* = 10 per treatment; ²*n* = 8 per treatment

($P < 0.001$) increase in the vitamin content in breast meat (Table 7). The highest values of both vitamins were recorded in the treatment with 0.25% of CA and 200 mg of vitamin E. The effect of CA and vitamin E supplementation on lipid oxidation in breast meat during storage is also shown in Table 7. The addition of CA significantly decreased the speed of lipid oxidation measured by MDA formation after 3 ($P < 0.02$) and 5 ($P < 0.05$) days of storage. The lower values were determined in chickens receiving 0.25% of CA and 50 mg of vitamin E in diet. However, the highest vitamin E concentration was found to be related to an increase in lipid oxidation.

DISCUSSION

Many *in vitro* experiments brought unambiguously effective results of CA application (Nair et al., 2005; Skřivanová et al., 2005, 2006; Vasudevan et al., 2005). Successful suppression of experimental infection of rabbits followed (Skřivanová et al., 2008). With the exception of experiments on pigs (Dierick et al., 2002; Marounek et al., 2004) and rabbits (Skřivanová et al., 2008), no data on performance after CA addition to diet have been published yet. An additional test with 10 chickens/treatment until the age of 10 days was published only (Solis de los Santos, 2008). The influence of polyunsaturated fatty acid (PUFA) on the production capability of chickens was studied (Zelenka et al., 2006, 2008 and others) as well as the effect of unsaturated FA and long-chain saturated FA (Skřivan et al., 2000). On the other hand, MCFA stays off in this course has not been mentioned up to the present.

The addition of 0.5% CA was found successful in pigs and rabbits. In chickens, it was necessary to decrease the supplementation from original 0.5% in EXP 1 and 2 to 0.25%. A higher CA dose strongly reduced the average live weight of chickens at the end of fattening. It was also observed when the diet with CA was applied only in the first 3 weeks of experiment. On the contrary, mortality of chickens significantly declined from 4% to 0% or 1%. The negative effect of higher CA supplementation was found out also in the following experiment. Chickens had insignificantly lower live weight and worse feed conversion. The worse results in consequence of a higher dose of CA cannot be explained by CA toxicity according to literature. MCFA have a high degree of safety documented by oral, parenteral and dermal tests (Traul et al., 2000). Furthermore, CA is a food-grade chemical approved by the FDA as GRAS (FDA, 1981). A further important decrease in chicken live weights was recorded at the higher supplementation of vitamin E. Vitamin E concentration in the basal diet was at the level 50 mg/kg. Vitamin-mineral premix added 30 mg of vitamin E/kg and components another 23 mg. The analytically determined vitamin E content was 52.7 mg/kg. Increased supplementation of vitamin E together with CA was found to be unsuitable because it either reduced the live weight of chickens or increased mortality. Similar findings were ascertained also with 0.25% CA addition. The cause is not known. CA is a saturated FA. The antioxidant ethoxyquin was added into the feed mixture by vitamin-mineral premix. The nutritious requirement of vitamin E was met by its content 53 mg. The effect of CA was also positive according to MDA production. Consequently, higher doses of vitamin E

were not only useless but also harmful in combination with CA. In experiment 4, the depression of chicken growth was eliminated but mortality was the highest when 200 mg/kg of vitamin E were supplemented in total. Chicken mortality decreased at the level of treatment with 50 mg of vitamin E due to a further increase in vitamin E to the total content of 300 mg/kg feed mixture. However, the MDA concentration remained high. Caprylic acid can be fully safe but data on interactions with other substances are missing.

Solis de los Santos et al. (2008) evaluated the effect of CA (from 0.35% to 1.4%) on the caecal *Campylobacter jejuni* colonization at experimental infection in 4 trials with 10 days old chicks. The dose of 0.35% CA, which is closer to our dose of 0.25% than to 0.5% CA, was more efficient compared with the spectrum of other five higher additions up to 1.4%. The authors simultaneously evaluated the live weight of chickens at 10 days, feed consumption and feed conversion which is less than orientation with 10 chicks/treatment. The lowest supplement (0.35%) did not decrease any of the three above-mentioned performance characteristics significantly in comparison with the basal diet. On the contrary, the highest doses were markedly worse, which is in accordance with the results presented by us.

Neither carcass analysis nor fat and crude protein content in meat brought any crucial knowledge in relation to CA. The negative effect of vitamin E up to 200 mg/kg on the weight of chickens did not prevent its higher deposition in breast meat. Moreover, the 2 highest additions of vitamin E significantly ($P < 0.001$) increased also vitamin A in meat. A comparison of the values of vitamin E and A content in meat with MDA shows that the worst results were ascertained in the combination of 0.25% of CA and 300 mg of vitamin E in diet. The concentration of MDA is in principle consistent with the control treatment. The relatively high addition of vitamin E did not increase the oxidation stability of meat stored for 3 or 5 days. Very low TBARS were found out at 0.25% of CA and total content 53 mg of vitamin E. Because the same content of vitamin E was also used in the control, it is possible that the enhancement of breast meat oxidation stability is due to CA supplementation.

On the basis of the results from presented experiments with chickens CA dosage in the amount 0.25% of diet can be recommended. Performance will not decrease, mortality can be reduced and

antibacterial activity will stay sufficiently effective according to the last reports. The oxidation stability of breast meat stored for 3–5 days at a temperature from 2.5 to 4°C will increase at dietary content of 0.25% CA and 50 mg/kg of vitamin E. It is not recommended to combine CA with a higher dose of vitamin E than 50 mg/kg.

REFERENCES

- Attia A.A., Abou-Zeid A.E., Mohamed F.F., Yakout H.M. (2001): Enhancement of broiler performance and immune response by α -tocopherol supplemented in diets. *Pakistan Journal of Biological Sciences*, 4, 1029–1035.
- Bergsson G., Arnfinnsson J., Karlsson S.M., Steingrímsson Ó., Thormar H. (1998): *In vitro* Inactivation of *Chlamydia trachomatis* by fatty acids and monoglycerides. *Antimicrobial Agents and Chemotherapy*, 42, 2290–2294.
- Dierick N., Decuypere J.A., Molly K., van Beek E., Vanderbeke E. (2002): Medium chain fatty acids as a nutritional alternative for in-feed antibiotics in piglet nutrition. *Reproduction, Nutrition, Development*, 42, 18 pp.
- EN 12822 (2000): Foodstuffs – Determination of vitamin E by high performance liquid chromatography – measurement of α -, β -, γ - and δ -tocopherols. European Standard. European Committee for Standardization, Brussels, Belgium.
- FDA (Food and Drug Administration) (1981): Code of Federal Regulations 21 CFR 184.1025. The Office of the Federal Register National Archives and Records Administration. Washington, USA.
- Guo Y., Tang Q., Yuan J., Jiang Z. (2001): Effects of supplementation with vitamin E on the performance and the tissue peroxidation of broiler chicks and the stability of thigh meat against oxidative deterioration. *Animal Feed Science and Technology*, 89, 165–173.
- Hassinen J.B., Durbin G.T., Bernhart F.W. (1951): The bacteriostatic effect of saturated fatty acids. *Archives of Biochemistry and Biophysics*, 31, 183–189.
- Jensen R.G. (2002): The composition of bovine milk lipids: January 1995 to December 2000. *Journal of Dairy Science*, 85, 295–350.
- Jensen R.G., Ferris A.M., Lammi-Keefe C.J., Henderson R.A. (1990): Lipids of bovine and human milks: A comparison. *Journal of Dairy Science*, 73, 223–340.
- Kabara J.J., Swieczkowski D.M., Conley A.J., Truant J.P. (1972): Fatty acids and derivatives as antimicrobial agents. *Antimicrobial Agents and Chemotherapy*, 2, 23–28.

- Marounek M., Skřivanová E., Skřivanová V. (2004): A note on the effect of caprylic acid and triacylglycerols of caprylic and capric acid on growth rate and shedding of coccidia oocysts in weaned piglets. *Journal of Animal and Feed Sciences*, 13, 269–275.
- Nair M.K., Joy J., Vasudevan P., Hinckley L., Hoagland T.A., Venkitanarayanan K.S. (2005): Antibacterial effect of caprylic acid and monocaprylin on major bacterial mastitis pathogens. *Journal of Dairy Science*, 88, 3488–3495.
- Nieman C. (1954): Influence of trace amounts of fatty acids on the growth of microorganisms. *Bacteriological Reviews*, 18, 147–163.
- Piette G., Raymond Y. (1999): Vergleichende bewertung verschiedener methoden. Bestimmung der Ranzigkeit in Fleischprodukten. *Fleischwirtschaft*, 7, 69–73.
- SAS (2003): The SAS System for Windows. Release 9.1.3. SAS Institute Inc., USA.
- Skřivan M., Skřivanová V., Marounek M., Tůmová E., Wolf J. (2000): Influence of dietary fat source and copper supplementation on broiler performance, fatty acid profile of meat and depot fat, and on cholesterol content in meat. *British Poultry Science*, 41, 608–614.
- Skřivanová E., Marounek M., Dlouhá G., Kaňka J. (2005): Susceptibility of *Clostridium perfringens* to C₂–C₁₈ fatty acids. *Letters in Applied Microbiology*, 41, 77–81.
- Skřivanová E., Marounek M., Benda V., Březina P. (2006): Susceptibility of *Escherichia coli*, *Salmonella* sp. and *Clostridium perfringens* to organic acids. *Veterinární medicína*, 51, 81–88.
- Skřivanová E., Molatová Z., Marounek M. (2008): Effects of caprylic acid experimentally infected with enteropathogenic *Escherichia coli* O103. *Veterinary Microbiology*, 126, 372–376.
- Solis de los Santos F., Donoghue A.M., Venkitanarayanan K., Dirain M.L., Reyes-Herrera I., Blore P.J., Donoghue D.J. (2008): Caprylic acid supplemented in feed reduces enteric *Campylobacter jejuni* colonization in ten-day-old broiler chickens. *Poultry Science*, 87, 800–804.
- Sprong R.C., Hulstein M.F.E., van der Meer R. (2001): Bactericidal activities of milk lipids. *Antimicrobial Agents and Chemotherapy*, 45, 1298–1301.
- Traul K.A., Driedger A., Ingle D.L., Nakhasi D. (2000): Review of the toxicologic properties of medium chain triglycerides. *Food and Chemical Toxicology*, 38, 79–98.
- Van Immerseel F., De Buck J., Boyen F., Bohez L., Pasmans F., Volf J., Sevcik M., Rychlik I., Haesebrouck F., Ducloux R. (2004): Medium-chain fatty acids decrease colonization and invasion through *hilA* suppression shortly after infection of chickens with *Salmonella enterica* serovar Enteritidis. *Applied and Environmental Microbiology*, 70, 3582–3587.
- Vasudevan P., Patrick M., Nair M., Annamalai T., Darre M., Khan M., Venkitanarayanan K. (2005): *In vitro* inactivation of *Salmonella* Enteritidis in chicken cecal contents by caprylic acid. *Journal of Applied Poultry Research*, 14, 122–125.
- Zelenka J., Schneiderová D., Mrkvicová E. (2006): Linseed oils with different fatty acid patterns in the diet of broiler chickens. *Czech Journal of Animal Science*, 51, 117–121.
- Zelenka J., Heger J., Zeman L. (2007): Nutrient requirements and nutritive values of feeds for poultry. ČAZV, 78 pp. (in Czech)
- Zelenka J., Jarošová A., Schneiderová D. (2008): Influence of n-3 and n-6 polyunsaturated fatty acids on sensory characteristics of chicken meat. *Czech Journal of Animal Science*, 53, 299–305.

Received: 2009–08–03

Accepted after corrections: 2009–10–25

Corresponding Author

Prof. Miloš Skřivan, Institute of Animal Science, Přátelství 815, 104 01 Prague 10-Uhřetěves, Czech Republic
Tel. +420 267 009 720, fax +420 267 711 448, e-mail: skřivan.milos@vuzv.cz
