



Milk Conjugated Linoleic Acid Response to Fish Oil and Linseed Oil Supplementation of Grazing Dairy Cows

Brown, W., A. A. AbuGhazaleh* and S. A. Ibrahim¹

Department of Animal Science, Food and Nutrition. Southern Illinois University Carbondale, IL 62901, USA

ABSTRACT : The effect of supplementing the diet of grazing dairy cows with fish oil (FO) and linseed oil (LSO) on milk conjugated linoleic acid (CLA) was investigated. Sixteen Holstein cows (170±19 DIM) were assigned into two groups and fed a grain supplement (8.0 kg/d; DM basis) containing 800 g of saturated animal fat (CONT) or 200 g FO and 600 g LSO (FOLSO). All cows grazed together on Sudan grass pasture *ad libitum* and were fed the treatment diets for 3 wks. Cows were milked twice a day and milk samples were collected during the last three days of the trial. Milk production (24.89 and 22.45 kg/d), milk protein percentage (2.76 and 2.82) and milk protein yield (0.68 and 0.64 kg/d) for the CONT and FOLSO diets, respectively, were not affected ($p>0.05$) by treatment diets. Milk fat percentage (3.90 and 2.86) and milk fat yield (0.97 and 0.64 kg/d) were lower ($p<0.05$) with the FOLSO diet compared with the CONT diet. The concentration and yield of milk *cis-9 trans-11* CLA were higher ($p<0.05$) with the FOLSO diet (2.56% of total FA and 16.44 g/d, respectively) than the CONT diet (0.66% of total FA and 6.44 g/d, respectively). The concentrations of milk *trans* C18:1 and vaccenic acid (VA) were higher ($p<0.05$) with the FOLSO diet (13.53 and 7.48% of total FA, respectively) than the CONT diet (3.69 and 2.27% of total FA, respectively). In conclusion, supplementing the diet of grazing cows with FO and LSO increased milk *cis-9 trans-11* CLA content but reduced milk fat content and yield. (**Key Words :** Dairy, Pasture, Milk, CLA, Fish Oil, Linseed Oil)

INTRODUCTION

Conjugated linoleic acid (CLA) refers to a mixture of positional and geometric isomers of octadecadienoic acid with conjugated double bonds. CLA are natural occurring fatty acids (FA) in foods derived from ruminants. In milk fat, the *cis-9 trans-11* CLA account for at least 82% of total CLA isomers (Sehat et al., 1998). The *cis-9 trans-11* CLA is synthesized either in the rumen as an intermediate during the biohydrogenation of linoleic acid or in tissues by Δ -9 desaturase from vaccenic acid (VA), another important intermediate in ruminal biohydrogenation of C18 unsaturated FA (Grinari and Bauman, 1999; AbuGhazaleh et al., 2005). Dietary *cis-9 trans-11* CLA is now receiving great attention because of its recognized health benefits as a cancer chemopreventive and antitherogetic (Berlury et al., 2002).

Concentration of CLA is higher in milk fat from cows offered fresh compared with conserved forages (Kelly et al.,

1998; Dhiman et al., 1999; Couvreur et al., 2006). Dhiman et al. (1999) conducted a study in which cows consumed either 1/3, 2/3, or all of their daily feed from pasture. They showed that CLA concentration in the milk fat was highest for cows on 100% pasture (22.1 mg/g of FA), intermediate for cows on 2/3 pasture (14.3 mg/g of FA), and least for cows receiving 1/3 pasture (8.9 mg/g of FA). Recently, Couvreur et al. (2006) reported a linear relationship between the proportion of fresh grass in dairy cows' diets and milk *cis-9 trans-11* CLA content. Milk *cis-9 trans-11* CLA content can be also enhanced by supplementing dairy cows' diets with lipid sources (Schroeder et al., 2003; Boken et al., 2005; Flowers et al., 2007). Schroeder et al. (2003) reported 173% increase in milk CLA when corn grain in grazing dairy cows' diet was partially replaced with Ca-salts of unsaturated FA. When cows on annual ryegrass pasture were supplemented with a grain mix containing 0 or 500 g/d of soybean oil, milk CLA content increased by 46% with the soybean oil supplementation (Boken et al., 2005). Looor et al. (2005a) reported a significant increase in milk *cis-9 trans-11* CLA content when dairy cows' diet was supplemented with linseed oil (LSO) at 3% of diet DM. Recently, Flowers et al. (2007) reported a linear increase in milk *cis-9 trans-11* CLA

* Corresponding Author: Amer A. AbuGhazaleh. Tel: +1-618-453-1767, Fax: +1-618-453-5231, E-mail: aabugha@siu.edu

¹ Department of Family and Consumer Sciences. North Carolina A&T State University. Greensboro, NC 27411, USA.

Received May 22, 2007; Accepted August 27, 2007

Table 1. Ingredient and nutrient content of grain supplement and pasture

Ingredient (% of DM)	Treatment diets ¹		Pasture
	CONT	FOLSO	
Corn ground cracked	42.20	42.20	
Linseed oil	0.00	7.40	
Fish oil	0.00	2.40	
Animal fat (Rumo-fat)	9.80	0.00	
Molasses dried	8.30	8.30	
Soybean meal (48% CP)	34.20	34.20	
Limestone	2.80	2.80	
Dicalcium phosphate	1.40	1.40	
Magnesium oxide	0.30	0.30	
Zn trace mineralized salt	0.50	0.50	
Vit ADE premix ²	0.30	0.30	
Vit E premix ³	0.30	0.30	
Chemical composition (% of DM)			
NDF	22.67±2.03	21.28±2.55	63.92±5.60
ADF	18.98±2.02	13.95±1.67	34.62±4.44
EE	10.91±0.58	10.07±1.47	0.74±0.17
CP	19.82±0.97	20.51±1.45	13.28±5.50

¹ CONT = Grain supplement with 800 g of Rumo fat; FOLSO = Grain supplement with 200 g FO and 600 g LSO.

² Contained 4,400,000 IU of vitamin A, 880,000 IU of vitamin D, and 400 IU of vitamin E/kg.

³ Contained 44,000 IU of vitamin E/kg.

content with increasing the level of LSO in grazing dairy cows' diet.

Inclusion of fish oil (FO) in dairy cows' diets has been also shown to increase milk *cis-9 trans-11* CLA content (Donovan et al., 2000; Whitlock et al., 2002). Milk fat *cis-9 trans-11* CLA response to FO is further enhanced when given in combination with linoleic or linolenic acid-rich oil sources (AbuGhazaleh et al., 2002, 2003). Despite the volume of research on the impact of dietary FO and plants oils on milk *cis-9 trans-11* CLA content under confinement feeding system, there is surprisingly little data in the literature evaluating the effect that dietary FO and plants oils may have under grazing feeding system on milk *cis-9 trans-11* CLA content. Therefore, the objective of this trial was to evaluate milk *cis-9 trans-11* CLA response to FO and LSO supplementation when cows fed a pasture-based diet.

MATERIALS AND METHODS

All procedures for this study involving the use of animals were conducted at The Southern Illinois University Carbondale Dairy Center and were approved by the Southern Illinois University Intuitional Animal Care and Use Committee. Sixteen multiparous Holstein cows of mean live weight 610±23.3 kg were divided into two groups (8 cows per group) based on milk yield (27±3.2 kg/d) and DIM (170±19), placed on Sudan-grass pasture, and offered 8 kg of grain supplement containing either 800 g saturated animal fat (rumo-fat; CONT) or 800 g of FO-LSO (1:3 w/w; FOLSO) for a total of 21 days. The first 18 days were

used for adjustment to diets, and the last 3 days for milk samples collection. Grain supplements were formulated to be isocaloric and isonitrogenous (Table 1). Menhaden FO (Omega Protein Inc., Hammond, LA) and Rumo-fat (Robt Morgan Inc. Paris, IL), a source of saturated animal fat, were used in this study. Linseed oil was purchased from a local store. Cows were offered the grain supplement into two equal meals at 06:30 and 16:30 using Calan Broadbent feeder doors (American Calan, Inc., Northwood, NH) to monitor individual intakes. The Sudan-grass pasture area was a twelve ha area divided into fifteen 0.8 ha paddocks for rotational grazing where cows moved to a new paddock every day. Cows were on the experimental pasture for 21 h a day and had continuous access to water. Cows were milked twice daily at 06:00 and 16:00.

Milk yields were recorded daily, but measurements collected during the last 5 days were used for statistical analysis. Milk samples were collected from each cow during the last 3 days (d 19, 20, and 21) from the morning and afternoon milkings, made into daily composites based on milk yield at each milking, and then split into two portions for analyses. One aliquot was stored at 4°C and sent to Prairie Farms (Carlinville, IL) for analyses of fat, protein, total solids, and lactose (AOAC, 1997) by mid-infrared spectrophotometry (Multispec; Foss Food Technology Corp., Eden Prairie, MN) and somatic cell counts (SCC; AOAC, 1997) using a Fossomatic 90 (Foss Food Technology Corp.). The second aliquot was stored at -20°C for FA analysis as described by AbuGhazaleh and Holmes (2007).

Grain supplements offered were recorded daily, but only

Table 2. Fatty acid composition for grain supplements, pasture, rumo fat, linseed oil, and fish oil

Fatty acid	Treatment diets ¹		Pasture ²	Lipid Supplements		
	CONT	FOLSO		Rumo-fat	LSO ³	FO ⁴
	----- g/100 g FA -----					
C14:0	1.41	1.34	0.33	2.35	0.05	8.25
C16:0	23.68	9.30	13.46	20.63	4.58	20.27
C18:0	46.38	3.49	3.31	52.05	2.85	4.14
C18:1 <i>cis</i> -9	6.88	16.09	2.28	0.28	17.31	11.22
C18:2 <i>cis</i> -9, <i>cis</i> -12	13.70	17.18	13.95	0.04	13.63	1.37
C18:3n3	0.93	34.38	37.66	nd ⁵	52.11	1.81
C20:5n3, EPA	0.01	1.42	0.02	nd	nd	13.03
C22:6n3, DHA	0.02	1.25	0.07	nd	nd	15.99
Unknown	6.85	15.42	28.86	24.49	9.41	23.82

¹ CONT = Grain supplement with 800 g of Rumo fat; FOLSO = Grain supplement with 200 g FO and 600 g LSO.

² Pasture = Sudan grass. ³ LSO = Linseed oil. ⁴ FO = Fish oil. ⁵ nd = Not detected or detected at <0.01.

Table 3. Effect of treatment diets on milk yield, composition and grain supplement intake.

Item	Treatment diets ¹		SEM ²	p-value
	CONT	FOLSO		
Milk (kg/d)	24.89	22.45	0.88	0.07
Fat				
%	3.90	2.86	0.13	0.01
kg/d	0.97	0.64	0.04	0.01
Protein				
%	2.76	2.82	0.11	0.74
kg/d	0.68	0.64	0.03	0.30
Lactose				
%	4.60	4.63	0.07	0.77
kg/d	1.14	1.04	0.05	0.20
Total solids				
%	12.16	11.46	0.14	0.01
kg/d	3.02	2.57	0.10	0.01
SCC×10 ³ /ml	133.93	938.76	290.80	0.07
Grain supplement intake (kg/d)	7.58	7.83	0.21	0.42
Body weight (kg)	585	577	15.50	0.43
BCS ³	3.10	2.90	0.25	0.33

¹ CONT = Grain supplement with 800 g of Rumo fat; FOLSO = Grain supplement with 200 g FO and 600 g LSO.

² SEM = Standard error mean. ³ Scored on a five-point scale where 1 = emaciated to 5 = overly fat.

measurements collected during the last 5 days were used for statistical analysis. Orts were removed and weighed after the morning feeding. Representative samples of Sudan grass (collected from 4 different locations per paddock then composited into one) and grain supplements were collected at d 7, 14 and 21 and stored at -20°C until analysis for chemical and FA composition. Grain supplements and pasture samples were freeze dried (Labconco Freeze Dry System, Labconco, Kansas City, MO) and then ground through a standard model No. 3 Wiley mill (Arthur H. Thomas Co., Philadelphia, PA) with a 2-mm screen. Samples were analyzed for CP and EE according to AOAC methods (1997). A portion of each sample was reground through an ultracentrifuge mill (Brinkman Instruments Co., Westbury, NY) with a 1-mm screen and analyzed for NDF (Procedure B, Van Soest et al., 1991), ADF (Robertson and Van Soest, 1981) using ANKOM fiber analyzer using the

filter bag technique (ANKOM Technology Corp., Fairport, NY) and FA as described for milk. The C17:0, dissolved in benzene, was used as an internal standard. Cows body weights and body condition score (BCS; Wildman et al., 1982) were recorded at the beginning and end of the trial.

Experimental data was subjected to the PROC MIXED procedure of SAS (SAS, INC. Cary, NC) for a completely randomized design using following statistical model:

$$Y_{ijk} = \mu + D_i + C_j + S_k + e_{ijk}$$

where Y_{ijk} = the dependent variable; μ = overall mean; D_i = diet effect; C_j = cow effect; S_k = sample effect and e_{ijk} = residual error associated with Y_{ijk} . The values for least squares means are reported followed by standard error of mean (SEM). Treatment effects were declared significance at $p < 0.05$.

RESULTS AND DISCUSSION

Linolenic acid was the main FA in pasture, it accounted for 37.66% of total FA. Both palmitic and linoleic acids were the second main FA in pasture; they accounted for approximately 13.46 and 13.95% of total FA, respectively (Table 2). As expected, the FA profile in the CONT diet was characterized by its high stearic acid content. In the FOLSO diet, linolenic acid was the main FA followed both linoleic and oleic acids. Stearic and Palmitic acids were the main FA in the rumo-fat. As expected linolenic acid was the main FA in LSO whereas; eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) were the two main omega-3 FA in FO.

Milk yield, milk composition, and grain supplement intake are presented in Table 3. Inclusion of FO and LSO in the diet had no effect ($p>0.05$) on milk yield (24.89 and 22.45 kg/d for the CONT and the FOLSO diets, respectively). Similar results were reported under confinement (Boken et al., 2005; Loor et al., 2005a) and grazing (Rego et al., 2005b; AbuGhazaleh and Holms, 2007; Flowers et al., 2007) feeding conditions when diets were supplemented with oils. Supplementing diets with oils rich in polyunsaturated FA often results in a reduction in feed intake and therefore milk yield (Donovan et al., 2000; Lock and Shingfield, 2004). The equal intake of grain supplements in this experiment may explain the similar milk yield between treatment diets.

Milk fat percentages (3.90 and 2.86) and yields (0.97 and 0.64 kg/d) were higher ($p<0.01$) in cows fed the CONT diet compared with the FOLSO diet (Table 3). Milk fat depression is frequently seen in dairy cows fed oil supplements (AbuGhazaleh et al., 2002; Rego et al., 2005a,b; Shingfield et al., 2006; Holmes and AbuGhazaleh, 2007). After reviewing eighteen experiments involving 25 comparisons to describe the effects of lipid supplementation on milk production and composition with grazing dairy cows, Schroeder et al. (2004) concluded that the decrease in milk fat was a common result when unsaturated lipid supplements were added to grazing dairy cows' diets. Different explanations have been suggested for the milk fat depression associated with lipids feeding. Jenkins (1993) and Schroeder et al. (2004) suggested that the reduction in milk fat content with dietary lipids may results from their negative effects in rumen fiber digestion with a consequent reduction in acetate production, the precursor for short and medium-chain FA synthesis in mammary glands (Harfoot and Hazelwood, 1997). Others (Grummer, 1991; Chillard and Doreau, 1997) suggested that the increase uptake and incorporation of dietary long-chain FA into triglycerides inhibits the *de novo* FA synthesis in the mammary gland and therefore, reduce milk fat content. Recently, Bauman and Griinari (2001) suggested that milk fat depression was

related to the direct action on the mammary gland of specific FA isomers derived from the ruminal metabolism of dietary unsaturated FA. Piperova et al. (2004) and Shingfield et al. (2006) reported a close association between milk fat depression and levels of milk *trans*-10 C18:1. In the current study, the concentration of *trans*-10 C18:1 in milk fat was 808% higher with the FOLSO diet than the CONT diet.

Treatment diets had no effect ($p>0.05$) on milk protein percentages (2.76 and 2.82) and yields (0.68 and 0.64 kg/d). Supplementing grazing cows' diet with 500 g of sunflower oil or soybean oil did not affect milk protein content or yield (Rego et al., 2005b). Similarly, AbuGhazaleh and Holmes (2007) and Flowers et al. (2007) reported no difference in milk protein content in pasture cows fed a diet with or without plant oils. However, others (Dhiman et al., 2000; Shingfield et al., 2006; Whitlock et al., 2006) reported a significant reduction in milk protein content with oil supplementation, changes that have often been attributed to increases in milk yield (dilution effect) rather than decreases in milk protein synthesis (DePeters and Cant, 1992). Similar milk yield in the current study may explain the similar milk protein content between treatment diets. Treatment diets had no effect ($p>0.05$) on grain supplement intake SCC, body weight, and BCS.

The FOLSO diet resulted in marked alterations in milk FA composition and yield relative to the CONT diet (Tables 4 and 5). The concentrations of milk short and medium chain FA (C4:0-C16:0) were significantly reduced ($p<0.05$) with the FOLSO diet compared with the CONT diet. Even short and medium chain FA (C4:0-C14:0) are derived from the *de novo* synthesis by the mammary gland and from the mammary uptake of preformed FA (Jenkins, 1993). In the current study, the lower concentrations of even-number saturated FA with the FOLSO diet compared with the CONT diet suggest less *de novo* FA synthesis occurred with cows fed the FOLSO diet. As stated earlier, feeding oils are typically associated with a decrease in the *de novo* synthesis of short and medium chain FA (Loor et al., 2005a,c; Rego et al., 2005b).

The objective of this study was to evaluate the effects of feeding a CLA-stimulating diet to grazing dairy cows on milk *cis*-9, *trans*-11 CLA content. Milk *cis*-9, *trans*-11 CLA concentration and yield increased by 288 and 155% with the FOLSO diet relative to the CONT diet, respectively (Tables 4 and 5). The concentration of *cis*-9, *trans*-11 CLA averaged 0.66 and 2.56% of total FA with the CONT and the FOLSO diets, respectively. Diets containing FO, plant oils or their combinations consistently increased milk *cis*-9, *trans*-11 CLA content under confinement (Whitlock et al., 2002; Loor et al., 2005a; Bu et al., 2007) and grazing (Boken et al., 2005; Rego et al., 2005b; AbuGhazaleh and Holmes, 2007) feeding conditions. Supplementing grazing

dairy cows with 500 g/d of sunflower oil or soybean oil resulted in 68 and 53% increase in milk CLA compared with control cows (Rego et al., 2005b). Compared with pasture cows on control diet, supplementing grazing dairy cows diet with 160 and 320 g of FO/d resulted in 44 and 62% increase in milk CLA concentration, respectively (Rego et al., 2005a). The higher milk *cis-9, trans-11* CLA concentration seen in this study with the FOLSO diet compared with that reported by AbuGhazaleh et al. (2003; 1.06% of milk total FA), Loor et al. (2005a; 1.34% of milk total FA) and Bu et al. (2007; 1.60% of milk total FA) when dairy cows fed LSO at 524, 588 and 636 g/d, respectively, may have resulted from grazing and/or FO supplementation. Cows on pasture diets have been shown to have higher levels of *cis-9, trans-11* CLA (Boken et al., 2005; Couvreur et al., 2006) in their milk than those on conserved forages. Recently, Couvreur et al. (2006) reported a linear relationship between the proportion of fresh grass in dairy cows' diets and milk *cis-9, trans-11* CLA. AbuGhazaleh

and Jenkins (2004) showed that FO promotes VA accumulations in rumen by inhibiting the reduction of VA to C18:0. As mentioned earlier, *cis-9, trans-11* CLA in milk originates from either the ruminal biohydrogenation of linoleic acid (Harfoot and Hazlewood, 1997) or from the endogenous synthesis in mammary gland from VA (Griinari and Bauman, 1999). The endogenous synthesis of *cis-9, trans-11* CLA from VA has been proposed as the major pathway of *cis-9, trans-11* CLA synthesis in lactating cows, accounting for an estimated 78 to 90% of the *cis-9, trans-11* CLA in milk fat (Corl et al., 2001; Piperova et al., 2002).

Concentrations and yields of milk *trans* C18:1 were also affected by treatment diets (Tables 4 and 5). The concentration of *trans* C18:1 averaged 3.69 and 13.53% of total FA with the CONT and the FOLSO diets, respectively (Table 4). Vaccenic acid was the major *trans* C18:1 isomer in milk accounting for 61.5 and 55.3% of total *trans* C18:1 with the CONT and the FOLSO diets, respectively. Milk VA concentration was higher ($p < 0.05$) with the FOLSO diet

Table 4. Effect of treatment diets on milk fatty acid composition

Fatty acids	Treatment diets ¹		SEM ²	p-value
	CONT	FOLSO		
	----- g/100 g FA -----			
C4:0	1.43	0.91	0.08	0.01
C6:0	1.24	0.68	0.07	0.01
C8:0	0.77	0.38	0.05	0.01
C10:0	1.57	0.76	0.11	0.01
C12:0	1.85	1.08	0.09	0.01
C14:0	7.78	5.76	0.17	0.01
C16:0	27.40	19.85	0.59	0.01
C17:0	0.87	0.48	0.03	0.01
C18:0	15.60	9.45	0.94	0.01
C18:1 <i>trans</i>	3.69	13.53	0.89	0.01
<i>trans-6/8</i>	0.29	0.72	0.05	0.01
<i>trans-9</i>	0.25	0.73	0.06	0.01
<i>trans-10</i>	0.23	2.09	0.47	0.02
<i>trans-11</i> (VA)	2.27	7.48	0.76	0.04
<i>trans-12</i>	0.40	1.46	0.13	0.01
<i>trans-16</i>	0.23	1.03	0.06	0.01
C18:1 <i>cis-9</i>	23.96	22.41	1.19	0.37
C18:2 <i>trans-11, cis-15</i>	0.22	2.09	0.12	0.01
C18:2 <i>cis-9, cis-12</i>	1.99	1.66	0.13	0.09
C18:3n3	0.56	1.26	0.08	0.01
CLA <i>cis-9, trans-11</i>	0.66	2.56	0.28	0.05
CLA <i>trans, trans</i>	0.10	0.38	0.06	0.03
C20:5n3, EPA	0.03	0.07	<0.01	0.01
C22:4n6	0.03	0.04	<0.001	0.04
C22:5n3	0.05	0.07	<0.001	0.01
C22:6n3, DHA	0.01	0.06	<0.001	0.01
SMCFA ³	43.73	30.64	1.04	0.01
SAT ⁴	59.37	40.77	0.99	0.01
UNSAT ⁵	34.75	47.93	0.99	0.01
Unknown	5.73	11.18	1.76	0.01

¹ CONT = Grain supplement with 800 g of Rumo fat; FOLSO = Grain supplement with 200 g FO and 600 g LSO.

² SEM = Standard error mean. ³ SMCFA = Short and medium chain FA (C4:0-C16:0).

⁴ SAT = Sum of saturated FA. ⁵ UNSAT = Sum of unsaturated FA.

Table 5. Effect of treatment diets on milk fatty acid yield

Fatty acids	Treatment diets ¹		SEM ²	p-value
	CONT	FOLSO		
	----- g/d -----			
C4:0	13.60	6.02	0.08	0.01
C6:0	11.90	4.49	0.08	0.01
C8:0	7.35	2.46	0.55	0.01
C10:0	15.06	4.85	1.12	0.01
C12:0	17.88	6.82	1.12	0.01
C14:0	75.40	36.00	3.39	0.01
C16:0	262.80	127.00	10.52	0.01
C17:0	8.37	3.05	0.31	0.01
C18:0	143.50	65.10	7.40	0.01
C18:1 <i>trans</i>	36.61	84.85	6.33	0.01
<i>trans</i> -6/8	2.73	4.48	0.34	0.01
<i>trans</i> -9	2.40	4.60	0.36	0.01
<i>trans</i> -10	2.27	12.39	1.92	0.01
<i>trans</i> -11 (VA)	22.88	47.79	5.53	0.01
<i>trans</i> -12	3.94	9.12	0.99	0.01
<i>trans</i> -16	2.29	6.45	0.28	0.01
C18:1 <i>cis</i> -9	236.17	144.67	14.78	0.01
C18:2 <i>trans</i> -11, <i>cis</i> -15	2.03	13.49	1.02	0.01
C18:2 <i>cis</i> -9, <i>cis</i> -12	19.30	10.82	1.32	0.01
C18:3n3	5.43	8.34	0.78	0.02
CLA <i>cis</i> -9, <i>trans</i> -11	6.44	16.44	2.01	0.01
CLA <i>trans</i> , <i>trans</i>	0.90	2.52	0.04	0.01
C20:5n3, EPA	0.32	0.46	0.06	0.05
C22:4n6	0.28	0.20	0.03	0.12
C22:5n3	0.51	0.48	<0.01	0.54
C22:6n3, DHA	0.05	0.38	0.03	0.01
SMCFA ³	404.84	186.85	15.65	0.01
SAT ⁴	549.47	245.70	16.46	0.01
UNSAT ⁵	317.16	299.96	17.89	0.82

¹ CONT = Grain supplement with 800 g of Rumo fat; FOLSO = Grain supplement with 200 g FO and 600 g LSO.

² SEM = Standard error mean. ³ SMCFA = Short and medium chain FA (C4:0-C16:0).

⁴ SAT = Sum of saturated FA. ⁵ UNSAT = Sum of unsaturated FA.

(7.48 g/100 g FA) than the CONT diet (2.27 g/100 g FA). The increase in concentrations of milk *trans* C18:1 isomers, VA in particular, with the FOLSO diet was expected. Similar increases in milk *trans* C18:1 and VA concentrations were observed under confinement (Loor et al., 2005a,b; Bu et al., 2007) and grazing (Flowers et al., 2007; Holmes and AbuGhazaleh, 2007) feeding conditions when cows were fed FO and/or LSO. AbuGhazaleh and Holmes (2007) observed a significant increase in milk VA content when partially grazed dairy cows were supplemented with a blend of sunflower oil (300 g/d) and FO (100 g/d) when compared to partially grazed dairy cows received control diet (5.11 vs. 2.20 g/100 g of milk total FA). Milk VA accounted for 5.43, 3.23 and 3.04% of milk FA when dairy cows fed LSO at 860, 588 and 636 g/d, respectively (Loor et al., 2005a,c; Bu et al., 2007). The higher milk VA concentration seen in this study (7.48%) may have resulted from grazing, FO or their combinations. Recently, Felton and AbuGhazaleh (2007) and Holmes and AbuGhazaleh (2007) showed that the increase in milk VA

content was higher when oil supplements were fed to dairy cows managed under grazing than confinement feeding systems. Previously, AbuGhazaleh et al. (2002) and Whitlock et al. (2002) showed that combining FO with plant oils results in more increase in milk VA than when fed separately. Human body can convert some of the VA into *cis*-9, *trans*-11 CLA (Salminen et al., 1998); therefore, producing milk with high VA concentration could be also beneficial.

Concentrations of milk *trans*-6/8 C18:1, *trans*-9 C18:1, *trans*-10 C18:1, *trans*-12 C18:1 and *trans*-16 C18:1 were all higher ($p < 0.05$) with the FOLSO diet than the CONT diet. The increase in concentrations of *trans*-6/8, *trans*-9, *trans*-10, *trans*-12 and *trans*-16 C18:1 in milk fat with the FOLSO diet is consistent with that reported by Holmes and AbuGhazaleh (2007), Loor et al. (2005a,c) and Flowers et al. (2007) when FO and/or LSO were added to dairy cows' diets. The presence of these different *trans* C18:1 isomers in milk fat provides comparative evidence for the existence of alternative pathways for the biohydrogenation of C18

polyunsaturated FA other than those established by Harfoot and Hazlewood (1997).

IMPLICATIONS

Supplementing grazing cows' diets with fish oil and linseed oil enhances milk conjugated linoleic acid content and may be used as a practical approach to increase conjugated linoleic acid as functional ingredient in milk.

ACKNOWLEDGMENTS

We thank the employees of Prairie Farms, Carlinville, IL, for milk analysis. We also thank Bruce Jacobson for helping with milk fatty acid analysis. This research was partially supported by a grant from the Illinois Council on Food and Agricultural Research.

REFERENCES

- AbuGhazaleh, A. A., D. J. Schingoethe, A. R. Hippen and K. F. Kalscheur. 2002. Feeding fish meal and extruded soybeans enhances the conjugated linoleic acid content of milk. *J. Dairy Sci.* 85:624-631.
- AbuGhazaleh, A. A., D. J. Schingoethe, A. R. Hippen and K. F. Kalscheur. 2003. Milk conjugated linoleic acid response to fed fish oil supplementation of diets differing in fatty acid profiles. *J. Dairy Sci.* 86:944-953.
- AbuGhazaleh, A. A. and T. C. Jenkins. 2004. Short Communication: Docosahexaenoic acid promotes vaccenic acid accumulation in mixed ruminal cultures when incubated with linoleic acid. *J. Dairy Sci.* 87:1047-1050.
- AbuGhazaleh, A. A., M. B. Riley, E. E. Thies and T. C. Jenkins. 2005. Dilution rate and pH effects on the conversion of oleic acid to trans C18:1 positional isomers in continuous culture. *J. Dairy Sci.* 88:4334-4341.
- AbuGhazaleh, A. A. and L. D. Holmes. 2007. Diet supplementation with fish oil and sunflower oil to increase conjugated linoleic acid levels in milk fat of partial grazing dairy cows. *J. Dairy Sci.* 90:2897-2904.
- Association of Official Analytical Chemists. 1997. Official Methods of Analysis. 16th ed. AOAC Int., Gaithersburg, MD.
- Bauman, D. E. and J. M. Griinari. 2001. Regulation and nutritional manipulation of milk fat: Low-fat milk syndrome. *Livest. Prod. Sci.* 70:15-29.
- Belury, M. A., S. Y. Moya-Camarena, M. Lu, L. Shi, L. M. Leesnitzer and S. G. Blanchard. 2002. Conjugated linoleic acid is an activator and ligand for peroxisome proliferator-activated receptor-gamma (PPAR γ). *Nutr. Res.* 22(7):817-824.
- Boken, S. L., C. R. Staples, L. E. Sollengberger, T. C. Jenkins and W. W. Thatcher. 2005. Effect of grazing and fat supplementation on production and reproduction of Holstein cows. *J. Dairy Sci.* 88:4258-4272.
- Bu, D. P., J. Q. Wang, T. R. Dhiman and S. J. Liu. 2007. Effectiveness of oils rich in linoleic and linolenic acids to enhance conjugated linoleic acid in milk from dairy cows. *J. Dairy Sci.* 90:998-1007.
- Chilliard, Y. and M. Doreau. 1997. Influence of supplementary fish oil and rumen-protected methionine on milk yield and composition in dairy cows. *J. Dairy Res.* 64:173-179.
- Corl, B. A., L. H. Baumgard, D. A. Dwyer, J. M. Griinari, B. S. Phillips and D. E. Bauman. 2001. The role of delta9 desaturase in the production of cis-9, trans-11 CLA. *J. Nutr. Bioch.* 12:622-630.
- Couvreur, S., C. Hurtaud, C. Lopez, L. Delaby and J. L. Peyraud. 2006. The linear relationship between the proportion of fresh grass in the cow diet, milk fatty acid composition, and butter properties. *J. Dairy Sci.* 89:1956-1969.
- DePeters, E. J. and J. P. Cant. 1992. Nutritional factors influencing the nitrogen composition of bovine milk: A review. *J. Dairy Sci.* 75:2043-2070.
- Dhiman, T. R., G. R. Anand, L. D. Satter and M. W. Pariza. 1999. Conjugated linoleic acid content of milk from cows fed different diets. *J. Dairy Sci.* 82:2146-2156.
- Dhiman, T. R., L. D. Satter, M. W. Pariza, M. P. Galli, K. Albright and M. X. Tolosa. 2000. Conjugated linoleic acid content of milk from cows offered diets rich in linoleic and linolenic acid. *J. Dairy Sci.* 83:1016-1027.
- Donovon, D. C., D. J. Schingoethe, R. J. Baer, J. Ryali, A. R. Hippen and S. T. Franklin. 2000. Influence of dietary fish oil on conjugated linoleic acid and other fatty acids in milk from lactating dairy cows. *J. Dairy Sci.* 83:1-9.
- Felton, D. and A. A. AbuGhazaleh. 2007. Milk conjugated linoleic acid response to fish oil and sunflower oil supplementation to dairy cows managed under two feeding systems. *J. Dairy Sci.* (Suppl.1) 90.
- Flowers, G. A. A. AbuGhazaleh and S. Ibrahim. 2007. Optimizing the levels of linseed oil in grazing cows diet to maximize conjugated linoleic acid in milk. *J. Dairy Sci.* (Suppl. 1) 90.
- Griinari, J. M. and D. E. Bauman. 1999. Biosynthesis of conjugated linoleic acid and its incorporation into meat and milk in ruminants (Ed. M. P. Yurawecz, M. M. Mossoba, J. K. G. Kramer, M. W. Pariza and G. J. Nelson). *Advances in Conjugated linoleic acid Research* pp. 180-200. AOCS Press, Champaign.
- Grummer, R. R. 1991. Effect of feed on the composition of milk fat. *J. Dairy Sci.* 74:3244-3257.
- Harfoot, C. G. and G. P. Hazelwood. 1997. Lipid metabolism in the rumen. (Ed. P. N. Hobson and C. S. Stewart) *The Rumen Microbial Ecosystem*. Chapman & Hall, London, UK. 382-426.
- Holmes, L. D. and A. A. AbuGhazaleh. 2007. Milk cis-9, trans-11 CLA and omega-3 fatty acids response to linseed oil and fish oil supplementation when dairy cows managed under confinement or grazing system. *J. Dairy Sci.* (Suppl.1)90.
- Jenkins, T. C. 1993. Lipid metabolism in the rumen. *J. Dairy Sci.* 76:3851-3863.
- Kelly, M. L., E. S. Kolver, D. E. Bauman, M. E. Van Amburgh and L. D. Muller. 1998. Effect of intake of pasture on concentrations of conjugated linoleic acid in milk of lactating cows. *J. Dairy Sci.* 81:1630-1637.
- Lock, A. L. and K. J. Shingfield. 2004. Optimizing milk composition. Nottingham University Press, 107-188.
- Loor, J. J., A. Ferlay, A. Ollier, M. Doreau and Y. Chilliard. 2005a. Relationship among *trans* and conjugated fatty acids and bovine milk fat yield due to dietary concentrate and linseed oil. *J. Dairy Sci.* 88:726-740.

- Loor, J. J., M. Doreau, J. M. Chardigny, A. Ollier, J. L. Sebedio and Y. Chilliard. 2005b. Effects of ruminal or duodenal supply of fish oil on milk fat secretion and profiles of *trans*-fatty acids and conjugated linoleic acid isomers in dairy cows fed maize silage. *Anim. Feed Sci. Tech.* 119:227-246.
- Loor, J. J., A. Ferlay, A. Ollier, M. Doreau and Y. Chilliard. 2005c. High-concentrate diets and polyunsaturated oils alter *trans* and conjugated isomers in bovine rumen, blood, and milk. *J. Dairy Sci.* 88:3986-3999.
- Piperova, L. S., B. B. Teter, K. F. Kalscheur, M. P. Yurawecz, Y. Ku, K. M. Morehouse and R. E. Erdman. 2002. Duodenal and milk *trans* octadecenoic acid and conjugated linoleic acid (CLA) isomers indicate that postabsorptive synthesis is predominant source of *cis*-9 containing CLA in lactating cows. *J. Nutr.* 132:1235-1241.
- Piperova, L. S., U. Moallem, B. B. Teter, J. Sampugna, M. P. Yurawecz, K. M. Morehouse, D. Luchini and R. A. Erdman. 2004. Changes in milk fat in response to dietary supplementation with calcium salts of *trans*-18:1 or conjugated linoleic fatty acids in lactating dairy cows. *J. Dairy Sci.* 87:3836-3844.
- Rego, O. A., H. J. D. Rosa, P. V. Portugal, R. Corderio, A. E. S. Borba, C. M. Vouzela and R. J. B. Bessa. 2005a. Influence of dietary fish on conjugated linoleic acid, omega-3 and other fatty acids in milk fat from grazing dairy cows. *Livest. Prod. Sci.* 95:27-33.
- Rego, O. A., H. J. D. Rosa, P. V. Portugal, T. Franco, C. M. Vouzela, A. E. S. Borba and R. J. B. Bessa. 2005b. The effects of supplementation with sunflower and soybean oils on the fatty acid profile of milk fat from grazing dairy cows. *Anim. Res.* 54:17-24.
- Robertson, J. B. and P. J. Van Soest. 1981. The detergent system of analysis and its application to human foods. In *The Analysis of Dietary Fiber* (Ed. W. P. T. James and O. Theander.). Marcel Dekker, New York, NY. 123-129.
- Salminen, I., M. Mutanen, M. Jauhiainen and A. Aro. 1998. Dietary *trans* fatty acids increase conjugated linoleic acid levels in human serum. *J. Nutr. Biochem.* 9(2):93-98.
- Schroeder, G. F., J. E. Delahoy, I. Vidaurreta, F. Bargo, G. A. Gagliostro and L. D. Muller. 2003. Milk fatty acid composition of cows fed a total mixed ration or pasture plus concentrates replacing corn with fat. *J. Dairy Sci.* 86:3237-3248.
- Schroeder, G. F., G. A. Gagliostro, F. Bargo, J. E. Delahoy and L. D. Muller. 2004. Effects of fat supplementation on milk production and composition by dairy cows on pasture: a review. *Livest. Prod. Sci.* 86:1-18.
- Sehat, N., J. K. G. Kramer, M. M. Mossoba, M. P. Yurawecz, J. A. G. Roach, K. Eulitz, K. M. Morehouse and Y. Ku. 1998. Identification of conjugated linoleic acid isomers in cheese by gas Chromatography, silver ion high performance liquid chromatography and mass spectral reconstructed ion profiles. Comparison of chromatographic elution sequences. *Lipids* 33 (10):963-971.
- Shingfield, K. J., C. K. Reynolds, G. Herva's, J. M. Griinari, A. S. Grandison and D. E. Beever. 2006. Examination of the persistency of milk fatty acid composition responses to fish oil and sunflower oil in the diet of dairy cows. *J. Dairy Sci.* 89:714-732.
- Van Soest, P. J., J. B. Robertson and B. A. Lewis. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysacchrides in relation to animal nutrition. *J. Dairy Sci.* 74:3583-3597.
- Whitlock, L. A., D. J. Schingoethe, A. R. Hippen, K. F. Kalscheur, R. J. Baer, N. Ramaswamy and K. M. Kasperson. 2002. Fish oil and extruded soybeans fed in combination increase conjugated linoleic acids in milk of dairy cows more than when fed separately. *J. Dairy Sci.* 85:234-243.
- Whitlock, L. A., D. J. Schingoethe, A. A. AbuGhazaleh, A. R. Hippen and K. R. Kalscheur. 2006. Milk production and composition from cows fed small amounts of fish oil and extruded soybeans. *J. Dairy Sci.* 89:3972-3980.
- Wildman, E. E., G. M. Jones, P. E. Wagner, R. L. Boman, J. R. Troutt and T. N. Lesch. 1982. A dairy cow body condition scoring system and its relationship to selected production characteristics. *J. Dairy Sci.* 65:495-501.