



Growth Performances and Carcass Characteristics of Indigenous Lambs Fed Halophyte *Sporobolus virginicus* Grass Hay

S. A. Al-Shorepy*, G. A. Alhadrami and A. J. Al-Dakheel¹

Department of Aridland Agriculture, College of Food and agriculture, UAE University, Al-Ain, P.O. Box, 17555, UAE

ABSTRACT : The objective of the present study was to compare feed and water intakes, growth performance and carcass characteristics of indigenous lamb fed diets containing various levels of halophyte *Sporobolus* grass hay. Forty male and female lambs were randomly and equally allotted with 5 lambs of each sex per treatment to four dietary treatment groups, which were initially formulated to have 100, 70, 30 or 0.0% *Sporobolus* grass hay, as a source of forage replacement for conventional Rhodes grass commonly used in the region. The lambs receiving 0.0% *Sporobolus* grass hay (100% Rhodes grass hay) served as the control. Feed and water were offered *ad libitum*. Male lambs were slaughtered at the end of the feeding trials. The average daily feed intake was significantly ($p < 0.05$) higher for the animals fed different levels of *Sporobolus* grass hay than for the control animals. Feed conversion ratio (FCR), i.e., kg feed/kg BW, was similar in all treatment groups. Although lambs fed the diet with 70% *Sporobolus* grass hay had heavier carcass weights, the differences were not significant. In conclusion, growth performance or carcass characteristics of fattening indigenous lambs were not influenced by the inclusion of different levels of *Sporobolus* grass hay in the diet. Because of this, *Sporobolus* grass hay represents an alternative forage resource for sustaining small ruminant production in the saline coastal and sub-coastal areas of the world. (**Key Words :** Halophyte, *Sporobolus* Grass, Indigenous Lambs, Growth Performance, Carcass Characteristics)

INTRODUCTION

Sheep numbers in the United Arab Emirates (UAE) have been estimated at about one million out of a total livestock population of over three million head (Ministry of Environment and Water, 2005). In the UAE, indigenous sheep constitute about 30-35% of the total sheep population (Al-Shorepy, 2001). Conventional production of sheep in the UAE depends primarily on feeding Rhodes grass hay and concentrates. However, fresh water available for irrigated Rhodes grass forage is under intense demand, while plenty of saline water is available for producing certain forages. Salt-tolerant forages, especially grasses that could grow well under saline irrigation, would be potentially valuable alternative forage resources and could play a major role in sustaining livestock production (Glenn and O'Laery, 1985; Gihad, 1993; Gihad and El Shaer, 1994; Masters et al., 2007).

The general nutritional characteristics of salt-tolerant forages have been well defined and the nutritional composition varies by species, stage of maturity and salinity of the irrigation source (Alhadrami et al., 1998; Robinson et al., 2004). However, many researchers reported that animals fed salt tolerant forage might have lower appetite and growth rate than animals fed on conventional forages (Gihad and El Shaer, 1994; Glenn et al., 1994; Miyamoto et al., 1994; Alhadrami et al., 2004). Intensive livestock production, integrated with mixed feeding of formulated rations offers other opportunities for the use of salt tolerant forages (Swingle et al., 1996; Kraidees et al., 1998; Abouheif et al., 2000).

Recently, attention has been given to the possibility of growing halophytes as irrigated crops on a large scale (Glenn and Watson 1993; Miyamoto et al., 1994; Alhadrami et al., 2004; Al-Dakheel et al., 2006). *Sporobolus* grass (*Sporobolus virginicus*), well known for its high salt tolerance, has been grown as irrigated forage under high salinity condition in the UAE (Alhadrami et al., 2004; Al-Dakheel et al., 2006). Unlike other salt-tolerant plants which tend to accumulate a high content of salt in their leaves, *Sporobolus* grass excludes salt at the root level

* Corresponding Author: S. A. Al-Shorepy. Tel: +971-3-7133442, Fax: +971-3-7632384, E-mail: salih.abdu@uaeu.ac.ae

¹ International Center for Biosaline Agriculture (ICBA), P. O. Box 14660, Dubai, UAE.

Received January 29, 2009; Accepted October 15, 2009

(Gallagher, 1985; Glenn, 1987). This characteristic would make Sporobolus grass an attractive alternative forage for hay production in the saline coastal and sub-coastal areas of the world. Several studies were conducted in the UAE University to evaluate the effect of feeding salt-tolerant grass on the performance of livestock such as camels, sheep and goats (Alhadrami, 2003; Alhadrami et al., 2004; Al-Shorepy, 2005). It was found that the performance of these animals was not affected by replacing Rhodes grass hay with Sporobolus grass hay either partially or completely. Thus, the objective of the present study was to assess the effects of inclusion of Sporobolus grass hay as a source of forage into whole-mixed diets on growth, feed and voluntary water intake and carcass composition of indigenous lambs.

MATERIALS AND METHODS

Experimental animals, feeds and management

This study was carried out at the College of Food and Agriculture Experimental Station, UAE University, located in Al Foah area, 20 km north of Al Ain City. A total of 20 male and 20 female indigenous lambs, weighing an average of 15.7 and 12.7 kg respectively were used to evaluate the effect of dietary inclusion of Sporobolus grass hay on growth performance and carcass characteristics. The animals were treated against internal and external parasites before commencement of the experiment. Five lambs of each sex were randomly and equally allotted to each of four dietary treatment groups, which were initially formulated to have 100, 70, 30 or 0.0% Sporobolus grass hay, as a source of forage replacement for the conventional Rhodes grass commonly used in the region. The group receiving 0.0 Sporobolus grass hay (100% Rhodes grass hay) served as the control. Sporobolus grass was grown in saline desert lands and irrigated with groundwater of high salt content (20,000 ppm). After cutting, the Sporobolus grass was sun dried till moisture content reached about 15%. The hay was baled into rectangular bales with an average weight of 11 kg

Table 1. Chemical composition of Rhodes and Sporobolus grass (DM basis)

Item	Rhodes grass	Sporobolus grass
Crude protein	10.10	7.28
Acid detergent fiber	35.90	32.93
Neutral detergent fiber	73.21	72.48
Ash	10.70	13.45

per bale. Rhodes grass hay was obtained from local commercial sources. The chemical composition of both Rhodes grass and Sporobolus grass is presented in Table 1. The diet was formulated using chopped Sporobolus grass hay and/or Rhodes grass hay and commercial concentrate to provide a ratio of 40:60 forage and concentrate, respectively. The lambs were fed in groups of the same sex. The composition of the diets offered as a total mixed ration is presented in Table 2.

Animals were subjected to the experimental diets for an adaptation period of 7 days before the commencement of the experiment. The amount of feed offered each time to the lambs was recorded daily. Refusals and wastage were weighed on the following morning. The total weight of the feed offered, adjusted for refusal and wastage, provided an estimate of the average feed consumed by each group per day throughout the study. The animals had free access to water and the volume of water offered each time was recorded daily and that remaining was recorded on the following morning. Daily water intake, corrected for losses, provided an estimate of the average water intake per day for each group. Animals were weighed on a weekly basis before offering fresh feed allowance. Samples of feed were collected biweekly for proximate analyses (AOAC, 1990).

Slaughtering procedures

At the end of a 63-day period, the male lambs (20) were slaughtered according to Islamic tradition. The lambs were fasted for at least 12 h and body weight (BW) was recorded before slaughter. After slaughtering, head, skin, feet and

Table 2. Ingredients of the experimental diets

Item	Treatment (% Sporobolus)			
	0.0	30	70	100
Ingredient (%)				
Sporobolus grass hay	0.0	13.6	26.4	40.0
Rhodes grass hay	40.0	26.4	13.6	0.0
Barley grain	28.0	28.0	28.0	28.0
Wheat Bran	17.0	17.0	17.0	17.0
Soya bean meal	13.0	13.0	13.0	13.0
Di-calcium	1.0	1.0	1.0	1.0
Vitamin and minerals	0.5	0.5	0.5	0.5
Trace mineralized salt	0.5	0.5	0.5	0.5
Crude protein of the diet (on DM Basis)	13.8	13.4	13.2	13.1

offal were removed and weighed. The reticulo-rumen was weighed both full and empty and gut content was calculated as the difference between the two weights. Empty body weight (EBW) was computed by subtracting the weight of digesta from live weight. The weights of other components of offal or non-carcass parts such as kidney and pelvic fat and different organs (liver, spleen, heart, kidneys, lung and trachea, diaphragm and testes) were recorded. The weight of the non-carcass components was expressed as a percentage of EBW. The dressing percentage was calculated as follows: dressing percentage = carcass weight/weight at slaughter; real dressing percentage = carcass weight/EBW. The carcass was then split longitudinally into two halves following the dorsal mid-line. From the left side, the 9th, 10th and 11th ribs were cut laterally to the vertebral column and parallel to the rib and subcutaneous fat thickness over *M. longissimus dorsi* (LD) muscle was measured with a caliper. At the same point, LD area was drawn on acetate paper and measured with a planimeter.

Measurements

The following measurements were recorded or calculated: average daily gain (ADG, g/d) was calculated as (final BW-initial BW/days on feed); intake was calculated on a DM basis (DMI, g/d); feed conversion ratio (FCR) was calculated by dividing DMI/weight gain; water

consumption per unit feed intake (l/kg) was calculated as (TWI/DMI).

Statistical analysis

Data were statistically analyzed using the General Linear Model procedures of SAS (1999). The linear model included effects of dietary treatment, sex of the lamb and their interaction in a 4×2 factorial design. A linear model that included only the effect of dietary treatment was used to analyze slaughtering characteristics for males. Significant differences among treatment means for dietary effect on growth and carcass parameters were analyzed using the least significant difference (LSD) method (Steel and Torrie, 1986). Significance was declared at $p < 0.05$.

RESULTS AND DISCUSSION

Feed and water intakes

Feed and water intakes for the indigenous male and female lambs are presented in Table 3. The average daily feed intake per animal for both males and females was significantly higher ($p < 0.05$) for animals fed the diet containing 100% *Sporobolus* grass hay than for those fed either 30 or 0.0% *Sporobolus* grass hay. On average, animals fed 100% *Sporobolus* grass hay consumed 13% more than animals fed the control diet (100% Rhodes grass

Table 3. Least square means of feed and water intakes in indigenous lambs fed different levels of *Sporobolus* grass hay

Sporobolus (%)	Sex	Daily feed intake (kg)	Daily water intake (L)	Water intake/feed (L/kg)
0.0	Male	0.94 ^b	2.31 ^d	2.48 ^{bc}
30.0	Male	0.98 ^b	3.35 ^b	3.44 ^a
70.0	Male	1.10 ^a	3.72 ^a	3.52 ^a
100.0	Male	1.10 ^a	2.65 ^c	2.617 ^b
0.0	Female	0.72 ^c	2.03 ^b	2.84 ^c
30.0	Female	0.79 ^b	2.66 ^a	3.38 ^b
70.0	Female	0.74 ^c	2.74 ^a	3.79 ^a
100.0	Female	0.83 ^a	2.62 ^a	3.191 ^b
Pooled SEM		0.01	0.05	0.07
Main effect				
Sex		Male	1.02 ^a	3.01 ^a
		Female	0.77 ^b	2.52 ^b
Pooled SEM		0.01	0.04	0.06
Diet type		0.0% <i>Sporobolus</i>	0.83 ^c	2.17 ^d
		30.0% <i>Sporobolus</i>	0.88 ^b	3.007 ^b
		70.0% <i>Sporobolus</i>	0.91 ^{ab}	3.24 ^a
		100.0% <i>Sporobolus</i>	0.94 ^a	2.64 ^c
Pooled SEM		0.01	0.06	0.07
Probabilities				
Sex		0.001	0.001	0.001
Diet type		0.001	0.001	0.001
Diet type×sex		0.001	0.001	0.03

Values in columns within each subcategory with different superscripts are significantly different ($p < 0.05$).

hay). Daily water consumption per animal was significantly higher ($p < 0.05$) for animals receiving the diet containing 70% Sporobolus grass hay compared to the control treatment. The volume of water intake per unit of feed was higher ($p < 0.05$) for animals fed 70% Sporobolus grass hay than for the other groups.

Similar results have been reported by Alhadrami et al. (2004), who observed higher feed intake for ewe lambs fed a diet containing 100% Sporobolus grass hay. Swingle et al. (1996), reported higher dry matter intake for lambs fed a diet containing halophyte forages than for lambs fed the control diet (Bermuda grass). Also, Swingle et al. (1996) found that lambs fed diets containing halophyte forages consumed up to 110% more water per day and 50% more water per kilogram of dry matter intake than those fed diets containing Bermuda grass. The increased water consumption of the lambs fed diets containing Sporobolus grass hay probably reflected a higher amount of sodium in the diet. The increased voluntary water intake is most likely due to the homeostatic requirement to excrete excess sodium (Reffett and Boling, 1985).

Growth rate and feed conversion ratio

Growth rate of indigenous lambs is presented in Table 4. Both initial and final weights and average daily gains of




Figure 1. Growth rate of indigenous lambs fed different levels of Sporobolus grass hay.

animals were not significantly different between groups. However, lambs fed diets containing Sporobolus grass hay tended to gain weight faster than those on the control treatment (100% Rhodes grass hay). The rate of growth for all diets was linear throughout the study (Figure 1).

In general, live-weight growth of indigenous lambs in the current study was comparable with better-known tropical breeds at similar age and given a traditional forage (Gatenby, 1986; Singh et al., 2003). Swingle et al. (1996) found that inclusion of halophyte forages supported the same weight gain of lambs as *cynodon* (Bermuda grass) hay.

Table 4. Least square means of growth rate and feed efficiency in indigenous lambs fed different levels of Sporobolus grass

Sporobolus %	Sex	Initial body weight (kg)	Final body weight (kg)	Average daily gain (g)	Feed conversion ratio
0.0	Male	15.4	24.7	147.6	6.5
30	Male	15.3	24.8	150.8	6.7
70	Male	16.4	27.2	171.4	6.4
100	Male	16.1	25.1	142.8	7.8
0.0	Female	12.7	18.7	95.2	8.1
30	Female	13.7	20.4	106.3	7.6
70	Female	12.1	18.7	104.8	7.2
100	Female	12.5	19.6	112.7	9.3
Pooled SEM		0.34	0.37	2.36	0.84
Main effect					
Sex					
Male		15.8	25.4 ^a	153.2 ^a	6.86
Female		12.7	19.3 ^b	104.8 ^b	8.06
Pooled SEM		0.78	0.88	5.81	0.57
Diet type					
0.0% Sporobolus		14.1	21.7	121.4	6.9
30% Sporobolus		14.5	22.6	128.6	6.9
70% Sporobolus		14.2	22.9	138.1	6.6
100% Sporobolus		14.3	22.3	127.8	7.5
Pooled SEM		1.25	1.63	11.51	0.82
Probabilities					
Sex		0.62	0.001	0.001	0.16
Diet type		0.98	0.90	0.57	0.49
Diet type×sex		0.85	0.71	0.46	0.96

Values in columns within each subcategory with different superscripts are significantly different ($p < 0.05$).

Kraidees et al. (1998) reported that Najdi lambs fed diets containing *Salicornia* stems tended to gain weight faster than those fed diets containing Rhodes grass hay. It was assumed that the improvement in body weight gain might be related to increased body-water retention and accumulation of sodium (Kraidees et al., 1998; Masters et al., 2005).

Feed conversion ratio (FCR; kg feed/kg BW) was lower for animals fed 70% *Sporobolus* grass hay than in the other three treatment groups. Though not significant, lambs fed a diet containing 100% *Sporobolus* grass hay had the highest FCR. FCR in the current study was similar to tropical (Al-Dabeeb, 2005) and temperate (Butterfield, 1988) sheep. The values of FCR obtained in the current study for animals fed diets containing *Sporobolus* grass hay were comparable to those reported for similar breeds. Mahgoub et al. (2000) recorded a FCR of 8.47 and 7.3 in sheep fed medium and high energy diets, respectively.

Carcass characteristics

Ram lambs fed various levels of *Sporobolus* grass hay and slaughtered at the end of the experiment at about 145

days of age had similar weights at slaughter and for hot carcass and empty body weights (EBW) (Table 5). Though not significant, lambs fed 70% *Sporobolus* grass hay tended to have higher slaughter weight compared with those lambs which received 0% *Sporobolus* grass hay (100% Rhode grass hay). Lambs fed 100% *Sporobolus* grass hay had significantly ($p < 0.05$) lower digestive tract contents than those fed either 0 or 30% *Sporobolus* grass hay. Expressed as % EBW, there were no differences in non-carcass components between lambs fed different levels of *Sporobolus* grass hay. Both physical and chemical composition of ribs were not influenced by the inclusion of different levels of *Sporobolus* grass hay in the diets.

Very little literature is available concerning effects of feeding *Sporobolus* grass hay on carcass characteristics. However, Swingle et al. (1996) concluded that inclusion of halophyte forages did not affect the carcass value of all experimental lambs. Kraidees et al. (1998) found that replacing Rhodes grass by *Salicornia* stems in the diet did not have any adverse effect on carcass characteristics of growing Najdi lambs if fresh water was available. The results found in this study were comparable to those

Table 5. Least squares means for proportion of non-carcass components and carcass measurements for indigenous lambs fed different levels of *Sporobolus* grass hay

	Treatment (% <i>Sporobolus</i>)				Pooled SEM	p
	0.0	30	70	100		
Slaughter weight (kg)	24.9	25.2	27.3	25.2	2.11	0.84
Empty body Weight ^a (kg)	20.32	21.0	22.6	21.3	1.96	0.86
Hot carcass weight (kg)	10.6	10.6	11.5	10.7	1.11	0.92
Hot dressing percentage (%)	42.2	41.9	42.0	41.9		
Digestive tract content (kg)	4.6 ^a	4.2 ^{ac}	4.7 ^a	3.9 ^{bc}	0.25	0.16
Proportion (%) on EBW ^b						
Hot carcass	51.8	50.5	50.8	49.8	0.99	0.56
Skin and limbs	14.8	14.4	14.9	14.3	0.55	0.86
Heart, lungs and liver	3.9	3.9	3.9	3.8	0.21	0.95
Fat (total)	7.2	6.5	8.2	7.0	0.84	0.52
Full digestive tract	32.9	30.4	30.3	28.1	2.12	0.47
Empty digestive tract	9.9	9.7	9.3	9.1	0.54	0.72
Head	7.3	7.1	7.0	7.4	0.21	0.62
Physical dissection of rib cut						
Rib weight (gm)	254.0	286.0	286.0	254.0	28.61	0.68
Rib area (cm)	9.7	9.3	9.4	11.1	0.89	0.48
Lean (%)	54.2	43.3	50.2	52.0	10.74	0.94
Fat (%)	21.7	21.8	22.7	20.0	6.10	0.98
Bone (%)	16.7	22.0	19.1	21.5	2.84	0.53
Chemical composition of <i>L. Dorsi</i> muscle (% DM)						
Protein	83.1	83.9	81.6	83.4	0.95	0.86
Ether extract	11.5	11.1	12.9	12.0	1.04	0.91
Ash	4.2	4.0	4.1	4.0	0.12	0.94

Values in columns within each subcategory with different superscripts are significantly different ($p < 0.05$).

^a Whole empty body = Live weight minus stomach and intestinal contents.

^b The weight of an organ or tissue relative to that of the whole empty body weight as a percentage.

reported for the Omani breed of sheep. Mahgoub and Early (2000) reported that hot carcass as a proportion of EBW ranged from 52-53%. Moreover, Mahgoub et al. (1998) found that slaughter weight of Omani lambs fed Rhodes grass hay at 162 days of age was 28.3 kg. Alhadrami et al. (1997) reported similar rib cut results for lambs of the same breed fed a diet containing 60% concentrate.

In the absence of mineral content of the *Sporobolus* hay, it appears that the rate of passage of digesta may have accelerated in lambs fed diets containing different levels of *Distichlis* grass hay. This implied a reduction of digesta-residue time in the rumen which might have increased the flow of dietary protein and soluble nutrients to the small intestine and resulted in increased absorption of nutrients of dietary origin, thereby accounting for the lower feed conversion efficiency and higher gain. This is consistent with the findings of Kraidees et al. (1998) who reported a higher gain for lambs fed *Salicornia* stems than on the control treatment. They further concluded that the changes in gain were presumably a direct consequence of higher mineral intake on the *Salicornia* diets. In addition, Kellaway et al. (1977) found that inclusion of Na up to 20 g/kg DM was well tolerated by calves; the growth rate was 44% greater at 20 g Na/kg DM in comparison to 2 g Na/kg DM. Moreover, Tomas et al. (1973) reported that, on mixed hay and concentrate diets, increased dilution rate in the rumen enhanced the efficiency of microbial protein synthesis. Findings of the current study indicated that diets based on 70% *Sporobolus* grass hay gave little advantage to experimental lambs over either Rhodes grass or other diets. This was reflected by higher daily gains, better feed conversion ratios (less feed per kg weight) and better slaughtering characteristics.

CONCLUSION

Findings of the present study suggest that inclusion of up to 100% *Sporobolus* grass hay in the diet did not have an adverse effect on growth performance or carcass characteristics of growing indigenous lambs. Therefore, *Sporobolus* grass could become an important forage resource for sustaining small ruminant production in the saline coastal and sub-coastal areas of the world. Although the water requirement of the animals is higher when feeding this grass, the net saving of fresh water is substantial.

ACKNOWLEDGMENT

This work was financially supported by the Research Affairs at the UAE University under a contract no. 07-6-12/02. Special thanks are due to Adil Alawad and Taha Aljuburi for their valuable assistance.

REFERENCES

- Abouheif, M. A., M. Al-Saiady, M. Kraidees, A. Tag Eldin and H. Metwally. 2000. Influence of inclusion of *salicornia* biomass in diets for rams on digestion and mineral balance. *Asian-Aust. J. Anim. Sci.* 13:967-973.
- Al-Dakheel, A. J., G. S. Al-Hadrami, S. A. Al-Shorepy and G. AbuRummani. 2006. Optimizing management practices for maximum production of two salt-tolerant grasses: *Sporobolus Virginicus* and *Distichlis Spicata*. *Proc. Seventh Annual UAE University Res. Conf.* 2:44-50.
- Al-Dabeeb, S. N. 2005. Effect of feeding low quality date palm on growth performance and apparent digestion coefficients in fattening Najdi sheep. *Small Rumin. Res.* 57:37-41.
- Alhadrami, G. A. 2003. *In situ* dry matter and fiber degradation of salt tolerant *Sporobolus* grass hay in camels fed yeast culture. *J. Camel Practice and Res.* 10(2):139-144.
- Alhadrami, G. A., S. A. Al-Shoropy, M. A. Ayoub and A. M. Yousef. 2004. Growth performance of Sheep fed *Sporobolus* grass hay grown in saline desert lands and irrigated with high salt content water. *Proc. Seventh Annual UAE University Res. Conf.* 2:27-34.
- Alhadrami, G. A., A. A. Nigm, A. M. Kholif and O. M. Abdalla. 1997. Effect of roughage to concentrate ratio on performance and carcass characteristics of local lambs in the United Arab Emirates. *Arab Gulf J. Sci. Res.* 15:137-148.
- Alhadrami, G. A., M. H. Abdel-Gawad and J. Jumma. 1998. *In-situ* SM and fiber degradation in camels and *in-vitro* gas production of two grasses irrigated with sea water. *Third Annual Meeting for Animal Production under arid Conditions*
- Al-Shorepy, S. A. 2001. Genetic parameters for growth traits of Local breed of sheep in the United Arab Emirates. *J. Agric. Sci. (Camb.)* 137:1-7.
- Al-Shoropy, S. A., G. A. Alhadrami and M. A. Ayoub. 2005. Growth performance indigenous goats fed *Distichlis* grass hay grown in saline desert lands and irrigated with high salt content water. *Proc. Seventh Annual UAE University Res. Conf.* 2:21-26.
- AOAC. 1990. *Official Methods of Analysis*, 15th Ed. Association of Official Analytical Chemist, Arlington, VA.
- Butterfield, R. M. 1988. *New concepts of sheep growth*. Department of Veterinary Anatomy. University of Sydney, Australia, p. 168.
- Gallagher, J. L. 1985. Halophytic crops for cultivation at seawater salinity. *Plant Soil* 89:323-336.
- Gatenby, R. M. 1986. *Sheep production in the tropics and the sub-tropics*. Longman, New York, USA, p. 351.
- Gihad, E. A. and H. M. El Shaer. 1994. Utilization of halophytes by livestock on rangelands: problems and prospects. In: *Halophytes as a resource for livestock and for rehabilitation of degraded lands* (Ed. V. R. Squires and A. T. Ayoub) pp. 77-96. Kluwer Academic, Dordrecht, Netherlands.
- Gihad, E. A. 1993. Utilization of high salinity tolerant plants and saline water by desert animals. In: *Towards the rational use of high salinity tolerant plants*, Vol. 1. (Ed. H. Lieth and A. A. Al Masoom). pp. 443-447. KluwerAcademic, Dordrecht, Netherlands.
- Glenn, E. P. 1987. Relationship between cation accumulation and water content of salt-tolerant grasses and a sedge. *Plant Cell*

- Environ. 10:205-212.
- Glenn, E. P. and J. W. O'Leary. 1985. Productivity and irrigation requirements of halophytes grown with seawater in the Sonoran Desert. *J. Arid Environ.* 9:81-91.
- Glenn, E. P. and M. C. Watson. 1993. Halophyte crops for direct seawater irrigation. In: *Towards the rational use of high salinity tolerant plants*. Vol. 1. (Ed. H. Lieth and A. A. Al Masoom). pp. 379-386. Kluwer Academic, Dordrecht, Netherlands.
- Glenn, E. P., R. S. Swingle, J. J. Riley, C. U. Mota, M. C. Watson, and V. R. Squires. 1994. North American halophytes: potential use in animal husbandry. In: *Halophytes as a resource for livestock and for rehabilitation of degraded lands* (Ed. V. R. Squires and A. T. Ayoub). pp. 165-174. Kluwer Academic, Dordrecht, Netherlands.
- Kellaway, R. C., D. J. Thomson, D. E. Beever and D. F. Osbourn. 1977. Effects of NaCl growth rate and acid±base balance in calves. *J. Agric. Sci. Camb.* 88:1-9.
- Kraidees, M. S., M. A. Abouheif, M. Y. Al-Saiady, A. Tag-Eldin, and H. Metwally. 1998. The effect of dietary inclusion of halophyte *Salicornia bigelovii* Torr on growth performance and carcass characteristics of lambs. *Anim. Feed Sci. Technol.* 76: 149-159.
- Mahgoub, O., D. J. Byerley, J. M. Chestworth and R. M. Myhara. 1998. Effect of testicular status and feeding diets containing palm by-products on growth and composition of the rack cut in Omani sheep. *Small Rumin. Res.* 28:281-288.
- Mahgoub, O., C. D. Lu and R. J. Early. 2000. Effects of dietary energy density on feed intake, body weight gain and carcass chemical composition of Omani growing lambs. *Small Rumin. Res.* 37:35-42.
- Masters, D. G., A. J. Rintoul, R. A. Dynes, K. L. Pearce and H. C. Norman. 2005. Feed intake and production in sheep fed diets high in sodium and potassium. *Aust. J. Agric. Res.* 56:427-434.
- Masters, D. G., S. E. Benes and H. C. Norman. 2007. Biosaline agriculture for forage and livestock production. *Agric. Ecosyst. Environ.* 119:234-248.
- Ministry of Environment and Water. 2005. *Agricultural census Results*, Ministry of Environment and Water, Dubai, United Arab Emirates.
- Miyamoto, S., E. P. Glenn and N. T. Singh. 1994. Utilization of halophytic plants for fodder production with brackish water in subtropical deserts. In: *Halophytes as a resource for livestock and for rehabilitation of degraded lands* (Ed. V. R. Squires and A. T. Ayoub). pp. 43-76. Kluwer Academic, Dordrecht, Netherlands.
- Reffett, J. K. and J. A. Boling. 1985. Nutrient utilization in lambs fed diets high in sodium or potassium. *J. Anim. Sci.* 61:1004-1009.
- Robinson, P. H., S. R. Grattan, G. Getachew, C. M. Grieve, J. A. Poss, D. L. Suarez and S. E. Benes. 2004. Biomass accumulation and potential nutritive value of some forages irrigated with saline-sodic drainage water. *Anim. Feed Sci. Technol.* 111:175-189.
- SAS. 1999. *Statistical Analysis System*. In: Release 9. 1. Institute INC, Cary, N.C., USA.
- Singh, N. P., S. K. Sankhyan and V. S. S. Prasad. 2003. Production performance and carcass characteristics of Malpura and Mutton synthetic lambs fed low and high energy rations in a semiarid region of India. *Asian-Aust. J. Anim. Sci.* 16(5):655-659.
- Steel, G. D. R. and H. J. Torrie. 1986. *Principles and Procedures of Statistics*, second ed. McGraw-Hill, USA.
- Swingle, R. S., E. P. Glenn and V. Squires. 1996. Growth performance of lambs fed mixed diets containing halophyte ingredients. *Anim. Feed Sci. Technol.* 63:137-148.
- Tomas, F. M., G. B. Jones, B. J. Potter and G. L. Langsford. 1973. Influence of saline drinking water on mineral balances in sheep. *Aust. J. Agric. Res.* 24:377-386.