



## Intermediary Metabolism of Plasma Acetic Acid, Glucose and Protein in Sheep Fed a Rice Straw-based Diet

M. K. Alam, Y. Ogata, Y. Sako, M. Al-Mamun and H. Sano\*

Iwate University, Ueda 3-18-8, Morioka, 020-8550, Japan

**ABSTRACT :** The present study was conducted to determine plasma acetate, glucose and protein metabolism using dilution of isotopes  $[1-^{13}\text{C}]$ Na acetate,  $[U-^{13}\text{C}]$ glucose and  $[1-^{13}\text{C}]$ leucine (Leu) in sheep fed rice straw (*Oriza japonica* L.). Four sheep were assigned to either rice straw (RS-diet) or mixed hay (MH-diet) with a crossover design. Nitrogen (N) intake and N digestibility were lower ( $p = 0.002$  and  $p = 0.02$ , respectively) for RS-diet than MH-diet, but N retention did not differ ( $p > 0.10$ ) between the diets. Concentrations of rumen acetate tended to be lower ( $p = 0.07$ ), and propionate was higher ( $p = 0.02$ ) for RS-diet than MH-diet. Concentrations of plasma lactate, non-esterified fatty acids, Leu and  $\alpha$ -ketoisocaproic acid did not differ ( $p > 0.10$ ) between the diets, but plasma glucose and urea concentrations were lower ( $p = 0.01$  and  $p = 0.003$ , respectively) for RS-diet than MH-diet. Turnover rate of plasma acetate did not differ ( $p = 0.39$ ) between the diets, and plasma glucose and Leu turnover rates were numerically lower ( $p = 0.15$  and  $p = 0.14$ , respectively) for RS-diet than MH-diet. Whole body protein synthesis and degradation did not differ ( $p > 0.10$ ) between the diets. Thus it can be concluded that the intermediary metabolism of acetate, glucose and protein on rice straw is comparable to mixed hay in sheep. (**Key Words :** Rice Straw, Intermediary Metabolism, Mixed Hay, Stable Isotope, Sheep)

### INTRODUCTION

Agricultural by-product is one of the most important feed resources in sustainable animal production. Rice is the world's second largest cereal crop after wheat, and produces the largest amount of crop residue at about 330 million tons (Van Soest, 2006). Abundant availability of rice straw makes it important in animal production. Rice straw is used as animal feed in many countries of the world, but it is widely used in the dry summer of developing countries to raise livestock production. In south and south-east Asian countries, the potential use of rice straw as animal feed is particularly important as it constitutes the staple diet of ruminants. Despite the large quantities of rice straw available in rice-producing countries, it is not extensively used as animal feed due to lack of information of its effect on ruminant production. Since rice straw with low nutritive value and low digestibility is the basal feed for ruminants (Hossain et al., 2002), the nutritive value of rice straw and its effect on digestion attributes in small and large

ruminants has been investigated (Acorda et al., 1992; Wu et al., 2005). Intermediary metabolism, particularly important for growth, lactation and production of ruminants, is influenced by dietary energy intake (Harris et al., 1992). However, information on intermediary metabolism in ruminants fed rice straw only is scanty. Therefore, it is necessary to know the effect of rice straw on intermediary metabolism of plasma nutrients in ruminants. The present study was designed to evaluate the effect of rice straw on the intermediary metabolism of plasma acetate, glucose and protein with simultaneous use of  $[1-^{13}\text{C}]$ Na acetate,  $[U-^{13}\text{C}]$ glucose and  $[1-^{13}\text{C}]$ leucine (Leu) isotopes as well as on digestion attributes such as nitrogen (N) balance and rumen characteristics in sheep fed rice straw and mixed hay.

### MATERIALS AND METHODS

#### Animals, diets and management

Four crossbred (Corriedale $\times$ Suffolk) shorn sheep (*Ovis aries* L.), average age 3 yr and body weight (BW)  $50 \pm 1$  kg, were used in this experiment. The sheep were assigned to two dietary treatments; one rice straw (*Oriza japonica* L.) only diet (RS-diet) and the other a mixed hay diet (MH-diet) of orchardgrass (*Dactylis glomerata* L.) and reed canarygrass (*Phalaris arundinaceae* L.) at a 60:40 ratio

\* Corresponding Author: Hiroaki Sano. Department of Animal Sciences, Faculty of Agriculture, Iwate University, Ueda 3-18-8, Morioka, 020-8550, Japan. Tel: +81-19-621-6165, Fax: +81-19-621-6165, E-mail: sano@iwate-u.ac.jp

Received March 3, 2010; Accepted June 11, 2010

**Table 1.** Chemical composition of feed

Item (%)	MH	RS
Dry matter	94.3	94.4
Crude protein	13.5	4.4
Crude ash	10.9	15.5
Neutral detergent fiber	69.7	73.8

MH = Mixed hay of orchardgrass (*Dactylis glomerata* L.) and reed canarygrass (*Phalaris arundinacea* L.).

RS = Rice straw (*Oriza japonica* L.).

(Table 1). The metabolizable energy (ME) was estimated at 1.30 kcal/g for rice straw (NARO, 2006) and 1.73 kcal/g for mixed hay (NRC, 1985). In the preliminary experiment 20% of the RS-diet remained as leftover and the MH-diet was completely consumed by the sheep when both diets were given at maintenance level. For this reason, feed allowance was 67.2 g/kg<sup>0.75</sup>/d for RS-diet based on energy at maintenance level and 40.5 g/kg<sup>0.75</sup>/d for MH-diet based on energy about 20% less than maintenance level to ensure similar energy intake for both diets. The experiment was performed using a crossover design with two 21-d periods. Two sheep were fed RS-diet during the first period and then fed MH-diet during the second period, and the other two sheep were fed these diets in the reverse order. The sheep were housed in individual pens in an animal barn during the first 14 d of each dietary period and were fed twice daily at 8:00 h and 20:00 h. Water was available *ad libitum*. On day 15, the sheep were moved to individual metabolism cages in a controlled environment chamber with an air temperature of 23±1°C and lighting from 7:00 h to 21:00 h. The sheep were weighed on days 1, 8, 15 and 21 of each dietary period. The handling of animals, including cannulation and blood sampling, was carried out according to the rules and regulations established by the Animal Care Committee of Iwate University.

#### Nitrogen balance test and collection of rumen fluid

A nitrogen balance trial was conducted for 5 days from d 16 to d 20 for each dietary treatment. Urine was collected from each sheep every 24 h in a bucket containing 50 ml of 6 N H<sub>2</sub>SO<sub>4</sub> and the volume was recorded. A sub-sample of urine was stored at -30°C until analysis. Feces were also collected from each sheep every 24 h and dried at 60°C in a forced air oven for 48 h and then weighed after being held at room temperature for 5 days. The required amount of feces was ground to pass through a 1 mm screen and stored at room temperature for further analysis. On d 20, rumen fluid was collected before feeding (BF), 3 h (3F) and 6 h (6F) after feeding via a stomach tube for measuring pH, ammonia-N (NH<sub>3</sub>-N) and volatile fatty acids (VFA). The pH of rumen fluid was measured immediately after collection with a pH meter (HM-10P, Toa Electronics Ltd., Japan). A sub-sample was centrifuged at 8,000×g for 10 min

at 0°C (RS-18IV, Tomy, Japan), then 1 ml of supernatant was taken and mixed with 1 ml of 0.1 N HCl and stored at -30°C until measurement of rumen NH<sub>3</sub>-N concentration. Residual rumen fluid samples were preserved at -30°C for further analysis.

#### Isotope dilution method

For determining the turnover rates (TR) of plasma acetate, glucose and Leu the isotope dilution methods using [1-<sup>13</sup>C]Na acetate, [U-<sup>13</sup>C]glucose and [1-<sup>13</sup>C]Leu were conducted simultaneously on d 21 of each dietary period. Two catheters, one for isotope infusion and another for blood sampling, were inserted into the left and right jugular veins on the morning of the study. The catheters were filled with sterile solution of tri-sodium citrate (0.13 mol/L) and at 12:00 h 87 μmol/kg<sup>0.75</sup> of [1-<sup>13</sup>C]Na acetate (1-<sup>13</sup>C, 99%, Cambridge Isotope Laboratories, Inc., USA), 3.1 μmol/kg<sup>0.75</sup> of [U-<sup>13</sup>C]glucose (D-glucose -<sup>13</sup>C<sub>6</sub>, 99 atom % excess <sup>13</sup>C; Cambridge Isotope Laboratories, USA) and 7.2 μmol/kg<sup>0.75</sup> of [1-<sup>13</sup>C]Leu (L-leucine-1-<sup>13</sup>C, 99 atom % excess <sup>13</sup>C; Cambridge Isotope Laboratories, USA) dissolved in saline solution were injected as a priming dose through the jugular infusion catheter. Immediately after the injection of priming dose, the isotopes were continuously infused at rates of 87, 3.1 and 7.2 μmol/kg<sup>0.75</sup>/h of [1-<sup>13</sup>C]Na acetate, [U-<sup>13</sup>C]glucose and [1-<sup>13</sup>C]Leu, respectively, by a multichannel peristaltic pump (AC-2120, Atto Co. Ltd., Japan) for 4 h through the same catheter. Blood samples (5 ml) were collected through the sampling catheter just before the priming dose injection and at 30 min intervals during the last 2 h of the isotope infusion. The collected blood samples were immediately transferred to heparinized tubes, stored in crushed ice until centrifugation at 10,000×g for 10 min at 2°C, and the plasma samples then stored at -30°C for further analysis.

#### Chemical analysis

Analyses of proximate composition of the experimental diets were performed using the methods described in AOAC (1995). Nitrogen in diets, feces, urine and feed residues was analyzed by the Kjeldahl method with the Foss Kjeltex System (Tecator Digester System and Kjeltex 2300, Foss Tecator, Sweden). Rumen VFA concentrations were determined by titrating the steam distillate of rumen fluid with 0.1 N NaOH. The titrated distillate was dried and then individual VFA were determined using gas chromatography (5890A, Hewlett Packard Co., USA). Ammonia-N content of rumen fluid was determined using a colorimetric method (Wetherburn, 1967).

Plasma [1-<sup>13</sup>C]acetate enrichment and concentrations of plasma VFA and lactate were determined using selected ion monitoring with a gas chromatography-mass spectrometry system (GC/MS) (QP-2010, Shimadzu, Japan), according to

the procedure of Moreau et al. (2003) as previously described by Al-Mamun et al. (2009). Plasma [ $U-^{13}C$ ]glucose enrichment was determined by the procedure of Tserng and Kalhan (1983) with slight modifications as described previously (Sano et al., 1996). The enrichment of [ $U-^{13}C$ ]glucose was determined using selected ion monitoring with GC/MS. Concentrations of plasma glucose were determined using the method described by Huggett and Nixon (1957). Plasma amino acids and  $\alpha$ -keto acids were separated and converted to N-methyl-N-t-butyl-dimethylsilyltrifluoroacetamide (MTBSTFA) derivatives according to the procedures of Rocchiccioli et al. (1981) and Calder and Smith (1988), as described previously (Sano et al., 2004). Isotopic enrichments of plasma [ $1-^{13}C$ ]Leu and  $\alpha$ -[ $1-^{13}C$ ]ketoisocaproic acid [ $\alpha$ -[ $1-^{13}C$ ]KIC] and concentrations of plasma Leu and  $\alpha$ -KIC were measured by selected ion monitoring using the GC/MS. Concentrations of plasma non-esterified fatty acids (NEFA) and urea were determined using kits (NEFA C and Urea NB, Wako Pure Chemicals, Japan).

### Calculation

For the isotope dilution method, the TR of plasma acetate, glucose and Leu were calculated using the equation given by Tserng and Kalhan (1983).

$$TR = I \times (1/E - 1)$$

Where, TR is the turnover rate of plasma acetate, glucose and Leu, I is the infusion rate of [ $1-^{13}C$ ]Na acetate, [ $U-^{13}C$ ]glucose and [ $1-^{13}C$ ]Leu and E is the plasma isotope enrichment of [ $1-^{13}C$ ]acetate, [ $U-^{13}C$ ]glucose and [ $1-^{13}C$ ]Leu or  $\alpha$ -[ $1-^{13}C$ ]KIC at steady state.

Whole body protein synthesis (WBPS) and degradation (WBPD) were calculated from the relationships between whole body protein flux (WBPF), N absorption and urinary N excretion according to the equations described by

Schroeder et al. (2006), as follows:

$$WBPS = WBPF - (\text{urinary N excretion} \times 6.25)$$

$$WBPD = WBPF - (N \text{ absorbed} \times 6.25)$$

Leucine concentrations in carcass protein (66 g/kg) were used as described by Harris et al. (1992). Thus the WBPF was obtained by dividing the turnover rate of plasma Leu by 0.066.

### Statistical analysis

All data were statistically analyzed with the MIXED procedure of SAS (1996). The least-squares means statement was used to test the effects of diet and period and the random effect was sheep. Results were considered significant at  $p < 0.05$  and a tendency was defined as  $0.05 \leq p < 0.10$ . The repeated measures statement and the Tukey adjustment were used for the time course of changes and the significance level was  $p < 0.05$ .

## RESULTS

Body weight change did not differ ( $p = 0.31$ ) between the diets. Dry matter intake (DMI) and estimated ME intake were higher ( $p = 0.002$  and  $p = 0.03$ , respectively) for RS-diet than MH-diet. Nitrogen intake, N excretion through urine and feces and N digestibility were lower ( $p < 0.05$ ) for RS-diet than MH-diet, but N retention did not differ ( $p = 0.39$ ) between the dietary treatments (Table 2).

Rumen pH was not affected ( $p = 0.26$ ) by the diets, and decreased ( $p < 0.05$ ) after feeding (Table 3). Rumen  $NH_3$ -N concentration was lower ( $p = 0.0002$ ) for RS-diet than MH-diet and decreased at 6F ( $p < 0.05$ ). The concentrations of rumen total VFA did not differ ( $p = 0.17$ ) between the diets and also did not differ ( $p = 0.28$ ) after feeding. The concentration of acetic acid tended to be lower ( $p = 0.07$ ),

**Table 2.** Effect dietary intake on body weight change, dry matter intake, estimated metabolizable energy (ME) intake, nitrogen (N) balance and digestibility of N in sheep

Item	MH-diet	RS-diet	SEM	p-value
No. of sheep	4	4		
Body weight change (kg/d)	-0.11	-0.07	0.04	0.31
Dry matter intake (g/kg <sup>0.75</sup> /d)	37	54	5	0.002
Estimated ME intake (kcal/kg <sup>0.75</sup> /d)	63	70	3	0.03
N intake (g/kg <sup>0.75</sup> /d)	0.85	0.47	0.12	0.002
N in faeces (g/kg <sup>0.75</sup> /d)	0.28	0.21	0.02	0.01
N in urine (g/kg <sup>0.75</sup> /d)	0.42	0.09	0.10	0.002
N absorption (g/kg <sup>0.75</sup> /d)	0.57	0.26	0.10	0.004
N retention (g/kg <sup>0.75</sup> /d)	0.15	0.17	0.02	0.39
N digestibility (%)	67	56	4	0.02

MH = Mixed hay of orchardgrass (*Dactylis glomerata* L.) and reed canarygrass (*Phalaris arundinacea* L.), RS = Rice straw (*Oriza japonica* L.), SEM = Standard error of means.

**Table 3.** Dietary effects on rumen pH, concentrations of rumen ammonia-N and volatile fatty acids (VFA) before feeding (BF) and at 3h (3F) and 6h (6F) after feeding in sheep

Rumen parameters	Treatments						SEM	p-value		
	MH-diet			RS-diet				Diet	Time	Diet×Time
	BF	3F	6F	BF	3F	6F				
No. of sheep	4	4	4	4	4	4				
pH	6.9	6.8	6.7	7.0	6.8	6.9	0.10	0.26	0.04	0.04
NH <sub>3</sub> -N (mg/dL)	8.4	9.6	7.4	2.3	1.5	1.0	2.2	0.0002	0.001	0.001
Total VFA (mmol/L)	91.8	92.5	98.3	88.0	89.9	89.2	2.1	0.17	0.28	0.56
Acetate (mmol/L)	69.7	69.8	73.8	62.9	59.5	62.2	3.2	0.07	0.40	0.84
Propionate (mmol/L)	14.9	14.6	17.1	18.1	23.2	19.7	1.9	0.02	0.26	0.10
iso-Butyrate (mmol/L)	0.9	0.9	0.7	0.5	0.4	0.5	0.1	0.04	0.18	0.44
Butyrate (mmol/L)	4.8	4.7	5.6	5.5	6.0	5.9	0.32	0.11	0.78	0.49
iso-Valerate (mmol/L)	1.2	1.6	0.7	0.6	0.4	0.5	0.3	0.08	0.29	0.22
Valerate (mmol/L)	0.4	1.0	0.4	0.4	0.4	0.4	0.1	0.20	0.13	0.25

MH = Mixed hay of orchardgrass (*Dactylis glomerata* L.) and reed canarygrass (*Phalaris arundinacea* L.),

RS = Rice straw (*Oriza japonica* L.), SEM = Standard error of means.

and the concentration of propionate was higher ( $p = 0.02$ ) for RS-diet than MH-diet.

Concentrations of plasma glucose and urea were lower ( $p = 0.01$  and  $p = 0.003$ , respectively) for RS-diet than MH-diet (Table 4). Concentrations of plasma acetate, Leu, NEFA and lactate did not differ ( $p > 0.10$ ) between the dietary treatments.

Plasma acetate concentration and enrichment of plasma [ $1-^{13}\text{C}$ ]acetate remained constant during the last 2 h period of the [ $1-^{13}\text{C}$ ]Na acetate infusion (data are not shown). Turnover rate of plasma acetate did not differ ( $p = 0.39$ ) between the dietary treatments. Plasma glucose concentration and enrichment of plasma [ $U-^{13}\text{C}$ ]glucose remained constant during the latter half of the

**Table 4.** Dietary effects on plasma acetate, glucose and protein metabolism and concentration of plasma non-esterified fatty acids (NEFA), urea and lactate in sheep

Item	MH-diet	RS-diet	SEM	p-value
No. of sheep	4	4		
Acetate				
Concentration (mmol/L)	0.363	0.410	0.047	0.48
TR (mmol/kg <sup>0.75</sup> /h)	4.92	4.12	0.86	0.39
Glucose				
Concentration (mmol/L)	3.54	3.20	0.11	0.01
TR (mmol/kg <sup>0.75</sup> /h)	1.58	1.37	0.15	0.15
Leu				
Concentration (μmol/L)	90.8	73.9	8.4	0.21
TR (mmol/kg <sup>0.75</sup> /h)	0.259	0.205	0.027	0.14
WBPS (g/kg <sup>0.75</sup> /h)	9.7	9.2	1.0	0.67
WBPD (g/kg <sup>0.75</sup> /h)	8.8	8.1	1.0	0.58
α-KIC				
Concentration (μmol/L)	15.3	13.8	2.1	0.54
TR (mmol/kg <sup>0.75</sup> /h)	0.338	0.250	0.039	0.16
WBPS (g/kg <sup>0.75</sup> /h)	13.5	11.3	1.5	0.39
WBPD (g/kg <sup>0.75</sup> /h)	12.5	10.3	1.5	0.37
NEFA concentration (mEq/L)	0.25	0.22	0.04	0.41
Urea concentration (mmol/L)	3.3	1.1	0.7	0.003
Lactate concentration (mmol/L)	0.352	0.404	0.066	0.16

MH = Mixed hay of orchardgrass (*Dactylis glomerata* L.) and reed canarygrass (*Phalaris arundinacea* L.), RS = Rice straw (*Oriza japonica* L.),

SEM = Standard error of means, TR = Turnover rate, Leu = Leucine, WBPS = Whole body protein synthesis, WBPD = Whole body protein degradation,

α-KIC = α-ketoisocaproic acid.

[U-<sup>13</sup>C]glucose infusion (data are not shown). Turnover rate of plasma glucose was not affected ( $p = 0.15$ ) by the dietary treatments. Concentrations of plasma Leu and  $\alpha$ -KIC and enrichments of plasma [1-<sup>13</sup>C]Leu and  $\alpha$ -[1-<sup>13</sup>C]KIC remained constant during the latter 2 h of the [1-<sup>13</sup>C]Leu infusion (data are not shown). Plasma LeuTR calculated from [1-<sup>13</sup>C]Leu and  $\alpha$ -[1-<sup>13</sup>C]KIC enrichment was numerically lower ( $p = 0.14$  and  $p = 0.16$ , respectively) for RS-diet than MH-diet, but WBPS and WBPD did not differ ( $p > 0.10$ ) between the diets.

## DISCUSSION

The present experiment demonstrated that the effect of rice straw on intermediary metabolism of acetate, glucose and protein in sheep could be comparable to mixed hay. In the present study N digestibility and N excretion through urine was lower for RS-diet than MH-diet. This might be due to lower dietary CP intake for RS-diet, because similar results were previously found in lactating cows (Castillo et al., 2001) and sheep (Sano et al., 2004; Al-Mamun et al., 2008). The lower urinary N excretion for RS-diet, which suggested an increased urea recycling or a decreased protein oxidation due to lower CP intake than the MH-diet, is in agreement with Al-Mamun et al. (2008). Although N intake and urinary N excretion were lower for the RS-diet than MH-diet, N balance remained similar between the diets. This may suggest that the RS-diet contained more rumen undegradable protein than the MH-diet, and this result is in accordance with Al-Mamun et al. (2008).

In the present study, rumen pH was not affected by diets, but declined after feeding both diets. This drop in pH may be associated with the fermentation of carbohydrate and similar production of total VFA in the rumen. This is in agreement with the findings of Salman et al. (2008) who suggested that pH values were inversely related to total VFA concentration in the rumen of goats. The same trend was found in sheep by Santoso et al. (2006). The lower NH<sub>3</sub>-N concentration in the rumen for RS-diet than MH-diet might be due to the lower dietary CP intake, because similar results were found in sheep (Al-Mamun et al., 2008). Although NH<sub>3</sub>-N concentration was considerably lower for RS-diet than MH-diet, rumen pH did not differ between the diets in the present study.

Energy intake is inversely related to plasma NEFA concentrations (Sticker et al., 1995) and plasma NEFA is the best indicator of body lipid loss (Chillard et al., 2000). In the present study, unchanged plasma NEFA concentration for both diets might be due to use of roughage diets with restricted energy. Restricted energy intake for both diets caused similar mobilization of fatty acids from adipose tissue, which was responsible for similar BW loss of sheep

on both diets. Lower plasma glucose concentration was found for RS-diet than MH-diet. Evans et al. (1974) also found a lower glucose concentration for a low quality than a high quality roughage diet.

The central role of rumen fermentation products in intermediary metabolism of ruminants is well recognized. In the present study, plasma acetate TR remained similar for both diets. This might be due to use of roughage diets which fermented equally in the rumen being responsible for similar plasma acetate concentration and TR. Prior (1978) determined plasma acetate TR in sheep fed restricted and *ad libitum* amounts of feed and stated that plasma acetate concentrations and the apparent turnover rate of acetate were not significantly influenced by level of feed intake. Plasma acetate TR in the present study was comparable with previous data as determined by [<sup>14</sup>C]acetate dilution technique in sheep fed lucerne hay plus a concentrate mixture (Pethick and Lindsay, 1982; Sunagawa et al., 1986).

In adult sheep, plasma glucose TR was correlated with dietary energy intake level, suggesting that the nutritional status of the animal had at least as much influence as the supply of glucose precursors (Ortigue-Marty et al., 2003). The estimated energy intake level might also influence the dynamics of glucose metabolism as shown in lactating cows (Konig et al., 1984). In the present study, dietary treatments had no significant effect on plasma glucose TR, although propionate, a major glucose precursor in the ruminant, was higher for RS-diet than MH-diet. This might be due to roughage diets resulting in lower absorption of propionate, because Sano et al. (1999) suggested that, in sheep fed a roughage diet, propionate absorption is not strongly increased by feeding and gluconeogenesis is sustained by variable contributions of different precursors over the feeding cycle. Rodriguez et al. (1985) also reported that propionate infusion into the rumen failed to influence percentage of glucose derived from propionate, amount of propionate converted to glucose, and glucose TR in lactating goats fed a forage-based diet with concentrate mixture. The numerical value of plasma glucose TR in the present findings was comparable with data previously reported by Sano et al. (1999) in which plasma glucose TR was determined in sheep fed roughage diets at restricted and *ad libitum* amounts.

Plasma LeuTR in the present study was comparable to data reported previously in sheep (Sano et al., 2004). In the present study, numerically lower plasma LeuTR for RS-diet than MH-diet might be due to lower intake of CP, since dietary CP intake is positively correlated with LeuTR in sheep (Al-Mamun et al., 2008). However, Sano et al. (2004) reported that plasma LeuTR in sheep was influenced only marginally by dietary CP intake when ME intake was constant.

In the present study, even though plasma LeuTR differed slightly, WBPS and WBPD calculated from the enrichments of plasma [1-<sup>13</sup>C]Leu and  $\alpha$ -[1-<sup>13</sup>C]KIC did not differ between the diets. In the equation used (Schroeder et al., 2006), lower urinary N excretion and lower N absorption result in higher protein synthesis (WBPS) and degradation (WBPD). In spite of different CP intake, WBPS and WBPD did not differ between the diets, which might be due to use of the above method of calculation. The values of WBPS and WBPD in this study were calculated from [1-<sup>13</sup>C]Leu and  $\alpha$ -[1-<sup>13</sup>C]KIC, which were very similar to previous values calculated in sheep (Al-Mamun et al., 2007) using the same equation.

No significant differences were found in plasma acetate, glucose and protein metabolism in sheep between the dietary treatments. It can be suggested that the performance of RS-diet was comparable to MH-diet in relation to intermediary metabolism of plasma acetate, glucose and protein in sheep and hence rice straw could serve as an alternative feed for ruminants.

## REFERENCES

- Acorda, J. A., M. Okamoto and N. Yoshida. 1992. Nutritive value of rice straw processed with sodium hydroxide, soybean meal, urea, cage layer manure and molasses. *J. Jpn. Soc. Grassl. Sci.* 37:405-411.
- Al-Mamun, M., K. Goto, S. Chiba and H. Sano. 2009. Responses of plasma acetate metabolism to hop (*Humulus lupulus* L.) in sheep. *Int. J. Biol. Sci.* 5:287-292.
- Al-Mamun, M., Y. Hanai, C. Tanaka, Y. Tamura and H. Sano. 2008. Responses of whole body protein synthesis and degradation to plantain herb in sheep exposed to heat. *Arch. Anim. Nutr.* 62:219-229.
- Al-Mamun, M., C. Ito, A. Sato, T. Fujita and H. Sano. 2007. Comparison of the [<sup>2</sup>H<sub>5</sub>]phenylalanine model with [1-<sup>13</sup>C]leucine method to determine whole body protein synthesis and degradation in sheep fed two levels. *Asian-Aust. J. Anim. Sci.* 20:1517-1524.
- AOAC. 1995. Official methods of analysis. 16<sup>th</sup> edn. Association of Official Analytical Chemists, Arlington, Virginia.
- Calder, A. G. and A. Smith. 1988. Stable isotope ratio analysis of leucine and ketoisocaproic acid in blood plasma by gas chromatography/mass spectrometry. Use of tertiary butyldimethylsilyl derivatives. *Rapid Commun. Mass Spectrom.* 2:14-16.
- Castillo, A. R., E. Kebreab, D. E. Beever, J. H. Barbi, J. D. Sutton, H. C. Kirby and J. France. 2001. The effect of protein supplementation on nitrogen utilization in lactating dairy cows fed grass silage diets. *J. Anim. Sci.* 79:247-253.
- Chilliard, Y., A. Feraly, Y. Faulconnier, M. Bonnet, J. Rouel and F. Bocquier. 2000. Adipose tissue metabolism and its role in adaptations to undernutrition in ruminants. *Proc. Nutr. Soc.* 59:127-134.
- Evans, E., J. G. Buchanan-Smith, G. K. MacLeod and J. B. Stone. 1974. Glucose metabolism in cows fed low and high-roughage diets. *J. Dairy Sci.* 58:672-677.
- Fujita, T., M. Kajita and H. Sano. 2006. Responses of whole body protein synthesis, nitrogen retention and glucose kinetics to supplemental starch in goats. *Comp. Biochem. Physiol. B.* 144:180-187.
- Fujita, T., M. Kajita and H. Sano. 2007. Effects of non protein energy intake on whole body protein synthesis, nitrogen retention and glucose turnover in goats. *Asian-Aust. J. Anim. Sci.* 20:536-542.
- Harris, P. M., P. A. Skene, V. Buchan, E. Milne, A. G. Calder, S. E. Anderson, A. Connell and G. E. Lobley. 1992. Effect of food intake on hind-limb and whole body protein metabolism in young growing sheep: Chronic studies based on arterio-venous techniques. *Br. J. Nutr.* 68:389-407.
- Hossain, M. S., M. N. Haque, S. A. Aziz, M. S. Mazumder, M. L. Ali, and A. T. M. M-E-Elahi. 2002. Effect of urea molasses straw on the productive and reproductive performance of indigenous cows under the village conditions of Bangladesh. *Pakistan J. Biol. Sci.* 9:997-999.
- Huggett, A. G. and D. A. Nixon. 1957. Enzymatic determination of blood glucose. *Biochem. J.* 66:12.
- Krishnamurti, C. R. and S. M. Janssens. 1988. Determination of leucine metabolism and protein turnover in sheep, using gas-liquid chromatography-mass spectrometry. *Br. J. Nutr.* 59:155-164.
- Konig, B. A., J. D. Oldham and D. S. Parker. 1984. The effect of abomasal infusion of casein on acetate, palmitate and glucose kinetics in cows during early lactation. *Br. J. Nutr.* 52:319-328.
- Lapierre, H., J. P. Blouin, J. F. Bernier, C. K. Reynolds, P. Dubreuil and G. E. Lobley. 2002. Effect of supply of metabolizable protein on whole body and splanchnic leucine metabolism in lactating cows. *J. Dairy Sci.* 48:193-198.
- Moreau, N. M., S. M. Goupy, J. P. Antiganac, F. J. Monteau, B. J. Le Biezec, M. M. Champ, L. J. Martin and H. J. Dumon. 2003. Simultaneous measurement of plasma concentrations and <sup>13</sup>C-enrichment of short-chain fatty acids, lactic acid and ketone bodies by gas chromatography coupled to mass spectrometry. *J. Chromatogr. B.* 784:395-403.
- National Agriculture and Food Research Organization. 2006. Japanese feeding standard for dairy cattle. Japan Livestock Industry Association.
- National Research Council. 1985. Nutrient requirements of sheep. 6<sup>th</sup> Ed. National Academy Press. Washington, DC.
- Ortigue-Marty, I., J. Vernet and L. Majdoub. 2003. Whole body glucose turnover in growing and non-productive adult ruminants: meta-analysis and review. *J. Reprod. Nutr. Dev.* 43:371-383.
- Pethick, D. W. and D. B. Lindsay. 1982. Acetate metabolism in lactating sheep. *Br. J. Nutr.* 48:319-328.
- Prior, R. L. 1978. Effect of level of feed intake on lactate and acetate metabolism and lipogenesis *in vivo* in sheep. *J. Nutr.* 108:926-935.
- Rocchiccioli, F., J. P. Leroux and P. Cartier. 1981. Quantitation of 2-ketoacids in biological fluids by gas chromatography chemical ionization mass spectrometry of o-trimethylsilylquinoxalinol derivatives. *Biomed. Mass Spectrom.* 8:160-164.

- Rodriguez, N. R., E. C. Prigge, D. C. Lough and W. H. Hoover. 1985. Glucogenic and hormonal responses to abomasal casein and ruminal volatile fatty acid infusions in lactating goats. *J. Dairy Sci.* 68:1968-1975.
- Salman, F. M., R. I. El-Kadi, H. Abdel-Rahman, S. M. Ahmed, M. I. Mohammad and M. M. Shoukry. 2008. Biologically treated sugar beet pulp as a supplement in goat rations. *Int. J. Agric. Biol.* 10:412-416.
- Sano, H. and T. Fujita. 2006. Effect of supplemental calcium propionate on insulin action to blood glucose metabolism in adult sheep. *Reprod. Nutr. Dev.* 46:9-18.
- Sano, H., T. Fujita, M. Murakami and A. Shiga. 1996. Stimulative effect of epinephrine on glucose production and utilization rates in sheep using a stable isotope. *Domest. Anim. Endocrinol.* 13:445-451.
- Sano, H., M. Kajita and T. Fujita. 2004. Effect of dietary protein intake on plasma leucine flux, protein synthesis, and degradation in sheep. *Comp. Biochem. Physiol. B.* 139:163-168.
- Sano, H., D. N. Mowat, R. O. Ball and D. R. Trout. 1997. Effect of supplemental chromium on whole-body kinetics of glucose, lactate and propionate in rams fed a high grain diet. *Comp. Biochem. Physiol. B.* 118:117-121.
- Sano, H., H. Sawada, A. Takenami and M. Al-Mamun. 2009. Effects of diet and cold exposure on rates of plasma leucine turnover and protein synthesis in sheep. *J. Agric. Sci. (Camb.)* 147:91-97.
- Sano, H., A. Takebayashi, Y. Kodama, K. Nakamura, H. Ito, Y. Arino, T. Fujita, H. Takahashi and K. Ambo. 1999. Effects of feed restriction and cold exposure on glucose metabolism in response to feeding and insulin in sheep. *J. Anim. Sci.* 77:2564-2573.
- Sano, H., Y. Tamura and A. Shiga. 2002. Metabolism and glucose kinetics in sheep fed plantain and orchard grass and exposed to cold. *NZ. J. Agric. Res.* 45:171-177.
- Santoso, B., B. Mwenya, C. Sar and J. Takahashi. 2006. Ruminal fermentation and nitrogen metabolism in sheep fed a silage-based diet supplemented with *Yucca schidigera* and nisin. *Anim. Feed Sci. Technol.* 129:187-195.
- SAS. 1996. SAS/STAT® Software: Changes and Enhancements through Release 6.11. SAS Inst. Inc. Cary, NC.
- Schroeder, G. F., E. C. Titgemeyer, M. S. Awadeh, J. S. Smith and D. P. Gnad. 2006. Effects of energy level on methionine utilization by growing steers. *J. Anim. Sci.* 84:1497-1504.
- Sticker, L. S., D. L. Thompson, L. D. Bunting, J. M. Fernandez and C. L. DePew. 1995. Dietary protein and (or) energy restriction in mares: plasma glucose, insulin, nonesterified fatty acid and urea nitrogen responses to feeding, glucose and epinephrine. *J. Anim. Sci.* 73:136-144.
- Sunagawa, K., F. Otani, A. Hagino, K. Takahashi, Y. Otomo, Y. Shoji, K. Ambo and T. Tsuda. 1986. Kinetics of exogenous and endogenous acetate metabolism in sheep exposed to cold environment. *Jpn. J. Zootech. Sci.* 57:201-208.
- Sutton, J. D., M. S. Dhanoa, S. V. Morant, J. France, D. J. Napper and E. Schuller. 2003. Rates of production of acetate, propionate and butyrate in the rumen of lactating dairy cows given normal and low roughage diets. *J. Dairy Sci.* 86:3620-3633.
- Tserng, K. Y. and S. C. Kalhan. 1983. Calculation of substrate turnover rate in stable isotope tracer studies. *Am. J. Physiol.* 245:E308-E311.
- Van Soest, P. J. 2006. Rice straw, the role of silica and treatments to improve quality. *J. Anim. Feed Sci. Technol.* 130:137-171.
- Weatherburn, M. W. 1967. Phenol-hypochlorite reaction for determination of ammonia. *J. Anal. Chem.* 39:971-974.
- Wu, Y., W. Hu and J. Liu. 2005. Effects of supplementary urea-mineral lick block on the kinetics of fiber digestion, nutrient digestibility and nitrogen utilization of low quality roughages. *J. Zhejiang Univ. Sci.* 8:793-797.